



HAL
open science

Classical and Effective Descriptive Complexities of omega-Powers

Olivier Finkel, Dominique Lecomte

► **To cite this version:**

Olivier Finkel, Dominique Lecomte. Classical and Effective Descriptive Complexities of omega-Powers. 2007. ensl-00168644v1

HAL Id: ensl-00168644

<https://ens-lyon.hal.science/ensl-00168644v1>

Preprint submitted on 29 Aug 2007 (v1), last revised 4 Aug 2009 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Classical and effective descriptive complexities of ω -powers *

Olivier Finkel

Equipe Modèles de Calcul et Complexité
Laboratoire de l'Informatique du Parallélisme[†]
 CNRS et Ecole Normale Supérieure de Lyon
 46, Allée d'Italie 69364 Lyon Cedex 07, France.
 Olivier.Finkel@ens-lyon.fr

Dominique Lecomte

Equipe d'Analyse Fonctionnelle
 Université Paris 6, tour 46-0, boîte 186,
 4, place Jussieu, 75 252 Paris Cedex 05, France.
 lecomte@moka.ccr.jussieu.fr

and

Université de Picardie, I.U.T. de l'Oise, site de Creil,
 13, allée de la faïencerie, 60 107 Creil, France.

Abstract

We prove that, for each countable ordinal $\xi \geq 1$, there exist some Σ_ξ^0 -complete ω -powers, and some Π_ξ^0 -complete ω -powers, extending previous works on the topological complexity of ω -powers [Fin01, Fin03, Fin04, Lec01, Lec05, DF06]. We prove effective versions of these results; in particular, for each recursive ordinal $\xi < \omega_1^{CK}$ there exist some recursive sets $A \subseteq 2^{<\omega}$ such that $A^\infty \in \Pi_\xi^0 \setminus \Sigma_\xi^0$ (respectively, $A^\infty \in \Sigma_\xi^0 \setminus \Pi_\xi^0$), where Π_ξ^0 and Σ_ξ^0 denote classes of the hyperarithmetical hierarchy. To do this, we prove effective versions of a result by Kuratowski, describing a Π_ξ^0 set as the range of a closed subset of the Baire space ω^ω by a continuous bijection. This leads us to prove closure properties for the pointclasses Σ_ξ^0 in arbitrary recursively presented Polish spaces. We apply our existence results to get better computations of the topological complexity of some sets of dictionaries considered in [Lec05].

Keywords. ω -power, Borel class, complete, effective descriptive set theory, hyperarithmetical hierarchy.

2000 Mathematics Subject Classification. Primary: 03E15, 03B70, Secondary: 54H05, 68Q15, 68Q45, 68R15.

*LIP Research Report RR 2007-38

[†]UMR 5668 - CNRS - ENS Lyon - UCB Lyon - INRIA

This paper is an extended version of a conference paper which will appear in the Proceedings of the 16th EACSL Annual Conference on Computer Science and Logic, CSL 07 [FL07]. Part of the results in this paper have been also presented at the International Conference Computability in Europe, CiE 07, Siena, Italy, June 2007.

1 Introduction.

We consider the finite alphabet $\Sigma = \{0, \dots, \Sigma - 1\}$, where $\Sigma \geq 2$ is an integer, and a language over this alphabet, i.e., a subset A of the set $\Sigma^{<\omega}$ of finite words with letters in Σ . Notice that a language of finite words will be also sometimes called a dictionary, as in [Lec05]. The set of infinite words over the alphabet Σ , i.e., of sequences of length ω of letters of Σ , is denoted Σ^ω .

Definition 1.1 *The ω -power associated with A is the set A^∞ of the infinite sentences constructible with A by concatenation. So we have $A^\infty := \{ a_0 a_1 \dots \in \Sigma^\omega \mid \forall i \in \omega \ a_i \in A \}$.*

Notice that we denote here A^∞ the ω -power associated with A , as in [Lec05], while it is often denoted A^ω in Theoretical Computer Science papers, as in [Sta97a, Fin01, Fin03, FL07]. Here we reserved the notation A^ω to denote the cartesian product of countably many copies of A since this will be often used in this paper.

In the theory of formal languages of infinite words, accepted by various kinds of automata, the ω -powers appear very naturally in the characterization of the class REG_ω of ω -regular languages (respectively, of the class CF_ω of context free ω -languages) as the ω -Kleene closure of the family REG of regular finitary languages (respectively, of the family CF of context free finitary languages) [Sta97a].

Since the set Σ^ω of infinite words over a finite alphabet Σ can be equipped with the usual Cantor topology, the question of the topological complexity of ω -powers of finitary languages naturally arises and has been posed by Niwinski [Niw90], Simonnet [Sim92], and Staiger [Sta97a].

What are the possible levels of topological complexity for the ω -powers?

As the concatenation map, from A^ω onto A^∞ , which associates $a_0 a_1 \dots$ to $(a_i)_{i \in \omega}$, is continuous, an ω -power is always an analytic set.

It has been recently proved, that for each integer $n \geq 1$, there exist some ω -powers of (context-free) languages which are Π_n^0 -complete Borel sets, [Fin01], and that there exists a (context-free) language L such that L^ω is analytic but not Borel, [Fin03]. Amazingly, the language L is very simple to describe and it is accepted by a simple 1-counter automaton. Notice that Louveau has proved independently that analytic-complete ω -powers exist, but the existence was proved in a non effective way. We refer the reader to [ABB96] for basic notions about context-free languages.

The first author proved in [Fin04] that there exists a finitary language V such that V^ω is a Borel set of infinite rank. However the only known fact on their complexity is that there is a (context-free) language W such that W^ω is Borel above Δ_ω^0 , [DF06]. In particular, it was still unknown which could be the possible infinite Borel ranks of ω -powers.

The basic notions of descriptive set theory used in this paper will be recalled in the next section. We now state our results which extend the previous ones.

Theorem 1.2 (a) Let $3 \leq \xi < \omega_1$, and $\Gamma \neq \check{\Gamma}$ be a Wadge class closed under finite unions satisfying the inclusions $\Delta_\xi^0(\Gamma) \subseteq \Gamma = \Delta_3^0\text{-}PU(\Gamma) \subseteq \Pi_{\xi+1}^0$. Then there is $A \subseteq 2^{<\omega}$ such that A^∞ is Γ -complete.

(b) Let $1 \leq \xi < \omega_1$. Then there is $A \subseteq 2^{<\omega}$ such that A^∞ is Σ_ξ^0 -complete.

(c) Let $1 \leq \xi < \omega_1$. Then there is $A \subseteq 2^{<\omega}$ such that A^∞ is Π_ξ^0 -complete.

(d) Let $1 \leq \xi < \omega_1$. Then there is $A \subseteq 2^{<\omega}$ such that A^∞ is $\check{D}_2(\Sigma_\xi^0)$ -complete.

(e) Let $3 \leq \xi < \omega_1$ and $\omega \leq \eta < \omega_1$ be an indecomposable ordinal. Then there is $A \subseteq 2^{<\omega}$ such that A^∞ is $\check{D}_\eta(\Sigma_\xi^0)$ -complete.

So we get a complete knowledge of the Borel classes Γ for which there is $A \subseteq 2^{<\omega}$ such that A^∞ is Γ -complete. Indeed, the only class Δ_ξ^0 admitting a complete set is Δ_1^0 . And $A := \{s \in 2^{<\omega} \mid 0 \prec s \text{ or } 1^2 \prec s\}$ implies that $A^\infty = 2^\omega \setminus N_{10}$ is a Δ_1^0 -complete set.

In this context coming from theoretical computer science, it is natural to wonder whether these examples are effective. We answer positively. The reader should see [Mos80] for basic notions of effective descriptive set theory. It is known that $B \subseteq 2^\omega$ is Σ_ξ^0 -complete if and only if $B \in \Sigma_\xi^0 \setminus \Pi_\xi^0$ (see 22.10 in [Kec95]). The effective version of Theorem 1.2 is the following:

Theorem 1.3 (1) Let $1 \leq \xi < \omega_1^{CK}$.

(a) There is $A \subseteq 2^{<\omega}$ such that $A^\infty \in \Sigma_\xi^0 \setminus \Pi_\xi^0$.

(b) There is $A \subseteq 2^{<\omega}$ such that $A^\infty \in \Pi_\xi^0 \setminus \Sigma_\xi^0$.

Moreover, A can be coded by a Δ_1^0 subset of ω .

(2) Similarly, let $\beta \in 2^\omega$ and $1 \leq \xi < \omega_1^\beta$.

(a) There is $A \subseteq 2^{<\omega}$ such that $A^\infty \in \Sigma_\xi^0(\beta) \setminus \Pi_\xi^0$.

(b) There is $A \subseteq 2^{<\omega}$ such that $A^\infty \in \Pi_\xi^0(\beta) \setminus \Sigma_\xi^0$.

Moreover, A can be coded by a $\Delta_1^0(\beta)$ subset of ω .

To prove Theorem 1.2, we use a theorem of Kuratowski which is a level by level version of a theorem of Lusin and Souslin stating that every Borel set $B \subseteq 2^\omega$ is the image of a closed subset of the Baire space ω^ω by a continuous bijection. This theorem of Lusin and Souslin had already been used by Arnold in [Arn83] to prove that every Borel subset of Σ^ω , for a finite alphabet Σ , is accepted by a non-ambiguous finitely branching transition system with Büchi acceptance condition and our first idea was to code the behaviour of such a transition system. This way, in the general case, we can manage to construct an ω -power of the same complexity as B . We now state Kuratowski's Theorem [Kur66] (see Corollary 33.II.1):

Theorem 1.4 Let $\xi \geq 1$ be a countable ordinal, X a zero-dimensional Polish space, and $B \in \Pi_{\xi+1}^0(X)$. Then there is $C \in \Pi_1^0(\omega^\omega)$ and a continuous bijection $f: C \rightarrow B$ such that f^{-1} is Σ_ξ^0 -measurable (i.e., $f[U]$ is $\Sigma_\xi^0(B)$ for each open subset U of C).

To prove Theorem 1.3, we first prove an effective version of Theorem 1.4. It has the following consequence.

Theorem 1.5 *Let $\xi \geq 1$ be a countable ordinal, and $B \in \Pi_{\xi+1}^0(2^\omega)$. Then there is $C \in \Pi_1^0(\omega^\omega)$, a partial function $f : \omega^\omega \rightarrow 2^\omega$, recursive on C , and a partial function $g : 2^\omega \rightarrow \omega^\omega$, Σ_ξ^0 -recursive on B , such that f defines a bijection from C onto B and g coincides with f^{-1} .*

To prove Theorems 1.3 and 1.5, we prove some results of effective descriptive set theory that cannot be found in [Mos80]. We prove that the pointclasses Σ_ξ^0 are, uniformly and in the codes, closed under taking sections at points in spaces of type at most 1, substitutions of partial recursive functions, finite intersections and unions, \exists^ω , among other things.

In [Lec05], the following question is asked. What is the topological complexity of the set of dictionaries whose associated ω -power is of a given level of complexity? More specifically, let $1 \leq \xi < \omega_1$. The following $\Sigma_2^1(2^{2^{<\omega}}) \setminus D_2(\Sigma_1^0)$ sets are introduced:

$$\Sigma_\xi := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Sigma_\xi^0\},$$

$$\Pi_\xi := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Pi_\xi^0\},$$

$$\Delta := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Delta_1^1\} = \{A \subseteq 2^{<\omega} \mid A^\infty \in \Pi_1^1\}.$$

The proof of Theorem 1.3 gives some more informations about the complexity of these sets. We will prove, using a result by J. Saint Raymond, that Σ_ξ and Π_ξ are Π_1^1 -hard if $\xi \geq 3$, which is a much better approximation of their complexity than the one in [Lec05]. The proof of this fact has the following consequence. Theorem 1.2 shows that the ω -powers are quite general objects. On the other hand, we will prove another result showing that they are not arbitrary.

Notation. Let Γ be a class having a universal set $\mathcal{U}_\Gamma^{2^\omega} \subseteq (2^\omega)^2$, and Γ' another class. We set

$$\mathcal{U}(\Gamma, \Gamma') := \{\alpha \in 2^\omega \mid (\mathcal{U}_\Gamma^{2^\omega})_\alpha \in \Gamma'\}.$$

Let X, Y be zero-dimensional Polish spaces and $A \subseteq X$, $B \subseteq Y$. We will use the following notation to denote the Wadge quasi-order:

$$(X, A) \leq_W (Y, B) \Leftrightarrow \exists f : X \rightarrow Y \text{ continuous with } A = f^{-1}(B).$$

We write $(X, A) <_W (Y, B)$ if $(X, A) \leq_W (Y, B)$ and $(Y, B) \not\leq_W (X, A)$.

The consequence we mentioned is the following. If we choose suitable universal sets, then the following inequalities hold:

$$\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0) \leq_W \Sigma_\xi <_W \mathcal{U}(\Sigma_1^1, \Sigma_\xi^0)$$

$$\mathcal{U}(\Sigma_\xi^0, \Pi_\xi^0) \leq_W \Pi_\xi <_W \mathcal{U}(\Sigma_1^1, \Pi_\xi^0)$$

$$\mathcal{U}(\Sigma_1^1, \Pi_1^1) \not\leq_W \Delta <_W \mathcal{U}(\Sigma_1^1, \Delta_1^1) = \mathcal{U}(\Sigma_1^1, \Pi_1^1).$$

This means that the ω -powers are analytic sets that do not behave like arbitrary analytic sets. This also means that there is a strong difference between the Borel levels on one side, and the level of analytic sets on the other side. Actually, our method to prove Theorem 1.3 is a method that works for the Borel levels, and it cannot be extended to the level of analytic sets, even if Theorem 1.3 can be extended to the level of analytic sets (see [Fin03]). Note that we will prove that $\mathcal{U}(\Sigma_1^1, \Delta_1^1)$ is Π_2^1 -complete.

This paper is organized as follows:

- In section 2 we prove Theorem 1.2.
- In section 3 we recall a few basic facts of effective descriptive set theory, and fix some notation. Then we prove the results of effective descriptive set theory that we need for the sequel. This is where the closure properties for the pointclasses Σ_ξ^0 are proved.
- In section 4 we prove Theorem 1.5.
- In section 5 we prove Theorem 1.3.
- In section 6 we study the complexity of some sets of dictionaries.

2 Proof of Theorem 1.2.

Basic facts and notation.

In descriptive set theory, we study the topological complexity of definable subsets of Polish spaces, i.e., of separable and completely metrizable topological spaces.

- The notation for the *Borel classes* in metrizable spaces is as follows: Σ_1^0 is the class of open sets, and if $\xi \geq 1$ is a countable ordinal, then Π_ξ^0 is the class of complements of Σ_ξ^0 sets, Σ_ξ^0 is the class of countable unions of sets in $\bigcup_{1 \leq \eta < \xi} \Pi_\eta^0$, and Δ_ξ^0 is the class $\Sigma_\xi^0 \cap \Pi_\xi^0$. The class of *Borel sets* is

$$\Delta_1^1 := \bigcup_{1 \leq \xi < \omega_1} \Sigma_\xi^0 = \bigcup_{1 \leq \xi < \omega_1} \Pi_\xi^0.$$

- The class of *analytic sets* is the class Σ_1^1 of subsets of Polish spaces that are continuous images of Polish spaces. One can prove that if X is a Polish space, then $A \subseteq X$ is analytic if and only if A is the projection on X of a closed subset of $X \times \omega^\omega$ (see 14.3 in [Kec95]). Then we can define the *projective classes* in Polish spaces as follows; if $n \geq 1$ is an integer, then Π_n^1 is the class of complements of Σ_n^1 sets, $\Sigma_{n+1}^1(X)$ is the class of projections on X of sets in $\Pi_n^1(X \times \omega^\omega)$, and Δ_n^1 is the class $\Sigma_n^1 \cap \Pi_n^1$.
- If Γ is a class of sets in Polish spaces and X is a Polish space, then a set $\mathcal{U}_\Gamma^X \in \Gamma(2^\omega \times X)$ is *universal* for $\Gamma(X)$ if $\Gamma(X) = \{(\mathcal{U}_\Gamma^X)_\alpha \mid \alpha \in 2^\omega\}$ (where $(\mathcal{U}_\Gamma^X)_\alpha := \{x \in X \mid (\alpha, x) \in \mathcal{U}_\Gamma^X\}$). For example, there are universal sets for $\Sigma_\xi^0(X)$, $\Pi_\xi^0(X)$, $\Sigma_1^1(X)$, $\Pi_1^1(X)$ for any Polish space X (see 22.3 and 26.1 in [Kec95]).
- Recall that a Polish space is *zero-dimensional* if it has a basis consisting of Δ_1^0 sets. Typically, let K be a countable set. If K is equipped with the discrete topology and $s \in K^{<\omega}$, then $N_s := \{\alpha \in K^\omega \mid s \prec \alpha\}$ is a basic Δ_1^0 set of K^ω ($s \prec \alpha$ means that s is a beginning of α). The length of $\gamma \in K^{\leq \omega}$ is denoted $|\gamma|$. If $\gamma \in K^{\leq \omega}$ and $k \in \omega$, then $\gamma \upharpoonright k$ is the beginning of length k of γ . If $s \prec \alpha = \alpha(0)\alpha(1)\dots$, then $\alpha - s$ is the sequence $\alpha(|s|)\alpha(|s|+1)\dots$
- If Γ is a class of sets in zero-dimensional Polish spaces, closed under continuous preimages, then a subset A of X is *Γ -hard* if for each $A' \in \Gamma(X')$ there is a continuous map $f: X' \rightarrow X$ with $A' = f^{-1}(A)$.

If $A \in \Gamma(X)$ is Γ -hard, then we say that A is Γ -complete. We say that Γ is a Wadge class if there is a Γ -complete set. We denote $\check{\Gamma} := \{\neg A \mid A \in \Gamma\}$. If $\Gamma \neq \check{\Gamma} \subseteq \Delta_1^1$ is a Wadge class, then A is Γ -complete if and only if $A \in \Gamma \setminus \check{\Gamma}$.

- If Γ is a Wadge class, then $\Delta_\xi^0(\Gamma) \subseteq \Gamma$ means that $E \in \Gamma(X)$ if $E \in \Delta_\xi^0(A)$ and $A \in \Gamma(X)$.
- If I is a set and Γ is a class or a set, then $(x_i)_{i \in I} \subseteq \Gamma$ means that $x_i \in \Gamma$ for each $i \in I$.
- We set $\Delta_\xi^0\text{-PU}(\Gamma) := \{\bigcup_{n \in \omega} A_n \cap P_n \mid (A_n)_{n \in \omega} \subseteq \Gamma \text{ and } (P_n)_{n \in \omega} \subseteq \Delta_\xi^0 \text{ partition}\}$ if $1 \leq \xi < \omega_1$. One can prove that if $\Delta_1^0 \subseteq \Gamma \neq \check{\Gamma} \subseteq \Delta_1^1$ is a Wadge class, then there is a bigger $1 \leq \xi < \omega_1$ (the level of Γ) such that $\Gamma = \Delta_\xi^0\text{-PU}(\Gamma)$ (see [LStR88]).

If $\eta < \omega_1$ and $(A_\theta)_{\theta < \eta}$ is an increasing sequence of subsets of some space X , then we set

$$D_\eta[(A_\theta)_{\theta < \eta}] := \{x \in X \mid \exists \theta < \eta \ x \in A_\theta \setminus \bigcup_{\theta' < \theta} A_{\theta'} \text{ and the parity of } \theta \text{ is opposite to that of } \eta\}.$$

If moreover $1 \leq \xi < \omega_1$, then we set $D_\eta(\Sigma_\xi^0) := \{D_\eta[(A_\theta)_{\theta < \eta}] \mid (A_\theta)_{\theta < \eta} \subseteq \Sigma_\xi^0\}$. One can prove that $D_\eta(\Sigma_\xi^0)$ has level ξ if $\eta \geq 1$ (see [LStR88]).

- We say that $\omega \leq \eta < \omega_1$ is indecomposable if η cannot be represented as $\eta_1 + \eta_2$ with $\eta_1, \eta_2 < \eta$. It is known that the indecomposable ordinals are the ω^θ with $1 \leq \theta < \omega_1$ (see IV.2.16 in [Lev79]).

Proof of Theorem 1.2.

- We have already said that the existence of the continuous bijection $f : C \rightarrow B$ given by Lusin and Souslin's Theorem had already been used by Arnold in [Arn83] to prove that every Borel subset of Σ^ω , for a finite alphabet Σ , is accepted by a non-ambiguous finitely branching transition system with Büchi acceptance condition. We now recall the definition of these transition systems.

A Büchi transition system is a tuple $\mathcal{T} = (\Sigma, Q, \delta, q_0, Q_f)$, where Σ is a finite input alphabet, Q is a countable set of states, $\delta \subseteq Q \times \Sigma \times Q$ is the transition relation, $q_0 \in Q$ is the initial state, and $Q_f \subseteq Q$ is the set of final states. A run of \mathcal{T} over an infinite word $\sigma \in \Sigma^\omega$ is an infinite sequence of states $(t_i)_{i \geq 0}$, such that $t_0 = q_0$, and for each $i \geq 0$, $(t_i, \sigma(i), t_{i+1}) \in \delta$. The run is said to be accepting iff there are infinitely many integers i such that t_i is in Q_f .

The transition system is said to be non-ambiguous if each infinite word $\sigma \in \Sigma^\omega$ has at most one accepting run by \mathcal{T} .

The transition system is said to be finitely branching if for each state $q \in Q$ and each $a \in \Sigma$, there are only finitely many states q' such that $(q, a, q') \in \delta$.

Our first idea was to code the behaviour of such a transition system. In fact this can be done on a part of infinite words of a special compact set $K_{0,0}$. However we shall have also to consider more general sets $K_{N,j}$ and then we shall need the hypothesis of the Σ_ξ^0 -measurability of the function f , which is given by Kuratowski's Theorem.

- We now come to the proof of Theorem 1.2.

(a) We may assume that $\Delta_1^0 \subseteq \Gamma$, otherwise $\Gamma = \{\emptyset\}$ since $\Delta_\xi^0(\Gamma) \subseteq \Gamma$, in which case $A := \emptyset$ is suitable. This implies that $\Delta_\xi^0 \subseteq \Gamma$ since $\Delta_\xi^0(\Gamma) \subseteq \Gamma$.

- Let $B \in \Gamma(2^\omega) \setminus \check{\Gamma}$, and $P_\infty := \{\alpha \in 2^\omega \mid \forall m \in \omega \exists n \geq m \alpha(n) = 1\}$, which is homeomorphic to ω^ω (we associate $0^{\beta(0)} 10^{\beta(1)} 1\dots$ to $\beta \in \omega^\omega$). As $B \in \Pi_{\xi+1}^0$, Theorem 1.4 gives $C \in \Pi_1^0(P_\infty)$ and f . By Proposition 11 in [Lec05], it is enough to find $A \subseteq 4^{<\omega}$. The dictionary A will be made of two pieces: we will have $A = \mu \cup \pi$. The set π will code f , and π^∞ will look like B on some nice compact sets $K_{N,j}$. Outside a countable family of compact sets, we will hide f , so that A^∞ will be the simple set μ^∞ .

- We set $Q := \{(s, t) \in 2^{<\omega} \times 2^{<\omega} \mid |s| = |t|\}$. We enumerate Q as follows. We start with $q_0 := (\emptyset, \emptyset)$. Then we put the sequences of length 1 of elements of 2×2 , in the lexicographical ordering: $q_1 := (0, 0)$, $q_2 := (0, 1)$, $q_3 := (1, 0)$, $q_4 := (1, 1)$. Then we put the 16 sequences of length 2: $q_5 := (0^2, 0^2)$, $q_6 := (0^2, 01)$, \dots And so on. We will sometimes use the coordinates of $q_N := (q_N^0, q_N^1)$. We put $M_j := \sum_{i < j} 4^{i+1}$. Note that the sequence $(M_j)_{j \in \omega}$ is strictly increasing, and that q_{M_j} is the last sequence of length j of elements of 2×2 .

- If $l \in \omega$ and $(a_i)_{i < l} \in (\omega^{<\omega})^l$, then $\frown_{i < l} a_i$ is the concatenation $a_0 \dots a_{l-1}$. Similarly, $\frown_{i \in \omega} a_i$ is the concatenation $a_0 a_1 \dots$.

- Now we define the “nice compact sets”. We will sometimes view 2 as an alphabet, and sometimes view it as a letter. To make this distinction clear, we will use the boldface notation $\mathbf{2}$ for the letter, and the lightface notation 2 otherwise.

We will have the same distinction with 3 instead of 2, so that $2 = \{0, 1\}$, $3 = \{0, 1, \mathbf{2}\}$, $4 = \{0, 1, \mathbf{2}, \mathbf{3}\}$. Let N, j be non-negative integers with $N \leq M_j$. We set

$$K_{N,j} := \{ \gamma = \mathbf{2}^N \frown [\frown_{i \in \omega} m_i \mathbf{2}^{M_{j+i+1}} \mathbf{3} \mathbf{2}^{M_{j+i+1}}] \in 4^\omega \mid \forall i \in \omega \ m_i \in 2 \}.$$

As the map $\varphi_{N,j} : K_{N,j} \rightarrow 2^\omega$ defined by $\varphi_{N,j}(\gamma) := (m_i)_{i \in \omega}$ is a homeomorphism, $K_{N,j}$ is compact.

- Now we will define the sets that “look like B ”.

- We define a function $c : B \times \omega \rightarrow Q$ by $c(\alpha, l) := [f^{-1}(\alpha), \alpha][l]$. Note that Q is countable, so that we equip it with the discrete topology. In these conditions, we prove that c is Σ_ξ^0 -measurable.

For any $q \in Q$, it holds that $c^{-1}(\{q\}) = \{(\alpha, l) \in B \times \omega \mid f^{-1}(\alpha)[l] = q^0 \text{ and } \alpha[l] = q^1\}$. But $\alpha[l] = q^1$ means that “ $l = |q^1|$ and α belongs to the basic open set N_{q^1} ”. In the same fashion, $f^{-1}(\alpha)[l] = q^0$ means that “ $l = |q^0|$ and $f^{-1}(\alpha)$ belongs to the basic open set N_{q^0} ”, or equivalently that “ $l = |q^0|$ and $\alpha = f(f^{-1}(\alpha))$ belongs to $f[C \cap N_{q^0}]$ ”. As $f[C \cap N_{q^0}]$ is a Σ_ξ^0 subset of B , $c^{-1}(\{q\})$ is a Σ_ξ^0 subset of $B \times \omega$ and c is Σ_ξ^0 -measurable.

- Let N be an integer. We put

$$E_N := \{ \alpha \in 2^\omega \mid q_N^1 \alpha \in B \text{ and } c(q_N^1 \alpha, |q_N^1|) = q_N \}.$$

Note that $E_0 = \{ \alpha \in 2^\omega \mid \alpha \in B \text{ and } c(\alpha, 0) = \emptyset \} = B$. Let us prove that $E_N \in \Gamma(2^\omega)$ for each integer N .

As c is Σ_ξ^0 -measurable and $\{q_N\} \in \Delta_1^0(Q)$, we get $c^{-1}(\{q_N\}) \in \Delta_\xi^0(B \times \omega)$. Note that the map $S: \{\alpha \in 2^\omega \mid q_N^1 \alpha \in B\} \rightarrow B \times \omega$ defined by $S(\alpha) := (q_N^1 \alpha, |q_N^1|)$ is continuous, so that $E_N = S^{-1}[c^{-1}(\{q_N\})]$ is in $\Delta_\xi^0(\{\alpha \in 2^\omega \mid q_N^1 \alpha \in B\})$. As $B \in \Gamma(2^\omega)$ and the map $\alpha \mapsto q_N^1 \alpha$ is continuous, $\{\alpha \in 2^\omega \mid q_N^1 \alpha \in B\}$ is in $\Gamma(2^\omega)$. Thus $E_N \in \Gamma(2^\omega)$ since $\Delta_\xi^0(\Gamma) \subseteq \Gamma$.

Now we define the transition system obtained from f .

- If $m \in 2$ and $n, p \in \omega$, then we write $n \xrightarrow{m} p$ if $q_n^0 \prec q_p^0$ and $q_p^1 = q_n^1 m$.

- As f is continuous on C , the graph $\text{Gr}(f)$ of f is a closed subset of $C \times 2^\omega$. As C is $\Pi_1^0(P_\infty)$, $\text{Gr}(f)$ is also a closed subset of $P_\infty \times 2^\omega$. So there is a closed subset F of $2^\omega \times 2^\omega$ such that $\text{Gr}(f) = F \cap (P_\infty \times 2^\omega)$. We identify $2^\omega \times 2^\omega$ with $(2 \times 2)^\omega$, i.e., we view (β, α) as $[\beta(0), \alpha(0)], [\beta(1), \alpha(1)], \dots$. By Proposition 2.4 in [Kec95], there is $R \subseteq (2 \times 2)^{<\omega}$, closed under initial segments, such that $F = \{(\beta, \alpha) \in 2^\omega \times 2^\omega \mid \forall k \in \omega (\beta, \alpha) \upharpoonright k \in R\}$. Notice that R is a tree whose infinite branches form the set F . In particular, we get

$$(\beta, \alpha) \in \text{Gr}(f) \Leftrightarrow \beta \in P_\infty \text{ and } \forall k \in \omega (\beta, \alpha) \upharpoonright k \in R.$$

- We set $Q_f := \{(t, s) \in R \mid t \neq \emptyset \text{ and } t(|t| - 1) = 1\}$. Notice that Q_f is simply the set of pairs $(t, s) \in R$ such that the last letter of t is a 1.

We have in fact already defined the transition system \mathcal{T} obtained from f . This transition system has a countably infinite set Q of states and a set Q_f of accepting states. The initial state is $q_0 := (\emptyset, \emptyset)$. The input alphabet is $2 = \{0, 1\}$ and the transition relation $\delta \subseteq Q \times 2 \times Q$ is given by: if $m \in 2$ and $n, p \in \omega$ then $(q_n, m, q_p) \in \delta$ iff $n \xrightarrow{m} p$.

Recall that a run of \mathcal{T} is said to be Büchi accepting if final states occur infinitely often during this run. Then the set of ω -words over the alphabet 2 which are accepted by the transition system \mathcal{T} from the initial state q_0 with Büchi acceptance condition is exactly the Borel set B .

• Now we define the finitary language π . We set

$$\pi := \left\{ \begin{array}{l} s \in 4^{<\omega} \mid \exists j, l \in \omega \exists (m_i)_{i \leq l} \in 2^{l+1} \exists (n_i)_{i \leq l}, (p_i)_{i \leq l}, (r_i)_{i \leq l} \in \omega^{l+1} \\ \\ n_0 \leq M_j \\ \text{and} \\ \forall i \leq l \ n_i \xrightarrow{m_i} p_i \text{ and } p_i + r_i = M_{j+i+1} \\ \text{and} \\ \forall i < l \ p_i = n_{i+1} \\ \text{and} \\ q_{p_i} \in Q_f \\ \text{and} \\ s = \bigwedge_{i \leq l} \mathbf{2}^{n_i} m_i \mathbf{2}^{p_i} \mathbf{2}^{r_i} \mathbf{3} \mathbf{2}^{r_i} \end{array} \right\}.$$

• Let us prove that $\varphi_{N,j}[\pi^\infty \cap K_{N,j}] = E_N$ if $N \leq M_j$.

Let $\gamma \in \pi^\infty \cap K_{N,j}$, and $\alpha := \varphi_{N,j}(\gamma)$. We can write

$$\gamma = \bigwedge_{k \in \omega} [\bigwedge_{i \leq l_k} \mathbf{2}^{n_i^k} m_i^k \mathbf{2}^{p_i^k} \mathbf{2}^{r_i^k} \mathbf{3} \mathbf{2}^{r_i^k}].$$

As this decomposition of γ is in π , we have $n_i^k \xrightarrow{m_i^k} p_i^k$ if $i \leq l_k$, $p_i^k = n_{i+1}^k$ if $i < l_k$, and $q_{p_{l_k}^k} \in Q_f$, for each $k \in \omega$. Moreover, $p_{l_k}^k = n_0^{k+1}$, for each $k \in \omega$, since $\gamma \in K_{N,j}$ implies that $p_{l_k}^k + r_{l_k}^k = r_{l_k}^k + n_0^{k+1} = M_{j+1+m}$ for some integer m . So we get

$$N \xrightarrow{\alpha(0)} p_0^0 \xrightarrow{\alpha(1)} \dots \xrightarrow{\alpha(l_0)} p_{l_0}^0 \xrightarrow{\alpha(l_0+1)} p_0^1 \xrightarrow{\alpha(l_0+2)} \dots \xrightarrow{\alpha(l_0+l_1+1)} p_{l_1}^1 \dots$$

In particular we have

$$q_N^0 \prec q_{p_0^0}^0 \prec \dots \prec q_{p_{l_0}^0}^0 \prec q_{p_0^1}^0 \prec \dots \prec q_{p_{l_1}^1}^0 \dots$$

because $n \xrightarrow{m} p$ implies that $q_n^0 \prec q_p^0$. Note that $|q_{p_{l_k}^k}^1| = |q_N^1| + \sum_{j \leq k} (l_j + 1)$, so that the sequence $(|q_{p_{l_k}^k}^1|)_{k \in \omega}$ is strictly increasing since $|q_n^0| = |q_n^1|$ for each integer n . This implies the existence of $\beta \in P_\infty$ such that $q_{p_{l_k}^k}^0 \prec \beta$ for each $k \in \omega$. Note that $\beta \in P_\infty$ because, for each integer k , $q_{p_{l_k}^k} \in Q_f$. Note also that $(\beta, q_N^1 \alpha) \upharpoonright [k \in R$ for infinitely many k 's. As R is closed under initial segments, $(\beta, q_N^1 \alpha) \upharpoonright [k \in R$ for every $k \in \omega$, so that $q_N^1 \alpha = f(\beta) \in B$. Moreover,

$$c(q_N^1 \alpha, |q_N^1|) = (\beta \upharpoonright [q_N^1], q_N^1) = (q_N^0, q_N^1) = q_N,$$

and $\alpha \in E_N$.

Conversely, let $\alpha \in E_N$. We have to see that $\gamma := \varphi_{N,j}^{-1}(\alpha) \in \pi^\infty$. As $\gamma \in K_{N,j}$, we are allowed to write $\gamma = \mathbf{2}^N \bigwedge [\bigwedge_{i \in \omega} \alpha(i) \mathbf{2}^{M_{j+i+1}} \mathbf{3}^{M_{j+i+1}}]$. We set $\beta := f^{-1}(q_N^1 \alpha)$. There is a sequence of integers $(k_i)_{i \in \omega}$ such that $q_{k_i} = (\beta, q_N^1 \alpha) \upharpoonright [i$. Note that $N \xrightarrow{\alpha(0)} k_{|q_N^1|+1} \xrightarrow{\alpha(1)} k_{|q_N^1|+2} \dots$. As $N \leq M_j$ we get $k_{|q_N^1|+i+1} \leq M_{j+i+1}$. So we can define $n_0 := N$, $p_0 := k_{|q_N^1|+1}$, $r_0 := M_{j+1} - p_0$, $n_1 := p_0$. Similarly, we can define $p_1 := k_{|q_N^1|+2}$, $r_1 := M_{j+2} - p_1$. We go on like this until we find some $q_{p_i} \in Q_f$. This clearly defines a word in π . And we can go on like this, so that $\gamma \in \pi^\infty$.

Thus $\pi^\infty \cap K_{N,j}$ is in $\Gamma(K_{N,j}) \subseteq \Gamma(4^\omega)$. Notice that we proved, among other things, the equality $\varphi_{0,0}[\pi^\infty \cap K_{0,0}] = B$. In particular, $\pi^\infty \cap K_{0,0}$ is Γ -complete in $K_{0,0}$.

Notice that π^∞ codes on $K_{0,0}$ the behaviour of the transition system accepting B . In a similar way, π^∞ codes on $K_{N,j}$ the behaviour of the same transition system, but starting this time from the state q_N instead of the initial state q_0 . But some ω -words in π^∞ are not in $K_{0,0}$ and not even in any $K_{N,j}$ and we do not know what is exactly the complexity of this set of ω -words. However we remark that all the words in π have the same form $\mathbf{2}^N \bigwedge [\bigwedge_{i \leq l} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}]$.

• We are ready to define μ . The idea is that an infinite sequence containing a word in μ cannot be in the union of the $K_{N,j}$'s.

We set

$$\mu^0 := \left\{ \begin{array}{l} s \in 4^{<\omega} \mid \exists l \in \omega \exists (m_i)_{i \leq l+1} \in 2^{l+2} \exists N \in \omega \exists (P_i)_{i \leq l+1}, (R_i)_{i \leq l+1} \in \omega^{l+2} \\ \forall i \leq l+1 \exists j \in \omega P_i = M_j \\ \text{and} \\ P_l \neq R_l \\ \text{and} \\ s = \mathbf{2}^N \frown [\frown_{i \leq l+1} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}] \end{array} \right\},$$

$$\mu^1 := \left\{ \begin{array}{l} s \in 4^{<\omega} \mid \exists l \in \omega \exists (m_i)_{i \leq l+1} \in 2^{l+2} \exists N \in \omega \exists (P_i)_{i \leq l+1}, (R_i)_{i \leq l+1} \in \omega^{l+2} \\ \forall i \leq l+1 \exists j \in \omega P_i = M_j \\ \text{and} \\ \exists j \in \omega (P_l = M_j \text{ and } P_{l+1} \neq M_{j+1}) \\ \text{and} \\ s = \mathbf{2}^N \frown [\frown_{i \leq l+1} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}] \end{array} \right\},$$

$$\mu := \mu^0 \cup \mu^1.$$

All the words in A have the same form $\mathbf{2}^N \frown [\frown_{i \leq l} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}]$. Note that any finite concatenation of words of this form still has this form. Moreover, such a concatenation is in μ^i if its last word is in μ^i .

• Now we prove that μ^∞ is “simple”. The previous remarks show that

$$\mu^\infty = \{ \gamma \in 4^\omega \mid \exists i \in 2 \forall j \in \omega \exists k, n \in \omega \exists t \in (\mu^i)^{n+1} \ n \geq j \text{ and } \gamma \upharpoonright k = \frown_{l \leq n} t(l) \}.$$

This shows that $\mu^\infty \in \mathbf{\Pi}_2^0(4^\omega)$.

• Note again that all words in A have the same form $\mathbf{2}^N \frown [\frown_{i \leq l} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}]$. We set

$$P := \{ \mathbf{2}^N \frown [\frown_{i \in \omega} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}] \in 4^\omega \mid (m_i)_{i \in \omega} \in 2^\omega, N \in \omega, (P_i)_{i \in \omega}, (R_i)_{i \in \omega} \in \omega^\omega \text{ and} \\ \forall i \in \omega \exists j \in \omega P_i = M_j \}.$$

We define a map $F: P \setminus \mu^\infty \rightarrow (\{\emptyset\} \cup \mu) \times \omega^2$ as follows. Let $\gamma := \mathbf{2}^N \frown [\frown_{i \in \omega} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}] \in P \setminus \mu^\infty$, and $j_0 \in \omega$ with $P_0 = M_{j_0}$. If $\gamma \in K_{N, j_0-1}$, then we put $F(\gamma) := (\emptyset, N, j_0)$. If $\gamma \notin K_{N, j_0-1}$, then there is an integer l maximal for which $P_l \neq R_l$ or there is $j \in \omega$ with $P_l = M_j$ and $P_{l+1} \neq M_{j+1}$. Let $j_1 \in \omega$ with $P_{l+2} = M_{j_1}$. We put

$$F(\gamma) := (\mathbf{2}^N \frown [\frown_{i \leq l} m_i \mathbf{2}^{P_i} \mathbf{3} \mathbf{2}^{R_i}] \frown m_{l+1} \mathbf{2}^{P_{l+1}} \mathbf{3}, R_{l+1}, j_1).$$

• Fix $\gamma \in A^\infty$. If $\gamma \notin \mu^\infty$, then $\gamma \in P \setminus \mu^\infty$, $F(\gamma) := (t, S, j)$ is defined. Note that $t \mathbf{2}^S \prec \gamma$, and that $j > 0$. Moreover, $\gamma \upharpoonright t \mathbf{2}^S \in K_{0, j-1}$. Note also that $S \leq M_{j-1}$ if $t = \emptyset$, and that $t \mathbf{2}^S \gamma \upharpoonright (|t|+S) \mathbf{2}^{M_j} \mathbf{3} \notin \mu$. Moreover, there is an integer $N \leq \min(M_{j-1}, S)$ ($N = S$ if $t = \emptyset$) such that $\gamma \upharpoonright t \mathbf{2}^{S-N} \in \pi^\infty \cap K_{N, j-1}$, since the last word in μ in the decomposition of γ (if it exists) ends before $t \mathbf{2}^S$.

- In the sequel we will say that $(t, S, j) \in (\{\emptyset\} \cup \mu) \times \omega^2$ is suitable if $S \leq M_j$ if $t = \emptyset$, $t(|t|-1) = \mathbf{3}$ if $t \in \mu$, and $t \mathbf{2}^S m \mathbf{2}^{M_{j+1}} \mathbf{3} \notin \mu$ if $m \in 2$. We set, for (t, S, j) suitable,

$$P_{t,S,j} := \{ \gamma \in 4^\omega \mid t \mathbf{2}^S \prec \gamma \text{ and } \gamma - t \mathbf{2}^S \in K_{0,j} \}.$$

Note that $P_{t,S,j}$ is a compact subset of $P \setminus \mu^\infty$, and that $F(\gamma) = (t, S, j+1)$ if $\gamma \in P_{t,S,j}$. This shows that the $P_{t,S,j}$'s, for (t, S, j) suitable, are pairwise disjoint. Note also that μ^∞ is disjoint from $\bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j}$.

- We set, for (t, S, j) suitable and $N \leq \min(M_j, S)$ ($N = S$ if $t = \emptyset$),

$$A_{t,S,j,N} := \{ \gamma \in P_{t,S,j} \mid \gamma - t \mathbf{2}^{S-N} \in \pi^\infty \cap K_{N,j} \}.$$

Note that $A_{t,S,j,N} \in \Gamma(4^\omega)$ since $N \leq M_j$.

- The previous discussion shows that

$$A^\infty = \mu^\infty \cup \bigcup_{(t,S,j) \text{ suitable}} \bigcup_{\substack{N \leq \min(M_j, S) \\ N = S \text{ if } t = \emptyset}} A_{t,S,j,N}.$$

As Γ is closed under finite unions, the set

$$A_{t,S,j} := \bigcup_{\substack{N \leq \min(M_j, S) \\ N = S \text{ if } t = \emptyset}} A_{t,S,j,N}$$

is in $\Gamma(4^\omega)$.

- We can write

$$A^\infty = \mu^\infty \setminus \left(\bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j} \right) \cup \bigcup_{(t,S,j) \text{ suitable}} A_{t,S,j} \cap P_{t,S,j}.$$

Note that the $P_{t,S,j}$'s and $\bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j}$ are Δ_3^0 subsets of 4^ω since $(P_{t,S,j})_{(t,S,j) \text{ suitable}}$ is a countable family of closed sets. Moreover, μ^∞ is a $\Pi_2^0 \subseteq \Gamma$ subset of 4^ω . This implies that A^∞ is in $\Delta_3^0\text{-PU}(\Gamma) = \Gamma$. Moreover, the set $A^\infty \cap P_{\emptyset,0,0} = \pi^\infty \cap P_{\emptyset,0,0} = \pi^\infty \cap K_{0,0}$ is Γ -complete. This shows that A^∞ is Γ -hard (any reduction with values in $K_{0,0}$ is also a reduction with values in 4^ω). Thus A^∞ is Γ -complete.

We can now end the proof of Theorem 1.2.

(b) If $\xi = 1$, then we can take $A := \{s \in 2^{<\omega} \mid 0 \prec s \text{ or } \exists k \in \omega \ 10^k 1 \prec s\}$ and $A^\infty = 2^\omega \setminus \{10^\infty\}$ is Σ_1^0 -complete.

- If $\xi = 2$, then we will see in Theorem 2 the existence of $A \subseteq 2^{<\omega}$ such that A^∞ is Σ_2^0 -complete.
- So we may assume that $\xi \geq 3$, and we just have to apply (a) to $\Gamma := \Sigma_\xi^0$.

(c) If $\xi = 1$, then we can take $A := \{0\}$ and $A^\infty = \{0^\infty\}$ is Π_1^0 -complete.

- If $\xi = 2$, then we can take $A := \{0^k 1 \mid k \in \omega\}$ and $A^\infty = P_\infty$ is Π_2^0 -complete.

- So we may assume that $\xi \geq 3$, and we just have to apply (a) to $\Gamma := \Pi_\xi^0$.

(d) First notice that $D_2(\Sigma_\xi^0) = \{B \setminus C \mid B, C \in \Sigma_\xi^0\}$. Indeed, \subseteq is clear, and \supseteq comes from the fact that $B \setminus C = (B \cup C) \setminus C$. This implies that $\check{D}_2(\Sigma_\xi^0) = \{B \cup C \mid B \in \Sigma_\xi^0 \text{ and } C \in \Pi_\xi^0\}$. A consequence of this is the closure of $\check{D}_2(\Sigma_\xi^0)$ under finite unions. Another consequence is $\Delta_\xi^0[\check{D}_2(\Sigma_\xi^0)] \subseteq \check{D}_2(\Sigma_\xi^0)$. Indeed, if $D := B \cup C \in \check{D}_2(\Sigma_\xi^0)(X)$ and $E \in \Delta_\xi^0(D)$, then choose $\Sigma \in \Sigma_\xi^0(X)$ and $\Pi \in \Pi_\xi^0(X)$ such that $E = \Sigma \cap D = \Pi \cap D$. We get $E = (\Sigma \cap B) \cup (\Pi \cap C) \in \check{D}_2(\Sigma_\xi^0)(X)$.

- If $\xi = 1$, then we can take $A := \{s \in 2^{<\omega} \mid 0 \prec s \text{ or } \exists q \in \omega (101)^{q1^3} \prec s \text{ or } s = 10^2\}$ and $A^\infty = \bigcup_{p \in \omega} [N_{(10^2)^{p0}} \cup (\bigcup_{q \in \omega} N_{(10^2)^{p(101)^{q1^3}}})] \cup \{(10^2)^\infty\}$ is $\check{D}_2(\Sigma_1^0)$ -complete (see §7 in [Lec05], and also example 9 in [Sta97b]).

- If $\xi = 2$, then we can take $A := \{s \in 2^{<\omega} \mid 1^2 \prec s \text{ or } s = 0\}$ and

$$A^\infty = \left(\{0^\infty\} \cup \bigcup_{p \in \omega} N_{0^p 1^2} \right) \cap [(2^\omega \setminus P_\infty) \cup \{\alpha \in 2^\omega \mid \forall m \in \omega \exists n \geq m \alpha(n) = \alpha(n+1) = 1\}]$$

is $\check{D}_2(\Sigma_2^0)$ -complete (see §7 in [Lec05]).

- So we may assume that $\xi \geq 3$, and we just have to apply (a) to $\Gamma := \check{D}_2(\Sigma_\xi^0)$.

(e) Let X be a zero-dimensional Polish space, and $E, F \in D_\eta(\Sigma_\xi^0)(X)$. By Lemma 4.2 in [vEn], $E \times F$ is $D_\eta(\Sigma_\xi^0)$. Now let $\mathcal{C} \subseteq 2^\omega$ be $D_\eta(\Sigma_\xi^0)$ -complete, $h: 2^\omega \times 2^\omega \rightarrow 2^\omega$ continuous with $\mathcal{C} \times \mathcal{C} = h^{-1}(\mathcal{C})$, and $f, g: X \rightarrow 2^\omega$ continuous with $E = f^{-1}(\mathcal{C})$ and $F = g^{-1}(\mathcal{C})$. It is clear that the map $c: X \rightarrow 2^\omega$ defined by $c(x) := h[f(x), g(x)]$ satisfies $E \cap F = c^{-1}(\mathcal{C})$. This shows that $D_\eta(\Sigma_\xi^0)$ is closed under finite intersections. Thus $\check{D}_\eta(\Sigma_\xi^0)$ is closed under finite unions.

Note also that if $D \in D_\eta(\Sigma_\xi^0)$ and $B \in \Sigma_\xi^0$, then $B \cup D \in D_\eta(\Sigma_\xi^0)$. Indeed, let $(A_\theta)_{\theta < \eta} \subseteq \Sigma_\xi^0$ be an increasing sequence with $D = D[(A_\theta)_{\theta < \eta}]$. We set $B_0 := \emptyset$, $B_1 := B$, and $B_{2+\theta} := A_\theta \cup B$ if $\theta < \eta$. Then $(B_\theta)_{\theta < \eta} \subseteq \Sigma_\xi^0$ is increasing, and $D[(B_\theta)_{\theta < \eta}] = B \cup \bigcup_{2\rho+1 < \eta} (A_{2\rho+1} \cup B) \setminus (A_{2\rho} \cup B) = B \cup D$ since η is even. This shows that if $D \in \check{D}_\eta(\Sigma_\xi^0)$ and $B \in \Pi_\xi^0$, then $B \cap D \in \check{D}_\eta(\Sigma_\xi^0)$. This implies the inclusion $\Delta_\xi^0[\check{D}_\eta(\Sigma_\xi^0)] \subseteq \check{D}_\eta(\Sigma_\xi^0)$.

Now we can apply (a) to $\Gamma := \check{D}_\eta(\Sigma_\xi^0)$. □

As we have said above it remains a Borel class for which we have not yet got a complete ω -power: the class Σ_2^0 . Note that it is easy to see that the classical example of a Σ_2^0 -complete set, the set $2^\omega \setminus P_\infty$, is not an ω -power. However we are going to prove the following result.

Theorem 2 *There is a recursive (and even context-free) language $A \subseteq 2^{<\omega}$ such that $A^\infty \in \Sigma_2^0 \setminus \Pi_2^0$.*

Proof. By Proposition 11 in [Lec05], it is enough to find $A \subseteq 3^{<\omega}$. We set, for $j < 3$ and $s \in 3^{<\omega}$,

$$n_j(s) := \text{Card}\{i < |s| \mid s(i) = j\},$$

$$T := \{\alpha \in 3^{<\omega} \mid \forall l < 1 + |\alpha| \ n_2(\alpha[l]) \leq n_1(\alpha[l])\}.$$

• We inductively define, for $s \in T \cap 3^{<\omega}$, $s^{\leftrightarrow} \in 2^{<\omega}$ as follows:

$$s^{\leftrightarrow} := \begin{cases} \emptyset & \text{if } s = \emptyset, \\ t^{\leftrightarrow} \varepsilon & \text{if } s = t\varepsilon \text{ and } \varepsilon < 2, \\ t^{\leftrightarrow}, & \text{except that its last 1 is replaced with 0, if } s = t\mathbf{2}. \end{cases}$$

• We will extend this definition to infinite sequences. To do this, we introduce a notion of limit. Fix $(s_n)_{n \in \omega} \subseteq 2^{<\omega}$. We define $\lim_{n \rightarrow \infty} s_n \in 2^{<\omega}$ as follows. For each $t \in 2^{<\omega}$,

$$t \prec \lim_{n \rightarrow \infty} s_n \Leftrightarrow \exists n_0 \in \omega \ \forall n \geq n_0 \ t \prec s_n.$$

• If $\alpha \in T \cap 3^\omega$, then we set $\alpha^{\leftrightarrow} := \lim_{n \rightarrow \infty} (\alpha[n])^{\leftrightarrow}$. We define $e : T \cap 3^\omega \rightarrow 2^\omega$ by $e(\alpha) := \alpha^{\leftrightarrow}$. Note that $T \cap 3^\omega \in \Pi_1^0(3^\omega)$, and e is a Σ_2^0 -recursive partial function on $T \cap 3^\omega$, since for $t \in 2^{<\omega}$ we have

$$t \prec e(\alpha) \Leftrightarrow \exists n_0 \in \omega \ \forall n \geq n_0 \ t \prec (\alpha[n])^{\leftrightarrow}.$$

• We set $E := \{s \in T \cap 3^{<\omega} \mid n_2(s) = n_1(s) \text{ and } s \neq \emptyset \text{ and } 1 < [s[(|s|-1)]]^{\leftrightarrow}\}$. Note that $\emptyset \neq s^{\leftrightarrow} \prec 0^\infty$, and that $s(|s|-1) = \mathbf{2}$ changes $s(0) = [s[(|s|-1)]]^{\leftrightarrow}(0) = 1$ into 0 if $s \in E$.

• If $S \subseteq 3^{<\omega}$, then $S^* := \{\bigcap_{i < l} s_i \in 3^{<\omega} \mid l \in \omega \text{ and } (s_i)_{i < l} \subseteq S\}$. We put

$$A := \{0\} \cup E \cup \{\bigcap_{j \leq k} (c_j 1) \in 3^{<\omega} \mid [\forall j \leq k \ c_j \in (\{0\} \cup E)^*] \text{ and } [k > 0 \text{ or } (k = 0 \text{ and } c_0 \neq \emptyset)]\}.$$

Note that A is recursive.

• In the proof of Theorem 1.2.(b) we met the set $\{s \in 2^{<\omega} \mid 0 \prec s \text{ or } \exists k \in \omega \ 10^k 1 \prec s\}$. We will call this set B , and $B^\infty = 2^\omega \setminus \{10^\infty\}$ is Σ_1^0 -complete (and even Σ_1^0). Let us show that $A^\infty = e^{-1}(B^\infty)$.

- By induction on $|t|$, we get $(st)^{\leftrightarrow} = s^{\leftrightarrow} t^{\leftrightarrow}$ if $s, t \in T \cap 3^{<\omega}$. Let us show that $(s\beta)^{\leftrightarrow} = s^{\leftrightarrow} \beta^{\leftrightarrow}$ if moreover $\beta \in T \cap 3^\omega$.

Assume that $t \prec (s\beta)^{\leftrightarrow}$. Then there is $m_0 \geq |s|$ such that, for $m \geq m_0$,

$$t \prec [(s\beta)[m]]^{\leftrightarrow} = [s\beta[(m-|s|)]]^{\leftrightarrow} = s^{\leftrightarrow} [\beta[(m-|s|)]]^{\leftrightarrow}.$$

This implies that $t \prec s^{\leftrightarrow} \beta^{\leftrightarrow}$ if $|t| < |s^{\leftrightarrow}|$. If $|t| \geq |s^{\leftrightarrow}|$, then there is $m_1 \in \omega$ such that, for $m \geq m_1$, $\beta^{\leftrightarrow} \prec [(|t|-|s^{\leftrightarrow}|) \prec [\beta[(m-|s|)]]^{\leftrightarrow}$. Here again, we get $t \prec s^{\leftrightarrow} \beta^{\leftrightarrow}$. Thus $(s\beta)^{\leftrightarrow} = s^{\leftrightarrow} \beta^{\leftrightarrow}$.

Let $(s_i)_{i \in \omega} \subseteq T \cap 3^{<\omega}$. Then $\bigcap_{i \in \omega} s_i \in T$, and $(\bigcap_{i \in \omega} s_i)^{\leftrightarrow} = \bigcap_{i \in \omega} s_i^{\leftrightarrow}$, by the previous facts.

- Let $(a_i)_{i \in \omega} \in (A \setminus \{\emptyset\})^\omega$ and $\alpha := \bigwedge_{i \in \omega} a_i$. As $A \subseteq T$, $e(\alpha) = (\bigwedge_{i \in \omega} a_i)^{\leftrightarrow} = \bigwedge_{i \in \omega} a_i^{\leftrightarrow}$.

If $a_0 \in \{0\} \cup E$, then $\emptyset \neq a_0^{\leftrightarrow} \prec 0^\infty$, thus $e(\alpha) \in N_0 \subseteq 2^\omega \setminus \{10^\infty\} = B^\infty$.

If $a_0 \notin \{0\} \cup E$, then $a_0 = \bigwedge_{j \leq k} (c_j 1)$, thus $a_0^{\leftrightarrow} = \bigwedge_{j \leq k} (c_j^{\leftrightarrow} 1)$.

If $c_0 \neq \emptyset$, then $e(\alpha) \in B^\infty$ as before.

If $c_0 = \emptyset$, then $k > 0$, so that $e(\alpha) \neq 10^\infty$ since $e(\alpha)$ has at least two coordinates equal to 1.

We proved that $A^\infty \subseteq e^{-1}(B^\infty)$.

- Assume that $e(\alpha) \in B^\infty$. We have to find $(a_i)_{i \in \omega} \subseteq A \setminus \{\emptyset\}$ with $\alpha = \bigwedge_{i \in \omega} a_i$. We split into cases:

1. $e(\alpha) = 0^\infty$.

1.1. $\alpha(0) = 0$.

In this case $\alpha - 0 \in T$ and $e(\alpha - 0) = 0^\infty$. Moreover, $0 \in A$. We put $a_0 := 0$.

1.2. $\alpha(0) = 1$.

In this case there is a coordinate j_0 of α equal to **2** ensuring that $\alpha(0)$ is replaced with a 0 in $e(\alpha)$. We put $a_0 := \alpha \upharpoonright (j_0 + 1)$, so that $a_0 \in E \subseteq A$, $\alpha - a_0 \in T$ and $e(\alpha - a_0) = 0^\infty$.

Now the iteration of the cases 1.1 and 1.2 shows that $\alpha \in A^\infty$.

2. $e(\alpha) = 0^{k+1}10^\infty$ for some $k \in \omega$.

As in case 1, there is $c_0 \in (\{0\} \cup E)^*$ such that $c_0 \prec \alpha$, $c_0^{\leftrightarrow} = 0^{k+1}$, $\alpha - c_0 \in T$ and $e(\alpha - c_0) = 10^\infty$. Note that $\alpha(|c_0|) = 1$, $\alpha - (c_0 1) \in T$ and $e[\alpha - (c_0 1)] = 0^\infty$. We put $a_0 := c_0 1$, and argue as in case 1.

3. $e(\alpha) = (\bigwedge_{j \leq l+1} 0^{k_j} 1) 0^\infty$ for some $l \in \omega$.

The previous cases show the existence of $(c_j)_{j \leq l+1} \subseteq (\{0\} \cup E)^*$ such that $a_0 := \bigwedge_{j \leq l+1} c_j 1 \prec \alpha$, $\alpha - a_0 \in T$ and $e(\alpha - a_0) = 0^\infty$. We are done since $a_0 \in A$.

4. $e(\alpha) = \bigwedge_{j \in \omega} 0^{k_j} 1$.

An iteration of the discussion of case 3 shows that we can take a_i of the form $\bigwedge_{j \leq l+1} c_j 1$.

• The previous discussion shows that $A^\infty = e^{-1}(B^\infty)$. As e is Σ_2^0 -recursive, $e^{-1}(B^\infty) \in \Sigma_2^0(3^\omega)$.

It remains to see that $e^{-1}(B^\infty) \notin \Pi_2^0$. We argue by contradiction. We know that $B^\infty = 2^\omega \setminus \{10^\infty\}$, so $e^{-1}(\{10^\infty\}) = (T \cap 3^\omega) \setminus e^{-1}(B^\infty)$ is a Σ_2^0 subset of 3^ω since $T \cap 3^\omega$ is closed in 3^ω . Thus $e^{-1}(\{10^\infty\})$ is a countable union of compact subsets of 3^ω .

Consider now the cartesian product $(\{0\} \cup E)^\omega$ of countably many copies of $\{0\} \cup E$. The set $\{0\} \cup E$ is countable and it can be equipped with the discrete topology. The product $(\{0\} \cup E)^\omega$ is equipped with the product topology of the discrete topology on $\{0\} \cup E$. In these conditions, the topological space $(\{0\} \cup E)^\omega$ is homeomorphic to the Baire space ω^ω .

Consider now the map $h: (\{0\} \cup E)^\omega \rightarrow e^{-1}(\{10^\infty\})$ defined by $h(\gamma) := 1[\bigwedge_{i \in \omega} \gamma_i]$ for each sequence $\gamma = (\gamma_0, \gamma_1, \dots) \in (\{0\} \cup E)^\omega$. The map h is a homeomorphism by the previous discussion. As $(\{0\} \cup E)^\omega$ is homeomorphic to ω^ω , the Baire space ω^ω is also homeomorphic to $e^{-1}(\{10^\infty\})$. This implies that ω^ω is a countable union of compact sets. But this is absurd, by Theorem 7.10 in [Kec95].

• It remains to see that A is context-free. We assume here that the reader is familiar with the theory of formal languages and of context-free languages; basic notions may be found in the Handbook Chapter [ABB96].

It is easy to see that the language E is in fact accepted by a 1-counter automaton: it is the set of words $s \in 3^{<\omega}$ such that

$$\forall 1 \leq l < |s| \quad n_2(s[l]) < n_1(s[l]) \quad \text{and} \quad n_2(s) = n_1(s) \quad \text{and} \quad s(0) = 1 \quad \text{and} \quad s(|s| - 1) = 2.$$

This implies that A is also accepted by a 1-counter automaton because the class of 1-counter languages is closed under concatenation and star operation. In particular A is a context-free language because the class of languages accepted by 1-counter automata form a strict subclass of the class of context-free languages. \square

Remark. The operation $\alpha \rightarrow \alpha^{\leftarrow}$ we have defined is very close to the erasing operation defined by J. Duparc in his study of the Wadge hierarchy (see [Dup01]). However we have modified this operation in such a way that α^{\leftarrow} is always infinite when α is infinite, and that it has the good property with regard to ω -powers and topological complexity.

Question. What are the Wadge classes Γ for which there is $A \subseteq 2^{<\omega}$ such that A^∞ is Γ -complete? We have seen that Theorem 1.2 solves completely the case where Γ is a Borel class, and it also solves the problem for some more Wadge classes. The problem is solved for a few other Wadge classes in [Lec05]. We do not know (yet?) any Wadge class for which this problem cannot be solved.

3 Effective descriptive set theory background.

Basic facts and notation.

• In [Mos80], the classical *arithmetical hierarchy* is defined as follows (see 3E). Let X be a recursively presented Polish space, $[N(X, k)]_{k \in \omega}$ an effective enumeration of a neighborhood basis for the topology of X , and $B \subseteq X$. We say that $B \in \Sigma_1^0(X)$ if there is a recursive map $\varepsilon: \omega \rightarrow \omega$ such that $B = \bigcup_{i \in \omega} N[X, \varepsilon(i)]$. If $n \geq 1$ is an integer, then Π_n^0 is the class of complements of Σ_n^0 sets. We say that $B \in \Sigma_{n+1}^0$ if there is $C \in \Pi_n^0(\omega \times X)$ such that $B = \exists^\omega C := \{x \in X \mid \exists i \in \omega (i, x) \in C\}$. We also set $\Delta_n^0 := \Sigma_n^0 \cap \Pi_n^0$.

• We say that $\gamma \in \Sigma_1^0$ if $\{k \in \omega \mid \gamma \in N(\omega^\omega, k)\} \in \Sigma_1^0(\omega)$. Let $\beta \in 2^\omega$. The *relativization* $\Sigma_1^0(\beta)$ of Σ_1^0 to β is defined as follows. A set $P \subseteq X$ is in $\Sigma_1^0(\beta)$ if there is $Q \in \Sigma_1^0(2^\omega \times X)$ such that $P = Q_\beta$. As before we say that $\gamma \in \Sigma_1^0(\beta)$ if $\{k \in \omega \mid \gamma \in N(\omega^\omega, k)\} \in \Sigma_1^0(\beta)(\omega)$.

- Recall the existence of a *good parametrization* in Σ_n^0 for Σ_n^0 (see 3E.2, 3F.6 and 3H.1 in [Mos80]). This means that there is a system of sets $G^{\Sigma_n^0, X} \in \Sigma_n^0(\omega^\omega \times X)$ such that for each recursively presented Polish space X and for each $P \subseteq X$,

$$P \in \Sigma_n^0 \Leftrightarrow \exists \gamma \in \omega^\omega \ P = G_\gamma^{\Sigma_n^0, X},$$

$$P \in \Sigma_n^0 \Leftrightarrow \exists \gamma \in \Sigma_1^0 \ P = G_\gamma^{\Sigma_n^0, X}.$$

Moreover, if X is a recursively presented Polish space of type at most 1 (i.e., a finite product of spaces equal to ω , ω^ω or 2^ω), and Y is a recursively presented Polish space, then there is $S_{\Sigma_n^0}^{X, Y} : \omega^\omega \times X \rightarrow \omega^\omega$ recursive so that

$$(\gamma, x, y) \in G^{\Sigma_n^0, X \times Y} \Leftrightarrow [S_{\Sigma_n^0}^{X, Y}(\gamma, x), y] \in G^{\Sigma_n^0, Y}.$$

- Let $f : X \rightarrow Y$ be a partial function, $D \subseteq \text{Domain}(f)$ and $P \subseteq X \times \omega$. Then P computes f on D if

$$x \in D \Rightarrow \forall k \in \omega \ [f(x) \in N(Y, k) \Leftrightarrow (x, k) \in P].$$

If P is in some pointclass Γ and computes f on D , then we say that f is Γ -recursive on D . We also say recursive on D for Σ_1^0 -recursive on D .

- We also recall the notation for the coding of partial recursive functions from X into Y introduced in [Mos80] (see 7A). We first define a partial function $U : \omega^\omega \times X \rightarrow Y$ by

$$U(\gamma, x) \downarrow \Leftrightarrow U(\gamma, x) \text{ is defined} \Leftrightarrow \exists y \in Y \ \forall k \in \omega \ [y \in N(Y, k) \Leftrightarrow (\gamma, x, k) \in G^{\Sigma_1^0, X \times \omega}],$$

$$U(\gamma, x) := \text{the unique } y \in Y \text{ such that } \forall k \in \omega \ [y \in N(Y, k) \Leftrightarrow (\gamma, x, k) \in G^{\Sigma_1^0, X \times \omega}].$$

Now let $\gamma \in \omega^\omega$. The function $\{\gamma\}^{X, Y} : X \rightarrow Y$ is defined by $\{\gamma\}^{X, Y}(x) := U(\gamma, x)$. Then a partial function $f : X \rightarrow Y$ is recursive on its domain if and only if there is $\gamma \in \Sigma_1^0$ such that $f(x) = \{\gamma\}^{X, Y}(x)$ when $f(x)$ is defined. More generally, the functions of the form $\{\gamma\}^{X, Y}$ are the partial continuous functions from a subset of X into Y . We will write $\{\gamma\}$ instead of $\{\gamma\}^{X, Y}$ when $Y = \omega^\omega$, in order to simplify the notation.

If X is of type at most 1 and Z is a recursively presented Polish space, then there is a recursive map $S_{\Sigma_1^0}^{X, Y, Z} : \omega^\omega \times X \rightarrow \omega^\omega$ such that $\{\gamma\}^{X \times Y, Z}(x, y) = \{S_{\Sigma_1^0}^{X, Y, Z}(\gamma, x)\}^{Y, Z}(y)$ if $(\gamma, x) \in \omega^\omega \times X$.

Kleene's Recursion Theorem asserts that if $f : \omega^\omega \times X \rightarrow Y$ is recursive on its domain, then there is $\varepsilon^* \in \Sigma_1^0$ such that $f(\varepsilon^*, x) = \{\varepsilon^*\}^{X, Y}(x)$ when $f(\varepsilon^*, x)$ is defined (see 7A.2 in [Mos80]).

- We will use the following basic maps:

- We first define a one-to-one map $\langle \cdot \rangle : \omega^{<\omega} \rightarrow \omega$. Let $(p_n)_{n \in \omega}$ be the sequence of prime numbers. We set $\langle \emptyset \rangle := 1$, and, if $t := (t_0, \dots, t_l) \in \omega^{l+1}$, then we set $\bar{t} := \langle t_0, \dots, t_l \rangle := p_0^{t_0+1} \dots p_l^{t_l+1}$.

- If $k \in \omega$, then we say that “Seq(k)” (i.e., “ k is a sequence”) if $k = \langle t_0, \dots, t_{l-1} \rangle$ for some t_0, \dots, t_{l-1} .

- The length $\text{lh}(k)$ of $k \in \omega$ is l if $\text{Seq}(k)$ and $k = \langle t_0, \dots, t_{l-1} \rangle$, 0 otherwise.

- If $k, i \in \omega$, then we define $(k)_i := t_i$ if $\text{Seq}(k)$, $k = \langle t_0, \dots, t_{l-1} \rangle$ and $i < l$, 0 otherwise.

- If $\gamma \in \omega^\omega$ and $i \in \omega$, then we define $(\gamma)_i \in \omega^\omega$ by $(\gamma)_i(j) := \gamma(\langle i, j \rangle)$. But here we do not use the injection $(i, j) \mapsto \langle i, j \rangle$ above, since we want a bijection from ω^2 into ω . So we use the notation $\langle i, j \rangle$ for $2^i \cdot (2j+1) - 1$, when $(\gamma)_i$ is concerned. The inverse bijection is denoted $s \mapsto [(s)_0, (s)_1]$.

Borel codes and closure properties.

Notation. We give a coding of Borel sets slightly different from the one given in [Mos80] (see 7B), since there is a problem for Σ_1^0 . It can be found in [Lou??]. We define by induction on the countable ordinal $\xi \geq 1$ the set BC_ξ of Borel codes for Σ_ξ^0 as follows. If $\gamma \in \omega^\omega$, then we define $\gamma^* \in \omega^\omega$ by $\gamma^*(i) := \gamma(i+1)$. We set

$$BC_1 := \{ \gamma \in \omega^\omega \mid \gamma(0) = 0 \},$$

$$BC_\xi := \left\{ \gamma \in \omega^\omega \mid \gamma(0) = 1 \text{ and } \forall i \in \omega \{ \gamma^*(i) \downarrow \text{ and } \{ \gamma^*(i) \} \in \bigcup_{1 \leq \eta < \xi} BC_\eta \} \text{ if } \xi \geq 2. \right.$$

The set of Borel codes is $BC := \bigcup_{1 \leq \xi < \omega_1} BC_\xi$. We also set $BC^* := \bigcup_{2 \leq \xi < \omega_1} BC_\xi$.

Now let X be a recursively presented Polish space. We define $\rho^X : BC \rightarrow \Delta_1^1(X)$ by induction:

$$\rho^X(\gamma) := \begin{cases} \bigcup_{i \in \omega} N[X, \gamma^*(i)] & \text{if } \gamma \in BC_1, \\ \bigcup_{i \in \omega} X \setminus \rho^X[\{\gamma^*(i)\}] & \text{if } \gamma \in BC^*. \end{cases}$$

Clearly, $\rho^X[BC_\xi] = \Sigma_\xi^0(X)$, by induction on ξ . The following is a consequence of 7B.1.(ii).(a) in [Mos80]. It expresses the fact that the class of Borel sets is uniformly closed under complementation.

Lemma 3.1 *There is a recursive map $u_- : \omega^\omega \rightarrow \omega^\omega$ such that for each $1 \leq \xi < \omega_1$ and for each $\gamma \in BC_\xi$, $u_-(\gamma) \in BC_{\xi+1}$, and $\rho^X[u_-(\gamma)] = \neg \rho^X(\gamma)$ for each recursively presented Polish space X .*

Proof. Just copy the proof of 7B.1.(ii).(a) in [Mos80]: it gives more than the statement in [Mos80]. \square

In the sequel we will need a refinement of 7B.1.(iii) in [Mos80]:

Lemma 3.2 *Let X be a recursively presented Polish space of type at most 1. Then there is a recursive map $u_s^X : \omega^\omega \times X \rightarrow \omega^\omega$ such that for each $1 \leq \xi < \omega_1$, for each $\gamma \in BC_\xi$ and for each $x \in X$, $u_s^X(\gamma, x) \in BC_\xi$, and $\rho^Y[u_s^X(\gamma, x)] = \rho^{X \times Y}(\gamma)_x$ for each recursively presented Polish space Y .*

Some of the ideas of the proof are contained in 7A.3 in [Mos80].

Proof. For $\xi = 1$, we define a subset of $\omega^\omega \times X \times \omega$ by

$$(\gamma, x, k) \in P \Leftrightarrow \exists i \in \omega \left(k = \left\langle 0, \left(\gamma^*(i) \right)_2 \right\rangle \text{ and } x \in N \left[X, \left\langle 0, \left(\gamma^*(i) \right)_1 \right\rangle \right] \right).$$

By 3D.5 in [Mos80], Σ_1^0 is closed under recursive substitutions, so that $P \in \Sigma_1^0$. By 3C.4 in [Mos80], there is $P^* \in \Delta_1^0(\omega^\omega \times X \times \omega^2)$ with

$$(\gamma, x, k) \in P \Leftrightarrow \exists n \in \omega (\gamma, x, k, n) \in P^*.$$

We define a map $g: \omega^\omega \times X \rightarrow \omega^\omega$ by

$$g(\gamma, x)(j) := \begin{cases} (j-1)_0 & \text{if } j > 0 \text{ and } [\gamma, x, (j-1)_0, (j-1)_1] \in P^*, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, g is recursive and $g(\gamma, x) \in BC_1$.

- For the general case, we define a partial function $\psi: (\omega^\omega)^2 \times X \times \omega \rightarrow \omega^\omega$ by

$$\psi(\varepsilon, \gamma, x, i) := \begin{cases} g[\{\gamma^*\}(i), x] & \text{if } \{\gamma^*\}(i)(0) = 0, \\ \{\varepsilon\}[\{\gamma^*\}(i), x] & \text{if } \{\gamma^*\}(i)(0) = 1. \end{cases}$$

By 3G.1 and 3G.2 in [Mos80], the collection of partial functions which are recursive on their domain is closed under composition, so that ψ is recursive on its domain. Let $\nu \in \Sigma_1^0$ such that

$$\psi(\varepsilon, \gamma, x, i) = \{\nu\}(\varepsilon, \gamma, x, i)$$

if $\psi(\varepsilon, \gamma, x, i)$ is defined. We define a recursive map $\varphi: (\omega^\omega)^2 \times X \rightarrow \omega^\omega$ by

$$\varphi(\varepsilon, \gamma, x) := \begin{cases} g(\gamma, x) & \text{if } \gamma(0) = 0, \\ 1 \frown S_{\Sigma_1^0}^{(\omega^\omega)^2 \times X, \omega, \omega^\omega}(\nu, \varepsilon, \gamma, x) & \text{if } \gamma(0) \neq 0. \end{cases}$$

By Kleene's Recursion Theorem, there is $\varepsilon^* \in \Sigma_1^0$ such that $\varphi(\varepsilon^*, \gamma, x) = \{\varepsilon^*\}(\gamma, x)$ for each (γ, x) in $\omega^\omega \times X$. We put $u_s^X(\gamma, x) := \{\varepsilon^*\}(\gamma, x)$. Note that the map u_s^X is a total recursive map. We prove that $u_s^X(\gamma, x)$ satisfies the required properties by induction on ξ .

- Let $(\gamma, x) \in BC_1 \times X$. We have $u_s^X(\gamma, x) = \{\varepsilon^*\}(\gamma, x) = \varphi(\varepsilon^*, \gamma, x) = g(\gamma, x)$. So $u_s^X(\gamma, x)$ is in BC_1 , by the previous discussion. If moreover Y is a recursively presented Polish space, then using the proof of 3B.1 in [Mos80] we get

$$\begin{aligned} y \in \rho^{X \times Y}(\gamma)_x &\Leftrightarrow \exists i \in \omega \ (x, y) \in N[X \times Y, \gamma^*(i)] \\ &\Leftrightarrow \exists i \in \omega \ \left(y \in N\left[Y, \langle 0, (\gamma^*(i))_2 \rangle\right] \text{ and } x \in N\left[X, \langle 0, (\gamma^*(i))_1 \rangle\right] \right) \\ &\Leftrightarrow \exists k \in \omega \ [y \in N(Y, k) \text{ and } (\gamma, x, k) \in P] \\ &\Leftrightarrow \exists i \in \omega \ \left(y \in N[Y, (i)_0] \text{ and } [\gamma, x, (i)_0, (i)_1] \in P^* \right) \\ &\Leftrightarrow \exists i \in \omega \ y \in N\left(Y, [g(\gamma, x)]^*(i)\right) \\ &\Leftrightarrow y \in \rho^Y[g(\gamma, x)]. \end{aligned}$$

- Now let $(\gamma, x) \in BC_\xi \times X$, with $\xi \geq 2$. We have

$$u_s^X(\gamma, x) = \{\varepsilon^*\}(\gamma, x) = \varphi(\varepsilon^*, \gamma, x) = 1 \frown S_{\Sigma_1^0}^{(\omega^\omega)^2 \times X, \omega, \omega^\omega}(\nu, \varepsilon^*, \gamma, x).$$

As $\gamma \in BC_\xi$, $\{\gamma^*\}(i)$ is defined for each integer i . In particular, $\psi(\varepsilon^*, \gamma, x, i)$ is defined for each (γ, x, i) in $\omega^\omega \times X \times \omega$ since $\{\gamma^*\}(i)(0) \in 2$, and equal to

$$\{\nu\}(\varepsilon^*, \gamma, x, i) = \{S_{\Sigma_1^0}^{(\omega^\omega)^2 \times X, \omega, \omega^\omega}(\nu, \varepsilon^*, \gamma, x)\}(i).$$

This shows that $\{u_s^X(\gamma, x)^*\}(i)$ is defined for each integer i . If $\{\gamma^*\}(i)(0) = 0$, then

$$\{u_s^X(\gamma, x)^*\}(i) = g[\{\gamma^*\}(i), x] = u_s^X[\{\gamma^*\}(i), x].$$

As $\{\gamma^*\}(i) \in BC_1$, $u_s^X[\{\gamma^*\}(i), x]$ is in BC_1 too. Similarly, if $\{\gamma^*\}(i)(0) = 1$, then

$$\{u_s^X(\gamma, x)^*\}(i) = \{\varepsilon^*\}[\{\gamma^*\}(i), x] = u_s^X[\{\gamma^*\}(i), x].$$

Then $u_s^X[\{\gamma^*\}(i), x] \in BC_\eta$ for some $1 \leq \eta < \xi$, by induction assumption. This shows that $u_s^X(\gamma, x)$ is in BC_ξ . If Y is a recursively presented Polish space, then $\rho^Y(u_s^X[\{\gamma^*\}(i), x]) = \rho^{X \times Y}[\{\gamma^*\}(i)]_x$, by induction assumption. This shows that $\rho^Y[u_s^X(\gamma, x)] = \rho^{X \times Y}(\gamma)_x$. \square

Lemma 3.2 expresses, among other things, the fact that the pointclasses Σ_ξ^0 are uniformly closed under taking sections at points in spaces of type at most 1. Similarly, we now prove another lemma stating, among other things, that the pointclasses Σ_ξ^0 are uniformly closed under substitutions of partial recursive functions (when δ below is recursive).

Lemma 3.3 *Let X, Y be recursively presented Polish spaces. Then there is $u_r^{X, Y} : (\omega^\omega)^2 \rightarrow \omega^\omega$ recursive such that for each $1 \leq \xi < \omega_1$, for each $\gamma \in BC_\xi$ and for each $\delta \in \omega^\omega$, $u_r^{X, Y}(\gamma, \delta) \in BC_\xi$. Moreover, we have $x \in \rho^X[u_r^{X, Y}(\gamma, \delta)] \Leftrightarrow \{\delta\}^{X, Y}(x) \in \rho^Y(\gamma)$ if $\{\delta\}^{X, Y}(x)$ is defined.*

Proof. For $\xi = 1$, we choose $P \in \Sigma_1^0(\omega^\omega \times X \times \omega)$ such that

$$U(\delta, x) \downarrow \Rightarrow \forall k \in \omega [U(\delta, x) \in N(Y, k) \Leftrightarrow (\delta, x, k) \in P].$$

(this is possible since U is recursive on its domain; see 7A.1 in [Mos80]).

By 3C.4 and 3C.5 in [Mos80], there is $P^* \in \Delta_1^0(\omega^\omega \times \omega^3)$ with

$$(\delta, x, k) \in P \Leftrightarrow \exists i \in \omega \left(x \in N[X, (i)_0] \text{ and } [\delta, k, (i)_0, (i)_1] \in P^* \right).$$

We define a map $g : (\omega^\omega)^2 \rightarrow \omega^\omega$ by

$$g(\gamma, \delta)(j) := \begin{cases} ((j-1)_0)_0 & \text{if } j > 0 \text{ and } [\delta, \gamma^*[(j-1)_1], ((j-1)_0)_0, ((j-1)_0)_1] \in P^*, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, g is recursive and $g(\gamma, \delta) \in BC_1$.

- For the general case, we define a partial function $\psi: (\omega^\omega)^3 \times \omega \rightarrow \omega^\omega$ by

$$\psi(\varepsilon, \gamma, \delta, i) := \begin{cases} g[\{\gamma^*\}(i), \delta] & \text{if } \{\gamma^*\}(i)(0) = 0, \\ \{\varepsilon\}[\{\gamma^*\}(i), \delta] & \text{if } \{\gamma^*\}(i)(0) = 1. \end{cases}$$

We argue as in the proof of Lemma 3.2 to define $\varphi: (\omega^\omega)^3 \rightarrow \omega^\omega$, and we put $u_r^{X,Y}(\gamma, \delta) := \{\varepsilon^*\}(\gamma, \delta)$. The map $u_r^{X,Y}$ is a total recursive map. We show that $u_r^{X,Y}(\gamma, \delta)$ satisfies the required properties by induction on ξ .

- If $(\gamma, \delta) \in BC_1 \times \omega^\omega$ and $\{\delta\}^{X,Y}(x)$ is defined, then

$$\begin{aligned} x \in \rho^X[u_r^{X,Y}(\gamma, \delta)] &\Leftrightarrow \exists k \in \omega \ x \in N[X, g(\gamma, \delta)^*(k)] \\ &\Leftrightarrow \exists k \in \omega \ x \in N\left[X, \left((k)_0\right)_0\right] \text{ and } \left[\delta, \gamma^*[(k)_1], \left((k)_0\right)_0, \left((k)_0\right)_1\right] \in P^* \\ &\Leftrightarrow \exists j \in \omega \ \exists i \in \omega \ \left(x \in N[X, (i)_0] \text{ and } [\delta, \gamma^*(j), (i)_0, (i)_1] \in P^*\right) \\ &\Leftrightarrow \exists j \in \omega \ [\delta, x, \gamma^*(j)] \in P \\ &\Leftrightarrow \exists j \in \omega \ \{\delta\}^{X,Y}(x) \in N[Y, \gamma^*(j)] \\ &\Leftrightarrow \{\delta\}^{X,Y}(x) \in \rho^Y(\gamma). \end{aligned}$$

- Now let $\gamma \in BC_\xi$ with $\xi \geq 2$, and $\delta \in \omega^\omega$. As in the proof of Lemma 3.2, $u_r^{X,Y}(\gamma, \delta) \in BC_\xi$. If $\{\delta\}^{X,Y}(x)$ is defined, then

$$x \in \rho^X\left(u_r^{X,Y}[\{\gamma^*\}(i), \delta]\right) \Leftrightarrow \{\delta\}^{X,Y}(x) \in \rho^Y[\{\gamma^*\}(i)],$$

by induction assumption. This shows that $x \in \rho^X[u_r^{X,Y}(\gamma, \delta)] \Leftrightarrow \{\delta\}^{X,Y}(x) \in \rho^Y(\gamma)$. \square

As a corollary, one can prove the uniform closure of the pointclasses Σ_ξ^0 under fixations of recursive arguments. It is sometimes convenient to “view a code in BC_1 as an element of BC_2 ”, even it is not formally correct. The next lemma expresses this:

Lemma 3.4 *Let X be a recursively presented Polish space. Then there is $u_*^X: \omega^\omega \rightarrow \omega^\omega$ recursive such that for each $\gamma \in BC_1$ (resp., BC^*), $u_*^X(\gamma) \in BC_2$ (resp., $u_*^X(\gamma) = \gamma$), and $\rho^X[u_*^X(\gamma)] = \rho^X(\gamma)$.*

Proof. We define $R \in \Sigma_1^0(\omega^\omega \times X)$ by $(\gamma, x) \in R \Leftrightarrow \exists i \in \omega \ x \in N[X, \gamma^*(i)]$. As $R \in \Sigma_2^0$ there is $C \in \Pi_1^0(\omega \times \omega^\omega \times X)$ such that $R = \exists^\omega C$. Let $\varepsilon_0 \in \Sigma_1^0$ such that $\neg C = \bigcup_{i \in \omega} N[\omega \times \omega^\omega \times X, \varepsilon_0(i)]$. Note that $0 \frown \varepsilon_0 \in \Sigma_1^0 \cap BC_1$ and $\neg C = \rho^{\omega \times \omega^\omega \times X}(0 \frown \varepsilon_0)$. Using Lemma 3.2, we see the existence of $\gamma_0 \in \Sigma_1^0$ such that $\{\gamma_0\}(\gamma, i) = u_s^{\omega \times \omega^\omega}(0 \frown \varepsilon_0, i, \gamma)$ for each $(\gamma, i) \in \omega^\omega \times \omega$. Then we define $u_*^X(\gamma) := 1 \frown S_{\Sigma_1^0}^{\omega^\omega, \omega, \omega^\omega}(\gamma_0, \gamma)$ if $\gamma \in BC_1$, γ otherwise. \square

We now prove another lemma stating, among other things, that the pointclasses Σ_ξ^0 are uniformly closed under finite intersections and unions:

Lemma 3.5 *Let X be a recursively presented Polish space. There is $u_f^X: 2 \times \omega \times \omega^\omega \rightarrow \omega^\omega$ recursive such that for each $(\xi, a, n, \gamma) \in (\omega_1 \setminus \{0\}) \times 2 \times \omega \times \omega^\omega$,*

(a) *If $(\gamma)_i \in BC_1 \cup BC_\xi$ for each $i \leq n$, then $u_f^X(a, n, \gamma)$ is in $BC_1 \cup BC_\xi$. Moreover, the equalities $\rho^X[u_f^X(0, n, \gamma)] = \bigcap_{i \leq n} \rho^X[(\gamma)_i]$ and $\rho^X[u_f^X(1, n, \gamma)] = \bigcup_{i \leq n} \rho^X[(\gamma)_i]$ hold.*

(b) *If moreover $\xi \geq 2$ and $(\gamma)_i \in BC_\xi$ for some $i \leq n$, then $u_f^X(a, n, \gamma)$ is in BC_ξ .*

Proof. For $\xi = 1$, by 3B.2 in [Mos80] there is $f: \omega^3 \rightarrow \omega$ recursive such that, for $(u, n) \in \omega^2$,

$$\bigcap_{i \leq n} N[X, (u)_i] = \bigcup_{m \in \omega} N[X, f(u, n, m)].$$

We set $g(a, n, \gamma)(0) := 0$ and

$$g(0, n, \gamma)(i+1) := f\left(\langle [(\gamma)_0]^*[(i)_0], \dots, [(\gamma)_n]^*[(i)_n] \rangle, n, (i)_1\right),$$

$$g(1, n, \gamma)(i+1) := \begin{cases} [(\gamma)_{(i)_0}]^*[(i)_1] & \text{if } (i)_0 \leq n, \\ 0 & \text{otherwise.} \end{cases}$$

Note that $g(a, n, \gamma) \in BC_1$. If $(\gamma)_i \in BC_1$ for each $i \leq n$, then we get

$$\begin{aligned} \rho^X[g(0, n, \gamma)] &= \bigcup_{i \in \omega} N[X, g(0, n, \gamma)(i+1)] \\ &= \bigcup_{(j, m) \in \omega^2} N\left[X, f\left(\langle [(\gamma)_0]^*[(j)_0], \dots, [(\gamma)_n]^*[(j)_n] \rangle, n, m\right)\right] \\ &= \bigcup_{j \in \omega} \bigcap_{i \leq n} N\left(X, [(\gamma)_i]^*[(j)_i]\right) \\ &= \bigcap_{i \leq n} \bigcup_{j \in \omega} N\left(X, [(\gamma)_i]^*(j)\right) \\ &= \bigcap_{i \leq n} \rho^X[(\gamma)_i]. \end{aligned}$$

Moreover,

$$\rho^X[g(1, n, \gamma)] = \bigcup_{i \in \omega} N[X, g(1, n, \gamma)(i+1)] = \bigcup_{i \leq n} \bigcup_{j \in \omega} N\left(X, [(\gamma)_i]^*(j)\right) = \bigcup_{i \leq n} \rho^X[(\gamma)_i].$$

• For the general case, using Lemma 3.4 we define a partial function $h: \omega^\omega \times \omega^2 \rightarrow \omega^\omega$ by

$$\left(h(\gamma, n, j)\right)_i := \begin{cases} \left\{ \left(u_*^X[(\gamma)_i]\right)^* \right\} [(j)_i] & \text{if } i \leq n, \\ 0^\infty & \text{otherwise.} \end{cases}$$

It allows us to define another partial function $\psi: \omega^\omega \times 2 \times \omega \times \omega^\omega \times \omega \rightarrow \omega^\omega$ by

$$\psi(\varepsilon, a, n, \gamma, j) := \{\varepsilon\}[1 - a, n, h(\gamma, n, j)].$$

We argue as in the proof of Lemma 3.2 to define ν and a recursive map $\varphi: \omega^\omega \times 2 \times \omega \times \omega^\omega \rightarrow \omega^\omega$ by

$$\varphi(\varepsilon, a, n, \gamma) := \begin{cases} g(a, n, \gamma) & \text{if } (\gamma)_i(0) = 0 \text{ for each } i \leq n, \\ 1 \frown S_{\Sigma_1^0}^{\omega^\omega \times 2 \times \omega \times \omega^\omega, \omega, \omega^\omega}(\nu, \varepsilon, a, n, \gamma) & \text{if } (\gamma)_i(0) \neq 0 \text{ for some } i \leq n. \end{cases}$$

By Kleene's Recursion Theorem, there is $\varepsilon^* \in \Sigma_1^0$ such that $\varphi(\varepsilon^*, a, n, \gamma) = \{\varepsilon^*\}(a, n, \gamma)$ for each (a, n, γ) in $2 \times \omega \times \omega^\omega$. We put $u_f^X(a, n, \gamma) := \{\varepsilon^*\}(a, n, \gamma)$. The map u_f^X is a total recursive map. We show that $u_f^X(a, n, \gamma)$ satisfies the required properties by induction on ξ .

• Assume that $(\gamma)_i \in BC_1$ for each $i \leq n$. We have $u_f^X(a, n, \gamma) = g(a, n, \gamma)$, so we are done, by the previous discussion. Assume now that $\xi \geq 2$, and that $(\gamma)_i \in BC_1 \cup BC_\xi$ for each $i \leq n$. We may assume that $(\gamma)_i \in BC_\xi$ for some $i \leq n$. Then $\left\{ \left(u_*^X[(\gamma)_i] \right)^* \right\}(k)$ is defined for each integer k . In particular, $h(\gamma, n, j)$ and $\psi(\varepsilon^*, a, n, \gamma, j)$ are defined for each (a, j) in $2 \times \omega$. Thus $\{u_f^X(a, n, \gamma)^*(j) = u_f^X[1-a, n, h(\gamma, n, j)]\}$ is defined for each integer j . As $\left\{ \left(u_*^X[(\gamma)_i] \right)^* \right\}(k)$ is in some BC_{η_k} with $1 \leq \eta_k < \xi$ for each integer k , there is $1 \leq \eta < \xi$ such that $\left(h(\gamma, n, j) \right)_i$ is in $BC_1 \cup BC_\eta$ for each $i \leq n$. By induction assumption, we get $u_f^X[1-a, n, h(\gamma, n, j)] \in BC_1 \cup BC_\eta$. This shows that $u_f^X(a, n, \gamma) \in BC_\xi$. Moreover, by induction assumption we get

$$\begin{aligned}
\rho^X[u_f^X(0, n, \gamma)] &= \bigcup_{j \in \omega} \neg \bigcup_{i \leq n} \rho^X \left[\left(h(\gamma, n, j) \right)_i \right] \\
&= \bigcup_{j \in \omega} \bigcap_{i \leq n} \neg \rho^X \left[\left\{ \left(u_*^X[(\gamma)_i] \right)^* \right\} [(j)_i] \right] \\
&= \bigcap_{i \leq n} \bigcup_{j \in \omega} \neg \rho^X \left[\left\{ \left(u_*^X[(\gamma)_i] \right)^* \right\} (j) \right] \\
&= \bigcap_{i \leq n} \rho^X \left(u_*^X[(\gamma)_i] \right) \\
&= \bigcap_{i \leq n} \rho^X[(\gamma)_i].
\end{aligned}$$

Similarly, we get $\rho^X[u_f^X(1, n, \gamma)] = \bigcup_{i \leq n} \rho^X[(\gamma)_i]$. □

In the sequel we will need a last closure property, asserting, among other things, that the pointclasses Σ_ξ^0 are uniformly closed under \exists^ω :

Lemma 3.6 (a) *There is a recursive map $u_\exists: \omega^\omega \rightarrow \omega^\omega$ such that for each $1 \leq \xi < \omega_1$ and for each $\gamma \in BC_\xi$, $u_\exists(\gamma) \in BC_\xi$, and $x \in \rho^X[u_\exists(\gamma)] \Leftrightarrow \exists n \in \omega \ (n, x) \in \rho^{\omega \times X}(\gamma)$, for each recursively presented Polish space X and for each $x \in X$.*

(b) *There is a recursive map $u_{(\cdot)}: \omega^\omega \rightarrow \omega^\omega$ such that for each $1 \leq \xi < \omega_1$, $(\gamma)_n \in BC_\xi$ for each $n \in \omega$ implies that $u_{(\cdot)}(\gamma) \in BC_\xi$, and $x \in \rho^X[u_{(\cdot)}(\gamma)] \Leftrightarrow \exists n \in \omega \ x \in \rho^X[(\gamma)_n]$ for each recursively presented Polish space X and for each $x \in X$.*

Proof. (a) By 3B.1 in [Mos80], there are g and h recursive such that $N(\omega \times X, k) = N[\omega, g(k)] \times N[X, h(k)]$ for each integer k . If $\gamma(0) = 0$, then we put

$$u_\exists(\gamma)(j) := \begin{cases} h\left(\gamma^*[(j-1)_0]\right) & \text{if } j > 0 \text{ and } (j-1)_1 \in N\left[\omega, g\left(\gamma^*[(j-1)_0]\right)\right], \\ 0 & \text{otherwise.} \end{cases}$$

Using Lemma 3.2, we define a partial function $f: \omega^\omega \times \omega \rightarrow \omega^\omega$ by

$$f(\gamma, i) := u_s^\omega\left(\{\gamma^*\}[(i)_1], (i)_0\right).$$

As f is recursive on its domain, there is $\varepsilon_0 \in \Sigma_1^0$ such that $f(\gamma, i) = \{\varepsilon_0\}(\gamma, i)$ if $f(\gamma, i)$ is defined. If $\gamma(0) \neq 0$, then we put $u_{\exists}(\gamma) := 1 \frown S_{\Sigma_1^0}^{\omega, \omega, \omega}(\varepsilon_0, \gamma)$. This defines a recursive map u_{\exists} . If $\gamma(0) \neq 0$ and $\gamma \in BC_{\xi}$, then $\{[u_{\exists}(\gamma)]^*\}(i) = \{S_{\Sigma_1^0}^{\omega, \omega, \omega}(\varepsilon_0, \gamma)\}(i) = \{\varepsilon_0\}(\gamma, i) = f(\gamma, i)$. Thus $u_{\exists}(\gamma) \in BC_{\xi}$, even if $\gamma(0) = 0$. Let $x \in X$. If $\gamma(0) = 0$, then

$$\begin{aligned} \exists n \in \omega \ (n, x) \in \rho^{\omega \times X}(\gamma) &\Leftrightarrow \exists n \in \omega \ \exists p \in \omega \ (n, x) \in N[\omega \times X, \gamma^*(p)] \\ &\Leftrightarrow \exists n \in \omega \ \exists p \in \omega \ n \in N(\omega, g[\gamma^*(p)]) \text{ and } x \in N(X, h[\gamma^*(p)]) \\ &\Leftrightarrow \exists i \in \omega \ (i)_1 \in N[\omega, g(\gamma^*[(i)_0])] \text{ and } x \in N[X, h(\gamma^*[(i)_0])] \\ &\Leftrightarrow x \in \rho^X[u_{\exists}(\gamma)]. \end{aligned}$$

If $\gamma(0) \neq 0$, then

$$\begin{aligned} \exists n \in \omega \ (n, x) \in \rho^{\omega \times X}(\gamma) &\Leftrightarrow \exists n \in \omega \ \exists p \in \omega \ (n, x) \notin \rho^{\omega \times X}[\{\gamma^*\}(p)] \\ &\Leftrightarrow \exists i \in \omega \ \neg [(i)_0, x] \in \rho^{\omega \times X}(\{\gamma^*\}[(i)_1]) \\ &\Leftrightarrow \exists i \in \omega \ \neg x \in \rho^X[f(\gamma, i)] \\ &\Leftrightarrow x \in \rho^X[u_{\exists}(\gamma)]. \end{aligned}$$

(b) If $(\gamma)_0(0) = 0$, then we put

$$u_{(\cdot)}(\gamma)(j) := \begin{cases} 0 & \text{if } j = 0, \\ [(\gamma)_{(j-1)_0}]^*[(j-1)_1] & \text{otherwise.} \end{cases}$$

We define a partial function $f' : \omega^{\omega} \times \omega \rightarrow \omega^{\omega}$ by $f'(\gamma, i) := \{[(\gamma)_{(i)_0}]^*\}[(i)_1]$. As f' is recursive on its domain, there is $\gamma_0 \in \Sigma_1^0$ such that $f'(\gamma, i) = \{\gamma_0\}(\gamma, i)$ if $f'(\gamma, i)$ is defined. If $(\gamma)_0(0) \neq 0$, then we put $u_{(\cdot)}(\gamma) := 1 \frown S_{\Sigma_1^0}^{\omega, \omega, \omega}(\gamma_0, \gamma)$. This defines a recursive map $u_{(\cdot)}$. If $\xi \geq 2$ and $(\gamma)_n \in BC_{\xi}$ for each integer n , then $\{[u_{(\cdot)}(\gamma)]^*\}(i) = \{S_{\Sigma_1^0}^{\omega, \omega, \omega}(\gamma_0, \gamma)\}(i) = \{\gamma_0\}(\gamma, i) = f'(\gamma, i)$. Thus $u_{(\cdot)}(\gamma) \in BC_{\xi}$, even if $\xi = 1$. Let $x \in X$. If $(\gamma)_0(0) = 0$, then

$$\begin{aligned} \exists n \in \omega \ x \in \rho^X[(\gamma)_n] &\Leftrightarrow \exists n \in \omega \ \exists p \in \omega \ x \in N(X, [(\gamma)_n]^*(p)) \\ &\Leftrightarrow \exists i \in \omega \ x \in N(X, [(\gamma)_{(i)_0}]^*[(i)_1]) \\ &\Leftrightarrow x \in \rho^X[u_{(\cdot)}(\gamma)]. \end{aligned}$$

If $(\gamma)_0(0) \neq 0$, then

$$\begin{aligned} \exists n \in \omega \ x \in \rho^X[(\gamma)_n] &\Leftrightarrow \exists n \in \omega \ \exists p \in \omega \ x \notin \rho^X(\{[(\gamma)_n]^*\}(p)) \\ &\Leftrightarrow \exists i \in \omega \ \neg x \in \rho^X(\{[(\gamma)_{(i)_0}]^*\}[(i)_1]) \\ &\Leftrightarrow \exists i \in \omega \ \neg x \in \rho^X[f'(\gamma, i)] \\ &\Leftrightarrow x \in \rho^X[u_{(\cdot)}(\gamma)]. \end{aligned}$$

This finishes the proof. □

The hyperarithmetical hierarchy.

The notion of a hyperarithmetical set is defined in [Mos80] (see 7B): a subset of a recursively presented Polish space is hyperarithmetical if it is Borel and has a recursive Borel code. We can define a hyperarithmetical hierarchy, extending the arithmetical hierarchy. The following characterization of the arithmetical pointclasses Σ_n^0 can be found in [Lou??]:

Theorem 3.7 *Let X be a recursively presented Polish space, and $n \geq 1$ an integer. Then*

$$\Sigma_n^0(X) = \{\rho^X(\gamma) \mid \gamma \in \Sigma_1^0 \cap BC_n\}.$$

Actually, we will use only a small part of it. More specifically, we will only use the fact that if P is $\Sigma_1^0(X)$, then there is $\gamma \in \Sigma_1^0 \cap BC_1$ with $P = \rho^X(\gamma)$. It is very simple: there is $\varepsilon \in \Sigma_1^0$ such that $P = \bigcup_{i \in \omega} N[X, \varepsilon(i)] = \rho^X(0 \smallfrown \varepsilon)$. Thus $\gamma := 0 \smallfrown \varepsilon$ is suitable. The following definition comes naturally after Theorem 3.7, and can be found in [Lou??]:

Definition 3.8 *Let X be a recursively presented Polish space, and $1 \leq \xi < \omega_1$. Then we set*

$$\Sigma_\xi^0(X) = \{\rho^X(\gamma) \mid \gamma \in \Sigma_1^0 \cap BC_\xi\},$$

$$\Pi_\xi^0(X) = \check{\Sigma}_\xi^0(X),$$

$$\Delta_\xi^0(X) = \Sigma_\xi^0(X) \cap \Pi_\xi^0(X).$$

This defines the hyperarithmetical hierarchy.

Note that Lemma 3.3 (resp., 3.5, 3.6) implies that the hyperarithmetical pointclasses are closed under recursive substitutions (resp., finite intersections and unions, \exists^ω). Now we construct recursive maps giving codes for the basic neighborhoods and their complement in spaces of type at most 1.

Lemma 3.9 *Let X be a recursively presented Polish space of type at most 1.*

(a) *There is a recursive map $u_N : \omega \rightarrow \omega^\omega$ such that $u_N(k) \in BC_1$, and $\rho^X[u_N(k)] = N(X, k)$.*

(b) *There is a recursive map $u_{\neg N}^X : \omega \rightarrow \omega^\omega$ such that $u_{\neg N}^X(k) \in BC_1$, and $\rho^X[u_{\neg N}^X(k)] = \neg N(X, k)$.*

Proof. (a) Put $u_N(k) := 0k0^\omega$.

(b) By 3C.3 in [Mos80], the equivalence $(x, k) \in R \Leftrightarrow x \notin N(X, k)$ defines $R \in \Sigma_1^0(X \times \omega)$. By Theorem 3.7 there is $\gamma_0 \in \Sigma_1^0 \cap BC_1$ with $R = \rho^{X \times \omega}(\gamma_0)$. Using Lemma 3.2 we set $u_{\neg N}^X(k) := u_s^\omega(\gamma_0, k)$. \square

Now we use this to prove that, uniformly in $\xi \geq 2$, a set in the pointclass $\Sigma_\xi^0(X)$ (resp., $\Sigma_\xi^0(X)$) is the disjoint union of sets in $\Pi_{<\xi}^0$ (resp., $\Pi_{<\xi}^0$), if X is a space of type at most 1. We will use the notation

$$E = \dot{\bigcup}_{i \in \omega} E_i$$

to express the fact that E is the disjoint union of the E_i 's.

Lemma 3.10 *Let X be a recursively presented Polish space of type at most 1. Then there is a recursive map $u_d^X : \omega^\omega \rightarrow \omega^\omega$ such that $u_d^X(\gamma) \in BC_\xi$ if $\gamma \in BC_\xi$, for each $1 \leq \xi < \omega_1$. Moreover,*

(a) *There is a recursive map $u_c^X : \omega^\omega \times \omega \rightarrow \omega^\omega$ such that*

(1) $u_c^X(\gamma, i) \in BC_1$ for each $(\gamma, i) \in \omega^\omega \times \omega$.

(2) $\{[u_d^X(\gamma)]^*(i)\}$ is defined, in BC_1 , $\rho^X(\{[u_d^X(\gamma)]^*(i)\}) \in \Delta_1^0$ and $\neg\rho^X(\{[u_d^X(\gamma)]^*(i)\}) = \rho^X[u_c^X(\gamma, i)]$ for each $(\gamma, i) \in BC_1 \times \omega$.

(b) If $1 \leq \xi < \omega_1$ and $\gamma \in BC_\xi$, then $\rho^X(\gamma) = \bigcup_{i \in \omega} \neg\rho^X(\{[u_d^X(\gamma)]^*(i)\})$.

Proof. For $\xi = 1$, we first define a map $\tilde{f} : \omega^\omega \times \omega \rightarrow \omega^\omega$, using Lemma 3.9, as follows:

$$\left(\tilde{f}(\gamma, i)\right)_j := \begin{cases} u_{-N}^X[\gamma^*(j)] & \text{if } j < i, \\ u_N[\gamma^*(i)] & \text{if } j \geq i. \end{cases}$$

As \tilde{f} is recursive, the formula $u_c^X(\gamma, i) := u_f^X[0, i, \tilde{f}(\gamma, i)]$ defines u_c^X recursive such that $u_c^X(\gamma, i) \in BC_1$ for each $(\gamma, i) \in \omega^\omega \times \omega$ (see Lemma 3.5). Then, using Lemma 3.9, we define a map $f : \omega^\omega \times \omega \rightarrow \omega^\omega$:

$$\left(f(\gamma, i)\right)_j := \begin{cases} u_N[\gamma^*(j)] & \text{if } j < i, \\ u_{-N}^X[\gamma^*(i)] & \text{if } j \geq i. \end{cases}$$

As f is recursive, and using Lemma 3.5, there is $\varepsilon_0 \in \Sigma_1^0$ such that $\{\varepsilon_0\}(\gamma, i) = u_f^X[1, i, f(\gamma, i)] \in BC_1$ for each $(\gamma, i) \in \omega^\omega \times \omega$. We define a recursive map $g : \omega^\omega \rightarrow \omega^\omega$ by $g(\gamma) := 0 \frown S_{\Sigma_1^0}^{\omega^\omega, \omega^\omega}(\varepsilon_0, \gamma)$. If $(\gamma, i) \in BC_1 \times \omega$, then $\{[g(\gamma)]^*(i)\} = \{S_{\Sigma_1^0}^{\omega^\omega, \omega^\omega}(\varepsilon_0, \gamma)\}(i) = \{\varepsilon_0\}(\gamma, i) = u_f^X[1, i, f(\gamma, i)]$ is defined, $\rho^X(\{[g(\gamma)]^*(i)\}) \in \Delta_1^0$ since it is a finite union of clopen sets, and $\neg\rho^X(\{[g(\gamma)]^*(i)\}) = \rho^X[u_c^X(\gamma, i)]$. Moreover,

$$\begin{aligned} \rho^X(\gamma) &= \bigcup_{i \in \omega} N[X, \gamma^*(i)] = \bigcup_{i \in \omega} N[X, \gamma^*(i)] \setminus \left(\bigcup_{j < i} N[X, \gamma^*(j)] \right) \\ &= \bigcup_{i \in \omega} \neg \left[\rho^X(u_{-N}^X[\gamma^*(i)]) \cup \bigcup_{j < i} \rho^X(u_N[\gamma^*(j)]) \right] \\ &= \bigcup_{i \in \omega} \neg \rho^X(u_f^X[1, i, f(\gamma, i)]) = \bigcup_{i \in \omega} \neg \rho^X(\{[g(\gamma)]^*(i)\}). \end{aligned}$$

• For the general case, assume that $\gamma \in BC_\xi$, with $\xi \geq 2$. We set $B_j := \rho^X[\{\gamma^*\}(j)]$, so that we can write $\rho^X(\gamma) = \bigcup_{j \in \omega} \neg B_j$. Note that $\{\gamma^*\}(j) \in BC_{\eta_j}$, where $1 \leq \eta_j < \xi$. We set

$$B_{j,i} := \begin{cases} \neg N(X, [\{\gamma^*\}(j)]^*(i)) & \text{if } [\{\gamma^*\}(j)](0) = 0, \\ \rho^X(\{[\{\gamma^*\}(j)]^*(i)\}) & \text{otherwise,} \end{cases}$$

so that $B_j := \bigcup_{i \in \omega} \neg B_{j,i}$.

By Lemma 3.9, $B_{j,i} = \rho^X \left[u_{-N}^X \left([\{\gamma^*\}(j)]^*(i) \right) \right]$ if $[\{\gamma^*\}(j)](0) = 0$, and

$$\neg B_{j,i} = \begin{cases} \rho^X \left[u_N \left([\{\gamma^*\}(j)]^*(i) \right) \right] & \text{if } [\{\gamma^*\}(j)](0) = 0, \\ \rho^X \left[u_{-} \left(\left\{ [\{\gamma^*\}(j)]^* \right\} (i) \right) \right] & \text{otherwise,} \end{cases}$$

by Lemma 3.1. Thus

$$\begin{aligned} \rho^X(\gamma) &= \bigcup_{k \in \omega} \neg B_k \\ &= \bigcup_{k \in \omega} \bigcap_{j < k} B_j \setminus B_k \\ &= \bigcup_{k \in \omega} \bigcap_{j < k} \left(\bigcup_{i \in \omega} \neg B_{j,i} \right) \setminus B_k \\ &= \bigcup_{k \in \omega} \bigcap_{j < k} \left(\bigcup_{i \in \omega} \bigcap_{l < i} B_{j,l} \setminus B_{j,i} \right) \setminus B_k \\ &= \bigcup_{i \in \omega} \left(\bigcap_{j < lh(i)} \bigcap_{l < (i)_j} B_{j,l} \setminus B_{j,(i)_j} \right) \setminus B_{lh(i)} \\ &= \bigcup_{i \in \omega} \neg \left(B_{lh(i)} \cup \bigcup_{j < lh(i)} \bigcup_{l < (i)_j} B_{j,(i)_j} \cup \neg B_{j,l} \right). \end{aligned}$$

Note that the code for $B_{i,j}$ is a partial recursive function of γ, i and j . Using Lemma 3.5, this shows the existence of a partial function $f^X : \omega^\omega \times \omega \rightarrow \omega^\omega$, recursive on its domain, such that $f^X(\gamma, i)$ is in $\bigcup_{1 \leq \eta < \xi} BC_\eta$ and $\rho^X(\gamma) = \bigcup_{i \in \omega} \neg \rho^X[f^X(\gamma, i)]$ for each $\gamma \in BC_\xi$ with $\xi \geq 2$. There is $\varepsilon_1 \in \Sigma_1^0$ such that $f^X(\gamma, i) = \{\varepsilon_1\}(\gamma, i) = \{S_{\Sigma_1^0}^{\omega^\omega, \omega, \omega^\omega}(\varepsilon_1, \gamma)\}(i)$ if $f^X(\gamma, i)$ is defined. We define $h : \omega^\omega \rightarrow \omega^\omega$ by the formula $h(\gamma) := 1 \frown S_{\Sigma_1^0}^{\omega^\omega, \omega, \omega^\omega}(\varepsilon_1, \gamma)$. The map h is recursive, $h(\gamma) \in BC_\xi$ and

$$\rho^X(\gamma) = \bigcup_{i \in \omega} \neg \rho^X \left(\left\{ [h(\gamma)]^* \right\} (i) \right)$$

if $\gamma \in BC_\xi$ and $\xi \geq 2$.

- It remains to set $u_d^X(\gamma) := g(\gamma)$ if $\gamma(0) = 0$, $h(\gamma)$ otherwise. □

Now we will show that the hyperarithmetical hierarchy makes sense, i.e., the existence of sets of arbitrary complexity under ω_1^{CK} . The intuition is quite simple: we take universal sets. But we have to check that this is effective.

Notation. Recall that if $\alpha \in \omega^\omega$, then $\leq_\alpha := \{(m, n) \in \omega^2 \mid \alpha(\langle m, n \rangle) = 1\}$ (see 4A in [Mos80]), and $<_\alpha := \{(m, n) \in \omega^2 \mid \alpha(\langle m, n \rangle) = 1 \text{ and } \alpha(\langle n, m \rangle) \neq 1\}$. The first relation is used to define the set $WO := \{\alpha \in \omega^\omega \mid \leq_\alpha \text{ is a wellordering on its domain } \{n \in \omega \mid n \leq_\alpha n\}\}$, which is used to define

$$\omega_1^{CK} := \sup\{|\alpha| \mid \alpha \in WO \cap \Sigma_1^0\},$$

where $|\alpha|$ is the order type of \leq_α . If $\alpha \in \omega^\omega$ and $p \in \omega$, then we define $\alpha|_p \in 2^\omega \subseteq \omega^\omega$ by

$$\alpha|_p(q) = 1 \Leftrightarrow \text{Seq}(q) \text{ and } \text{lh}(q) = 2 \text{ and } \alpha(q) = 1 \text{ and } \forall i \in 2 (q)_i <_\alpha p.$$

If $\alpha \in WO$, then $\alpha|_p \in WO$ and $\leq_{\alpha|_p}$ is the restriction of \leq_α to the strict \leq_α -predecessors of p .

Lemma 3.11 *There is a partial function $\eta : \omega^\omega \times \omega \rightarrow \omega^\omega$, recursive on its domain, defined if $\alpha \in WO$ and $|\alpha|_p \geq 1$, such that $|\alpha|_p = \sup \uparrow \{|\alpha|_{\eta(\alpha, p)(n)}| + 1 \mid n \in \omega\}$.*

Proof. We define a partial function $g : \omega^\omega \times \omega \rightarrow \omega$ by $g(\alpha, p) := \min\{m \in \omega \mid m <_\alpha p\}$ if it exists. Note that g is recursive on its domain and defined on $D := \{(\alpha, p) \in WO \times \omega \mid |\alpha|_p \geq 1\}$. We define a map $h : \omega^\omega \times \omega^3 \rightarrow \omega$ by

$$h(\alpha, p, n, m) := \begin{cases} n & \text{if } m <_\alpha n <_\alpha p, \\ m & \text{otherwise.} \end{cases}$$

Note that h is recursive. This allows us to define a partial function $\psi : (\omega^\omega)^2 \times \omega^2 \rightarrow \omega$ by:

$$\psi(\varepsilon, \alpha, p, n) := \begin{cases} g(\alpha, p) & \text{if } n = 0, \\ h[\alpha, p, n-1, \{\varepsilon\}_{\Sigma_1^0}^{\omega^\omega \times \omega^2, \omega}(\alpha, p, n-1)] & \text{if } n \geq 1. \end{cases}$$

Note that ψ is recursive on its domain, so that there is $\varepsilon^* \in \Sigma_1^0$ such that $\{\varepsilon^*\}(\alpha, p, n) = \psi(\varepsilon^*, \alpha, p, n)$ if $\psi(\varepsilon^*, \alpha, p, n)$ is defined. Now it is clear that $\psi(\varepsilon^*, \alpha, p, n)$ is defined if $(\alpha, p) \in D$, by induction on n , and that $|\alpha|_p = \sup \uparrow \{|\alpha|_{\{\varepsilon^*\}(\alpha, p, n)}| + 1 \mid n \in \omega\}$. We put $\eta(\alpha, p)(n) := \{\varepsilon^*\}(\alpha, p, n)$ if $\psi(\varepsilon^*, \alpha, p, n)$ is defined. Clearly, η is defined on D and suitable. \square

Notation. In the next lemma we identify $(\omega^\omega)^\omega$ with ω^ω , using the formula $\left((\delta_q)_{q \in \omega} \right)_n = \delta_n$. Let $\alpha \in WO$, $\gamma_0 \in \omega^\omega$, and $u : \omega^\omega \rightarrow \omega^\omega$ a map. Using Lemma 3.11 we can define, by induction on p (with respect to the wellordering \leq_α), and if $|\alpha|_p \geq 1$, $\gamma_{|\alpha|_p} := u[(\gamma_{|\alpha|_{\eta(\alpha, p)(n)}})_{n \in \omega}]$.

Lemma 3.12 *Let $\theta < \omega_1^{CK}$, $\alpha \in WO \cap \Sigma_1^0$ with $\theta + 1 = |\alpha|$, $\gamma_0 \in \Sigma_1^0$, $u : \omega^\omega \rightarrow \omega^\omega$ a recursive map, and $p \in \omega$ such that $p \leq_\alpha p$. Then $\gamma_{|\alpha|_p}$ is Σ_1^0 .*

Proof. Fix $p_0 \in \omega$ with $|\alpha|_{p_0} = 0$. Using Lemma 3.11, we define a partial function $f : \omega^\omega \times \omega \rightarrow \omega^\omega$ by $f(\varepsilon, p) := u \left[\left(\{\varepsilon\}[\eta(\alpha, p)(n)] \right)_{n \in \omega} \right]$. Note that f is recursive on its domain. We define a partial function $\psi : \omega^\omega \times \omega \rightarrow \omega^\omega$ by

$$\psi(\varepsilon, p) := \begin{cases} \gamma_0 & \text{if } p = p_0, \\ f(\varepsilon, p) & \text{if } p \neq p_0. \end{cases}$$

As ψ is recursive on its domain, there is $\varepsilon^* \in \Sigma_1^0$ with $\{\varepsilon^*\}(p) = \psi(\varepsilon^*, p)$ if $\psi(\varepsilon^*, p)$ is defined. It remains to see that $\psi(\varepsilon^*, p)$ is defined and equal to $\gamma_{|\alpha|_p}$ if $p \leq \alpha$. We argue by induction on p (with respect to the wellordering \leq_α). If $p = p_0$, then $\psi(\varepsilon^*, p) = \gamma_0 = \gamma_{|\alpha|_{p_0}} = \gamma_{|\alpha|_p}$. Assume now that $|\alpha|_p \geq 1$, and that the statement is proved for q satisfying $|\alpha|_q < |\alpha|_p$. Then $\psi[\varepsilon^*, \eta(\alpha, p)(n)]$ is defined and equal to $\gamma_{|\alpha|_{\eta(\alpha, p)(n)}}$ for each $n \in \omega$. It is also equal to $\{\varepsilon^*\}[\eta(\alpha, p)(n)]$. Thus $f(\varepsilon^*, p)$ is defined and equal to $u[(\gamma_{|\alpha|_{\eta(\alpha, p)(n)}})_{n \in \omega}] = \gamma_{|\alpha|_p}$. \square

Notation. In the next lemma, $\alpha \in WO$. We denote $\eta_{\alpha, p} := |\alpha|_p$, and $\eta_{\alpha, p, n} := |\alpha|_{\eta(\alpha, p)(n)}$ if $|\alpha|_p \geq 1$.

Lemma 3.13 *There is $u : \omega^\omega \rightarrow \omega^\omega$ recursive such that $(\gamma)_n \in BC_{1+\eta_{\alpha, p, n}}$ for each integer n implies that $u(\gamma) \in BC_{1+\eta_{\alpha, p}}$ and $(\beta, \delta) \in \rho^{(2^\omega)^2}[u(\gamma)] \Leftrightarrow \exists n \in \omega [(\beta)_n, \delta] \notin \rho^{(2^\omega)^2}[(\gamma)_n]$.*

Proof. First note that there is $\varepsilon_0 \in \Sigma_1^0$ with $\{\varepsilon_0\}^{\omega \times (2^\omega)^2, (2^\omega)^2}(n, \beta, \delta) = [(\beta)_n, \delta]$ for each $(n, \beta, \delta) \in \omega \times (2^\omega)^2$. Similarly, using Lemmas 3.2 and 3.3, we see that there is $\varepsilon_1 \in \Sigma_1^0$ such that, for each $(\gamma, n) \in \omega^\omega \times \omega$, $\{\varepsilon_1\}(\gamma, n) = u_s^\omega \left(u_r^{\omega \times (2^\omega)^2, (2^\omega)^2}[(\gamma)_n, \varepsilon_0], n \right)$. We put $u(\gamma) := 1 \frown S_{\Sigma_1^0}^{\omega^\omega, \omega^\omega}(\varepsilon_1, \gamma)$, so that u is a recursive map. Moreover, $\{[u(\gamma)]^*\}(n) = \{\varepsilon_1\}(\gamma, n) = u_s^\omega \left(u_r^{\omega \times (2^\omega)^2, (2^\omega)^2}[(\gamma)_n, \varepsilon_0], n \right)$ is defined and in $BC_{1+\eta_{\alpha, p, n}}$, so that $u(\gamma) \in BC_{1+\eta_{\alpha, p}}$. Finally,

$$\begin{aligned} (\beta, \delta) \in \rho^{(2^\omega)^2}[u(\gamma)] &\Leftrightarrow \exists n \in \omega \quad (\beta, \delta) \notin \rho^{(2^\omega)^2}[\{[u(\gamma)]^*\}(n)] \\ &\Leftrightarrow \exists n \in \omega \quad (\beta, \delta) \notin \rho^{(2^\omega)^2} \left[u_s^\omega \left(u_r^{\omega \times (2^\omega)^2, (2^\omega)^2}[(\gamma)_n, \varepsilon_0], n \right) \right] \\ &\Leftrightarrow \exists n \in \omega \quad (n, \beta, \delta) \notin \rho^{\omega \times (2^\omega)^2} \left(u_r^{\omega \times (2^\omega)^2, (2^\omega)^2}[(\gamma)_n, \varepsilon_0] \right) \\ &\Leftrightarrow \exists n \in \omega \quad [(\beta)_n, \delta] \notin \rho^{(2^\omega)^2}[(\gamma)_n]. \end{aligned}$$

This finishes the proof. \square

Theorem 3.14 *Let $1 \leq \xi < \omega_1^{CK}$, and Γ be one of the classes Σ_ξ^0, Π_ξ^0 . Then there is $B_\xi \in \Gamma(2^\omega) \setminus \check{\Gamma}$.*

Proof. Assume first that $\Gamma = \Sigma_\xi^0$. As in 22.3 in [Kec95] we set

$$(\beta, \delta) \in \mathcal{U}_{\Sigma_1^0}^{2^\omega} \Leftrightarrow \exists k \in \omega \quad \beta(k) = 0 \text{ and } \delta \in N[2^\omega, k],$$

so that $\mathcal{U}_{\Sigma_1^0}^{2^\omega} \in \Sigma_1^0[(2^\omega)^2]$ is universal for $\Sigma_1^0(2^\omega)$. We define a recursive bijection $\psi : 2^\omega \rightarrow (2^\omega)^2$ by $\psi_i(\gamma)(k) := \gamma(2k + i)$, for $i \in 2$. We set $B_1 := \psi^{-1}(\mathcal{U}_{\Sigma_1^0}^{2^\omega})$, so that B_1 is Σ_1^0 . As in 22.4 in [Kec95], we see that $\mathcal{U}_{\Sigma_1^0}^{2^\omega} \notin \Pi_1^0$. Thus $B_1 \notin \Pi_1^0$ since ψ is a homeomorphism.

So we may assume that $\xi \geq 2$ and we will generalize this. Write $\xi = 1 + \theta$, with $1 \leq \theta < \omega_1^{CK}$. Let $\alpha \in WO \cap \Sigma_1^0$ with $\theta + 1 = |\alpha|$. Using the previous notation, we get $\eta_{\alpha, p} = \sup \uparrow \{\eta_{\alpha, p, n} + 1 \mid n \in \omega\}$ if $\eta_{\alpha, p} \geq 1$, by Lemma 3.11. As in 22.3 in [Kec95] again we inductively define, if $\eta_{\alpha, p} \geq 1$,

$$(\beta, \delta) \in \mathcal{U}_{\Sigma_{1+\eta_{\alpha, p}}^0}^{2^\omega} \Leftrightarrow \exists n \in \omega \quad [(\beta)_n, \delta] \notin \mathcal{U}_{\Sigma_{1+\eta_{\alpha, p, n}}^0}^{2^\omega},$$

so that $\mathcal{U}_{\Sigma_{1+\eta_{\alpha, p}}^0}^{2^\omega}$ is universal for $\Sigma_{1+\eta_{\alpha, p}}^0(2^\omega)$.

Note the existence of $q \in \omega$ with $\eta_{\alpha,q} = \theta$. As before we put $B_\xi := \psi^{-1}(\mathcal{U}_{\Sigma_\xi^0}^{2^\omega})$, so that B_ξ is not Π_ξ^0 . By Lemma 3.3, it remains to see that $\mathcal{U}_{\Sigma_{1+\eta_{\alpha,p}}^0}^{2^\omega}$ is $\Sigma_{1+\eta_{\alpha,p}}^0$. By Theorem 3.7 there is $\gamma_0 \in \Sigma_1^0 \cap BC_1$ such that $\mathcal{U}_{\Sigma_1^0}^{2^\omega} = \rho^{(2^\omega)^2}(\gamma_0)$. Lemma 3.13 gives u recursive. We can apply Lemma 3.12, so that $\gamma_{|\alpha|_p} \in \Sigma_1^0$ is defined for each p with $p \leq_\alpha p$. By induction we see that $\gamma_{\eta_{\alpha,p}} \in BC_{1+\eta_{\alpha,p}}$, by Lemma 3.13. Moreover,

$$(\beta, \delta) \in \rho^{(2^\omega)^2}(\gamma_{\eta_{\alpha,p}}) \Leftrightarrow (\beta, \delta) \in \rho^{(2^\omega)^2} \left(u[(\gamma_{\eta_{\alpha,p,n}})_{n \in \omega}] \right) \Leftrightarrow \exists n \in \omega [(\beta)_n, \delta] \notin \rho^{(2^\omega)^2}(\gamma_{\eta_{\alpha,p,n}}).$$

This inductively shows that $\rho^{(2^\omega)^2}(\gamma_{\eta_{\alpha,p}}) = \mathcal{U}_{\Sigma_{1+\eta_{\alpha,p}}^0}^{2^\omega}$. Thus $\mathcal{U}_{\Sigma_{1+\eta_{\alpha,p}}^0}^{2^\omega}$ is $\Sigma_{1+\eta_{\alpha,p}}^0$.

Assume now that $\Gamma = \Pi_\xi^0$. The previous facts give $B_\xi \in \Sigma_\xi^0(2^\omega) \setminus \Pi_\xi^0$. But it is clear that $A_\xi := \neg B_\xi$ is in $\Pi_\xi^0(2^\omega) \setminus \Sigma_\xi^0$. \square

Remark. We can define, for $\beta \in 2^\omega$, $\omega_1^\beta := \sup\{|\alpha| \mid \alpha \in WO \cap \Sigma_1^0(\beta)\}$. If X is a recursively presented Polish space, then we can define $\Sigma_\xi^0(\beta)(X) = \{\rho^X(\gamma) \mid \gamma \in \Sigma_1^0(\beta) \cap BC_\xi\}$, $\Pi_\xi^0(\beta) := \check{\Sigma}_\xi^0(\beta)$ and also $\Delta_\xi^0(\beta) := \Sigma_\xi^0(\beta) \cap \Pi_\xi^0(\beta)$. One can check that this definition of $\Sigma_1^0(\beta)$ is equivalent to the one we gave in section 3. The previous proof shows the existence of $B_\xi \in \Sigma_\xi^0(\beta)(2^\omega) \setminus \Pi_\xi^0$, for $1 \leq \xi < \omega_1^\beta$. Indeed, the only things to change in the proof are the following. In Lemma 3.12, $\theta < \omega_1^\beta$, $\alpha \in \Sigma_1^0(\beta)$, f and ψ become $\Sigma_1^0(\beta)$ on their domain by 3D.7, 3G.1 and 3G.2 in [Mos80]. Then we can apply 7A.2 in [Mos80] to get ε^* . The conclusion is that $\gamma_{|\alpha|_p} \in \Sigma_1^0(\beta)$. The result follows.

4 Effective versions of Kuratowski's theorem.

Notation. Let $\xi < \omega_1$. Then $\xi - 1$ will denote the predecessor of ξ if it exists, ξ otherwise. We also define $\xi^- := \xi - 1$ if $\xi \geq 3$, ξ otherwise.

Theorem 4.1 *Let $a \in 2$. There is a partial function $F^a : \omega^\omega \rightarrow (\omega^\omega)^3$, recursive on its domain, such that*

(a) *For each $1 \leq \xi \leq 2$ and for each $\gamma \in BC_\xi$, coding $B := \neg \rho^{2^\omega}(\gamma) \in \Pi_\xi^0$, $F^0(\gamma)$ is defined and*

(1) $F_0^0(\gamma) \in BC_1$ (codes $C := \neg \rho^{\omega^\omega}[F_0^0(\gamma)] \in \Pi_1^0$).

(2) $f := \{F_1^0(\gamma)\}_{|C}^{\omega^\omega, 2^\omega}$ defines a continuous bijection from C onto B .

(3) $F_2^0(\gamma) \in BC_1$ codes an open set computing a partial function $g : 2^\omega \rightarrow \omega^\omega$, defined and continuous on B , which coincides with f^{-1} .

(b) *For each $1 \leq \xi < \omega_1$ and for each $\gamma \in BC_\xi$, coding $B := \neg \rho^{2^\omega}(\gamma) \in \Pi_\xi^0$, $F^1(\gamma)$ is defined and*

(1) $F_0^1(\gamma) \in BC_1$ (codes $C := \neg \rho^{\omega^\omega}[F_0^1(\gamma)] \in \Pi_1^0$).

(2) $f := \{F_1^1(\gamma)\}_{|C}^{\omega^\omega, 2^\omega}$ defines a continuous bijection from C onto B .

(3) $F_2^1(\gamma) \in BC_{\xi^-}$ codes a $\Sigma_{\xi^-}^0$ set computing a partial function $g : 2^\omega \rightarrow \omega^\omega$, defined and $\Sigma_{\xi^-}^0$ -measurable on B , which coincides with f^{-1} .

Proof. Let us look at the case where $\xi = 1$ first. We define $\mu: \omega \rightarrow \omega$ by

$$\mu(k) := \begin{cases} 0 & \text{if } \binom{(k)_1}{1} = 0, \\ \min\{l \in \omega \mid \frac{1}{l+1} < \frac{\binom{(k)_1}{1}}{\binom{(k)_1}{2} + 1}\} & \text{otherwise.} \end{cases}$$

Clearly μ is recursive. Let us recall, for each $k \in \omega$, the definition of the basic neighborhood:

$$N(\omega^\omega, k) := \begin{cases} \emptyset & \text{if } \binom{(k)_1}{1} = 0, \\ \left\{ \delta \in \omega^\omega \mid \forall j < \mu(k) \delta(j) = \binom{\binom{(k)_1}{0}}{j} \right\} & \text{otherwise.} \end{cases}$$

We define a recursive map $\varphi: \omega \rightarrow \omega$ by

$$\varphi(k) := \begin{cases} 0 & \text{if } \binom{(k)_1}{1} = 0, \\ \left\langle \binom{\binom{(k)_1}{0}}{0}, \dots, \binom{\binom{(k)_1}{0}}{\mu(k)-1} \right\rangle & \text{otherwise.} \end{cases}$$

In 3A.2 in [Mos80] the recursive map $sg: \omega \rightarrow \omega$ is defined by $sg(n) := 0$ if $n = 0, 1$ otherwise. The recursive presentation of 2^ω ensures that

$$N(2^\omega, k) := \begin{cases} \emptyset & \text{if } \binom{(k)_1}{1} = 0, \\ \left\{ \alpha \in 2^\omega \mid \forall j < \mu(k) \alpha(j) = sg \left[\binom{\binom{(k)_1}{0}}{j} \right] \right\} & \text{otherwise.} \end{cases}$$

We view 2^ω as a subset of ω^ω . We denote by Id_{2^ω} the partial function defined on $2^\omega \subseteq \omega^\omega$, with values in 2^ω , by $\text{Id}_{2^\omega}(\alpha) := \alpha$. It is recursive on 2^ω , since the relation “ $\alpha \in N(2^\omega, k)$ ” is $\Sigma_1^0(\omega^\omega \times \omega)$ on $2^\omega \times \omega$. Thus there is $\delta_0 \in \Sigma_1^0$ with $\{\delta_0\}^{\omega^\omega, 2^\omega}(\alpha) = \text{Id}_{2^\omega}(\alpha)$ for each $\alpha \in 2^\omega$. By Lemma 3.3 we have $u_r^{\omega^\omega, 2^\omega}(\gamma, \delta_0) \in BC_1$ and $\alpha \in \rho^{\omega^\omega} [u_r^{\omega^\omega, 2^\omega}(\gamma, \delta_0)] \Leftrightarrow \alpha \in \rho^{2^\omega}(\gamma)$ if $\gamma \in BC_1$ and $\alpha \in 2^\omega$. As $2^\omega \in \Pi_1^0(\omega^\omega)$, there is $\gamma_0 \in \Sigma_1^0 \cap BC_1$ with $2^\omega = \neg \rho^{\omega^\omega}(\gamma_0)$, by Theorem 3.7. We define a recursive map $f: \omega^\omega \rightarrow \omega^\omega$ by $(f(\gamma))_i := \gamma_0$ if $i = 0$, $u_r^{\omega^\omega, 2^\omega}(\gamma, \delta_0)$ otherwise.

If $\gamma \in BC_1$, then using Lemma 3.5 we set $F_0^a(\gamma) := u_f^{\omega^\omega} [1, 1, f(\gamma)]$, so that $F_0^a(\gamma) \in BC_1$ and also $2^\omega \setminus \rho^{2^\omega}(\gamma) = \neg \rho^{\omega^\omega} [F_0^a(\gamma)]$ since

$$\rho^{\omega^\omega} [F_0^a(\gamma)] = \bigcup_{i \leq 1} \rho^{\omega^\omega} \left[(f(\gamma))_i \right] = \rho^{\omega^\omega}(\gamma_0) \cup \rho^{\omega^\omega} [u_r^{\omega^\omega, 2^\omega}(\gamma, \delta_0)] = \omega^\omega \setminus 2^\omega \cup \rho^{2^\omega}(\gamma).$$

Thus $B = 2^\omega \setminus \rho^{2^\omega}(\gamma) \in \Pi_1^0(2^\omega)$, and $C = B$. We set $F_1^a(\gamma) := \delta_0$ if $\gamma \in BC_1$, so that condition (2) is fulfilled.

We define $P \subseteq 2^\omega \times \omega$ by

$$(\alpha, k) \in P \Leftrightarrow \alpha \in N(2^\omega, k) \text{ and } \left[\forall j < \text{lh}[\varphi(k)] \left(\left((k)_1 \right)_0 \right)_j < 2 \right].$$

As P is Σ_1^0 , there is $\varepsilon_0 \in \Sigma_1^0 \cap BC_1$ with $P = \rho^{2^\omega \times \omega}(\varepsilon_0)$, by Theorem 3.7. We put $F_2^a(\gamma) := \varepsilon_0$ if $\gamma \in BC_1$, so that $F_2^a(\gamma)$ codes P computing the canonical injection from 2^ω into ω^ω since if $\alpha \in 2^\omega$, then we have $\alpha \in N(\omega^\omega, k) \Leftrightarrow P(\alpha, k)$. So we are done if $\gamma \in BC_1$.

• For the general case, we give the classical scheme of the construction before getting into the effective details, to make things easier to understand. So let $B \in \mathbf{\Pi}_\xi^0$. There is $(B_i)_{i \in \omega} \subseteq \bigcup_{1 \leq \eta < \xi} \mathbf{\Pi}_\eta^0$ such that $B = \bigcap_{i \in \omega} \neg B_i$. Using Lemma 3.10 we will find $(B_{i,j})_{i,j \in \omega} \subseteq \bigcup_{1 \leq \eta < \xi} \mathbf{\Delta}_\eta^0$ with $\neg B_i = \bigcup_{j \in \omega} B_{i,j}$. We will argue by induction on ξ , so that we will get $C_{i,j} \in \mathbf{\Pi}_1^0(\omega^\omega)$, $f_{i,j} : C_{i,j} \rightarrow B_{i,j}$, and $g_{i,j} := f_{i,j}^{-1}$. The objects we are looking for will be the following:

$$C := \left\{ \delta \in \omega^\omega \mid \forall i \in \omega \ [(\delta)_i]^* \in C_{i,(\delta)_i(0)} \text{ and } f_{i,(\delta)_i(0)} \left([(\delta)_i]^* \right) = f_{0,(\delta)_0(0)} \left([(\delta)_0]^* \right) \right\},$$

$f(\delta) := f_{0,(\delta)_0(0)} \left([(\delta)_0]^* \right)$. To define g , we define $h : B \rightarrow \omega^\omega$ by $h(\alpha)(i) := \min \{ j \in \omega \mid \alpha \in B_{i,j} \}$. Note that $h(\alpha)(i)$ is also the unique integer j satisfying $\alpha \in B_{i,j}$. We will have $(g(\alpha))_i := h(\alpha)(i) \frown g_{i,h(\alpha)(i)}(\alpha)$.

• We set

$$(\xi, \gamma) \in Q \Leftrightarrow \xi \geq 2 \text{ and } \gamma \in BC_\xi,$$

$$(\xi, \gamma, \varepsilon) \in Q^+ \Leftrightarrow (\xi, \gamma) \in Q \text{ and } \{\varepsilon\}_2^{\omega^\omega, (\omega^\omega)^3}(\delta) \text{ is defined and in } BC \text{ for each } \delta \in \bigcup_{1 \leq \eta < \xi} BC_\eta,$$

$$(\xi, \gamma, \varepsilon, \alpha) \in Q^{++} \Leftrightarrow (\xi, \gamma, \varepsilon) \in Q^+ \text{ and } \alpha \in B.$$

Assume that $(\xi, \gamma) \in Q$ and γ codes B , so that $\{\gamma^*\}(i)$ is defined for each integer i , and in BC_{η_i} for some $1 \leq \eta_i < \xi$. Using Lemma 3.10, we set $\gamma_{i,j} := \left\{ \left(u_d^{2^\omega} [\{\gamma^*\}(i)] \right)^* \right\}(j)$ for each j . Note that $\gamma_{i,j}$ is recursive in (γ, i, j) , $\gamma_{i,j} \in BC_1$ if $\{\gamma^*\}(i) \in BC_1$, and $\gamma_{i,j} \in \bigcup_{1 \leq \eta < \eta_i} BC_\eta$ if $\{\gamma^*\}(i) \in BC^*$. We also have $B_{i,j} = 2^\omega \setminus \rho^{2^\omega}(\gamma_{i,j})$.

The map F^a will be obtained by Kleene's Recursion Theorem, so that we will have, for some suitable ε^a , $F^a(\gamma) = \varphi^a(\varepsilon^a, \gamma) = \{\varepsilon^a\}^{\omega^\omega, (\omega^\omega)^3}(\gamma)$. In order to describe $\neg C$, we define $R \in \Sigma_1^0[(\omega^\omega)^3]$ as follows:

$$(\varepsilon, \gamma, \delta) \in R \Leftrightarrow \exists i \in \omega \left[\exists j \in \omega \ [(\delta)_i]^* \in N \left(\omega^\omega, [\{\varepsilon\}_0^{\omega^\omega, (\omega^\omega)^3}(\gamma_{i,(\delta)_i(0)})]^*(j) \right) \right] \text{ or}$$

$$\left[\left\{ \{\varepsilon\}_1^{\omega^\omega, (\omega^\omega)^3}(\gamma_{i,(\delta)_i(0)}) \right\}^{\omega^\omega, 2^\omega} \left([(\delta)_i]^* \right) \neq \left\{ \{\varepsilon\}_1^{\omega^\omega, (\omega^\omega)^3}(\gamma_{0,(\delta)_0(0)}) \right\}^{\omega^\omega, 2^\omega} \left([(\delta)_0]^* \right) \right].$$

By 3C.4 and 3C.5 in [Mos80], there is $R^* \in \Delta_1^0[(\omega^\omega)^2 \times \omega^2]$ with

$$(\varepsilon, \gamma, \delta) \in R \Leftrightarrow \exists i \in \omega \left(\delta \in N[\omega^\omega, (i)_0] \text{ and } [\varepsilon, \gamma, (i)_0, (i)_1] \in R^* \right).$$

We define a map $\psi_0 : (\omega^\omega)^2 \rightarrow \omega^\omega$ by

$$\psi_0(\varepsilon, \gamma)(i) := \begin{cases} (i)_0 & \text{if } [\varepsilon, \gamma, (i)_0, (i)_1] \in R^*, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, ψ_0 is recursive and $(\varepsilon, \gamma, \delta) \in R \Leftrightarrow \exists i \in \omega \ \delta \in N[\omega^\omega, \psi_0(\varepsilon, \gamma)(i)]$. We define a recursive map φ_0 by $\varphi_0(\varepsilon, \gamma) := 0 \frown \psi_0(\varepsilon, \gamma)$. Note that $\varphi_0(\varepsilon, \gamma) \in BC_1$ (we will have $F_0^a(\gamma) = \varphi_0(\varepsilon^a, \gamma)$, for ε^a suitable, if $\gamma \in BC^*$).

• We define a partial function $\psi_1 : (\omega^\omega)^3 \rightarrow 2^\omega$ by

$$\psi_1(\varepsilon, \gamma, \delta) := \left\{ \{\varepsilon\}_1^{\omega^\omega, (\omega^\omega)^3}(\gamma_{0,(\delta)_0(0)}) \right\}^{\omega^\omega, 2^\omega} \left([(\delta)_0]^* \right).$$

As ψ_1 is recursive on its domain, there is $\varepsilon_1 \in \Sigma_1^0$ such that $\psi_1(\varepsilon, \gamma, \delta) = \{\varepsilon_1\}^{(\omega^\omega)^3, 2^\omega}(\varepsilon, \gamma, \delta)$ if $\psi_1(\varepsilon, \gamma, \delta)$ is defined. We put $\varphi_1(\varepsilon, \gamma) := S_{\Sigma_1^0}^{(\omega^\omega)^2, \omega^\omega, 2^\omega}(\varepsilon_1, \varepsilon, \gamma)$, so that $\psi_1(\varepsilon, \gamma, \delta)$ is equal to $\{\varphi_1(\varepsilon, \gamma)\}^{\omega^\omega, 2^\omega}(\delta)$ when it is defined. Note that φ_1 is a total recursive map.

• Now we have to describe $\varphi_2^a(\varepsilon, \gamma)$ coding a set computing g . By the proof of 3C.3 in [Mos80] there are recursive maps $g' : \omega \rightarrow \omega$ and $h' : \omega^2 \rightarrow \omega$ such that, for each $(\delta, j, k) \in \omega^\omega \times \omega^2$,

$$\delta(j) = k \Leftrightarrow \exists i \in \omega \ [\delta \in N(\omega^\omega, i) \text{ and } j < g'(i) \text{ and } h'(i, j) = k].$$

We set $k_j := \left(\left((k)_1 \right)_0 \right)_j$. We have, for $\alpha \in B$,

$$\begin{aligned} & g(\alpha) \in N(\omega^\omega, k) \\ & \Leftrightarrow \left((k)_1 \right)_1 \neq 0 \text{ and } \forall j < \mu(k) \ g(\alpha)(j) = k_j \\ & \Leftrightarrow \left((k)_1 \right)_1 \neq 0 \text{ and } \forall j < \mu(k) \ [g(\alpha)]_{(\mathbf{j})_0}[(\mathbf{j})_1] = k_j \\ & \Leftrightarrow \left((k)_1 \right)_1 \neq 0 \text{ and } \forall j < \mu(k) \ \left[\left((\mathbf{j})_1 = 0 \text{ and } h(\alpha)[(\mathbf{j})_0] = k_j \right) \text{ or} \right. \\ & \quad \left. \left((\mathbf{j})_1 > 0 \text{ and } g_{(\mathbf{j})_0, h(\alpha)[(\mathbf{j})_0]}(\alpha)[(\mathbf{j})_1 - 1] = k_j \right) \right] \\ & \Leftrightarrow \left((k)_1 \right)_1 \neq 0 \text{ and } \forall j < \mu(k) \ \left[\left((\mathbf{j})_1 = 0 \text{ and } \alpha \in B_{(\mathbf{j})_0, k_j} \right) \text{ or} \right. \\ & \quad \left. \left((\mathbf{j})_1 > 0 \text{ and } \exists i \in \omega \ \left[g_{(\mathbf{j})_0, h(\alpha)[(\mathbf{j})_0]}(\alpha) \in N(\omega^\omega, i) \text{ and } (\mathbf{j})_1 \leq g'(i) \text{ and } h'[i, (\mathbf{j})_1 - 1] = k_j \right] \right) \right]. \end{aligned}$$

But $\alpha \in B_{(\mathbf{j})_0, k_j} \Leftrightarrow \alpha \notin \rho^{2^\omega}(\gamma_{(\mathbf{j})_0, k_j}) \Leftrightarrow \exists l \in \omega [k_j = l \text{ and } \alpha \notin \rho^{2^\omega}(\gamma_{(\mathbf{j})_0, l})]$. There is $\delta_1 \in \Sigma_1^0$ such that $\{\delta_1\}^{2^\omega \times \omega, 2^\omega}(\alpha, k) = \alpha$ if $(\alpha, k) \in 2^\omega \times \omega$.

If $\{\gamma^*\}[(\mathbf{j})_0](0) = 0$, then by Lemmas 3.10 and 3.3 we get

$$\begin{aligned} \alpha \notin \rho^{2^\omega}(\gamma_{(\mathbf{j})_0, l}) &\Leftrightarrow \alpha \in \rho^{2^\omega} \left[u_c^{2^\omega} \left(\{\gamma^*\}[(\mathbf{j})_0], l \right) \right] \\ &\Leftrightarrow (\alpha, k) \in \rho^{2^\omega \times \omega} \left(u_r^{2^\omega \times \omega, 2^\omega} \left[u_c^{2^\omega} \left(\{\gamma^*\}[(\mathbf{j})_0], l \right), \delta_1 \right] \right). \end{aligned}$$

If $\{\gamma^*\}[(\mathbf{j})_0](0) \neq 0$, then by Lemmas 3.3 and 3.1 we get

$$\begin{aligned} \alpha \notin \rho^{2^\omega}(\gamma_{(\mathbf{j})_0, l}) &\Leftrightarrow (\alpha, k) \notin \rho^{2^\omega \times \omega} [u_r^{2^\omega \times \omega, 2^\omega}(\gamma_{(\mathbf{j})_0, l}, \delta_1)] \\ &\Leftrightarrow (\alpha, k) \in \rho^{2^\omega \times \omega} \left(u_{-1} [u_r^{2^\omega \times \omega, 2^\omega}(\gamma_{(\mathbf{j})_0, l}, \delta_1)] \right). \end{aligned}$$

This shows the existence of a partial function $\tilde{g}_0 : \omega^\omega \times \omega^2 \rightarrow \omega^\omega$, recursive on its domain, such that $\tilde{g}_0(\gamma, j, l)$ is defined if $(\xi, \gamma) \in Q$. In this case, $\tilde{g}_0(\gamma, j, l) \in BC_1$ if $\{\gamma^*\}[(\mathbf{j})_0] \in BC_1$, $\tilde{g}_0(\gamma, j, l) \in BC_{\eta_{(\mathbf{j})_0}}$ if $\{\gamma^*\}[(\mathbf{j})_0] \in BC^*$, and

$$\alpha \in B_{(\mathbf{j})_0, l} \Leftrightarrow \alpha \notin \rho^{2^\omega}(\gamma_{(\mathbf{j})_0, l}) \Leftrightarrow (\alpha, k) \in \rho^{2^\omega \times \omega} [\tilde{g}_0(\gamma, j, l)].$$

Choose $\gamma_1 \in \Sigma_1^0 \cap BC_1$ such that $(\alpha, k, j, l) \in \rho^{2^\omega \times \omega^3}(\gamma_1) \Leftrightarrow k_j = l$. Using Lemma 3.2 we see that $u_s^{\omega^2}(\gamma_1, j, l) \in BC_1$ and $(\alpha, k) \in \rho^{2^\omega \times \omega} [u_s^{\omega^2}(\gamma_1, j, l)] \Leftrightarrow k_j = l$, for each $(\alpha, k, j, l) \in 2^\omega \times \omega^3$. Using Lemmas 3.5 and 3.6.(b), we get the existence of a partial function $g_0 : \omega^\omega \times \omega \rightarrow \omega^\omega$, recursive on its domain, such that $g_0(\gamma, j)$ is defined if $(\xi, \gamma) \in Q$. In this case, $g_0(\gamma, j) \in BC_1$ if $\{\gamma^*\}[(\mathbf{j})_0] \in BC_1$, and $g_0(\gamma, j) \in BC_{\eta_{(\mathbf{j})_0}}$ if $\{\gamma^*\}[(\mathbf{j})_0] \in BC^*$. If moreover $\alpha \in B$, then $\alpha \in B_{(\mathbf{j})_0, k_j} \Leftrightarrow (\alpha, k) \in \rho^{2^\omega \times \omega} [g_0(\gamma, j)]$.

- We will have

$$\begin{aligned} g_{(\mathbf{j})_0, h(\alpha)[(\mathbf{j})_0]}(\alpha) \in N(\omega^\omega, i) &\Leftrightarrow (\alpha, i) \in \rho^{2^\omega \times \omega} [\{\varepsilon\}_2^{\omega^\omega, (\omega^\omega)^3}(\gamma_{(\mathbf{j})_0, h(\alpha)[(\mathbf{j})_0]})] \\ &\Leftrightarrow \exists l \in \omega \left[(\alpha, i) \in \rho^{2^\omega \times \omega} [\{\varepsilon\}_2^{\omega^\omega, (\omega^\omega)^3}(\gamma_{(\mathbf{j})_0, l})] \text{ and } h(\alpha)[(\mathbf{j})_0] = l \right] \\ &\Leftrightarrow \exists l \in \omega \left[(\alpha, i) \in \rho^{2^\omega \times \omega} [\{\varepsilon\}_2^{\omega^\omega, (\omega^\omega)^3}(\gamma_{(\mathbf{j})_0, l})] \text{ and } \alpha \in B_{(\mathbf{j})_0, l} \right] \end{aligned}$$

if $(\xi, \gamma, \varepsilon, \alpha) \in Q^{++}$. If we apply Lemmas 3.5 and 3.6.(a), then we see the existence of a partial function $g_1^0 : (\omega^\omega)^2 \times \omega^2 \rightarrow \omega^\omega$, recursive on its domain, such that $g_1^0(\varepsilon, \gamma, j, l)$ is defined and in BC if $(\xi, \gamma, \varepsilon) \in Q^+$, in which case $(\alpha, k) \in \rho^{2^\omega \times \omega} [g_1^0(\varepsilon, \gamma, j, l)]$ is equivalent to

$$\exists i \in \omega \left[(\alpha, i) \in \rho^{2^\omega \times \omega} [\{\varepsilon\}_2^{\omega^\omega, (\omega^\omega)^3}(\gamma_{(\mathbf{j})_0, l})] \text{ and } (\mathbf{j})_1 \leq g'(i) \text{ and } h'[i, (\mathbf{j})_1 - 1] = k_j \right].$$

If $(\xi, \gamma, \varepsilon, \alpha) \in Q^{++}$, then

$$\begin{aligned} \exists i \in \omega \left[g_{(\mathbf{j})_0, h(\alpha)[(\mathbf{j})_0]}(\alpha) \in N(\omega^\omega, i) \text{ and } (\mathbf{j})_1 \leq g'(i) \text{ and } h'[i, (\mathbf{j})_1 - 1] = k_j \right] \\ \Leftrightarrow \exists l \in \omega \left[(\alpha, k) \in \rho^{2^\omega \times \omega} [g_1^0(\varepsilon, \gamma, j, l)] \text{ and } \alpha \in B_{(\mathbf{j})_0, l} \right]. \end{aligned}$$

But $g_1^0(\varepsilon, \gamma, j, l)$ could be in BC_1 for some l 's, and in BC^* for some others. This may happen if $\xi \geq 3$. This is a problem since we want to apply Lemma 3.6.(b). We will solve this problem with Lemma 3.4. We define a partial function $g_1^1 : (\omega^\omega)^2 \times \omega^2 \rightarrow \omega^\omega$ by $g_1^1(\varepsilon, \gamma, j, l) := u_*^{2^\omega \times \omega}[g_1^0(\varepsilon, \gamma, j, l)]$. As $\rho^{2^\omega \times \omega}[g_1^1(\varepsilon, \gamma, j, l)] = \rho^{2^\omega \times \omega}[g_1^0(\varepsilon, \gamma, j, l)]$ if $(\xi, \gamma, \varepsilon) \in Q^+$, it satisfies the previous properties of g_1^0 .

- Lemmas 3.5 and 3.6.(b) imply the existence of a partial function $g_2^1 : (\omega^\omega)^2 \times \omega \rightarrow \omega^\omega$, recursive on its domain, such that $g_2^1(\varepsilon, \gamma, j)$ is defined and in BC if $(\xi, \gamma, \varepsilon) \in Q^+$. If moreover $(\xi, \gamma, \varepsilon, \alpha) \in Q^{++}$, then $(\alpha, k) \in \rho^{2^\omega \times \omega}[g_2^1(\varepsilon, \gamma, j)]$ is equivalent to

$$\exists i \in \omega \left[g_{(\mathbf{j})_0, h(\alpha)[(\mathbf{j})_0]}(\alpha) \in N(\omega^\omega, i) \text{ and } (\mathbf{j})_1 \leq g'(i) \text{ and } h'[i, (\mathbf{j})_1 - 1] = k_j \right].$$

We also define a partial function $g_2^0 : (\omega^\omega)^2 \times \omega \rightarrow \omega^\omega$. It is defined relatively to g_1^0 exactly like g_2^1 was defined relatively to g_1^1 . It will satisfy the previous properties of g_2^1 if $\xi = 2$, and we will have, for $(\xi, \gamma, \varepsilon, \alpha) \in Q^{++}$ and $a \in 2$,

$$g(\alpha) \in N(\omega^\omega, k) \Leftrightarrow \left((k)_1 \right)_1 \neq 0 \text{ and } \forall j < \mu(k) \left[\left((\mathbf{j})_1 = 0 \text{ and } (\alpha, k) \in \rho^{2^\omega \times \omega}[g_0(\gamma, j)] \right) \text{ or } \left((\mathbf{j})_1 > 0 \text{ and } (\alpha, k) \in \rho^{2^\omega \times \omega}[g_2^a(\varepsilon, \gamma, j)] \right) \right].$$

- We define a partial function $g_3^1 : (\omega^\omega)^2 \times \omega \rightarrow \omega^\omega$ by

$$g_3^1(\varepsilon, \gamma, j) := \begin{cases} u_*^{2^\omega \times \omega}[g_0(\gamma, j)] & \text{if } (\mathbf{j})_1 = 0, \\ g_2^1(\varepsilon, \gamma, j) & \text{if } (\mathbf{j})_1 > 0. \end{cases}$$

Note that g_3^1 is recursive on its domain, and $g_3^1(\varepsilon, \gamma, j)$ is defined and in BC if $(\xi, \gamma, \varepsilon) \in Q^+$. We also define a partial function $g_3^0 : (\omega^\omega)^2 \times \omega \rightarrow \omega^\omega$. It is defined relatively to g_2^0 , like g_3^1 was defined relatively to g_2^1 , except that $g_3^0(\varepsilon, \gamma, j) := g_0(\gamma, j)$ if $(\mathbf{j})_1 = 0$. The function g_3^0 will satisfy the previous properties of g_3^1 if $\xi = 2$.

- By Lemma 3.5, we get the existence of a partial function $g_4^1 : (\omega^\omega)^2 \times \omega \rightarrow \omega^\omega$, recursive on its domain, such that $g_4^1(\varepsilon, \gamma, m)$ is defined and in BC if $(\xi, \gamma, \varepsilon) \in Q^+$ and, if moreover $(\xi, \gamma, \varepsilon, \alpha) \in Q^{++}$, then

$$(\alpha, k) \in \rho^{2^\omega \times \omega}[g_4^1(\varepsilon, \gamma, m)] \Leftrightarrow \left((k)_1 \right)_1 \neq 0 \text{ and } \mu(k) = m \text{ and } \forall j < m \ (\alpha, k) \in \rho^{2^\omega \times \omega}[g_3^1(\varepsilon, \gamma, j)].$$

Thus $(\xi, \gamma, \varepsilon, \alpha) \in Q^{++}$ will imply that

$$\begin{aligned} g(\alpha) \in N(\omega^\omega, k) &\Leftrightarrow \left((k)_1 \right)_1 \neq 0 \text{ and } \forall j < \mu(k) \ (\alpha, k) \in \rho^{2^\omega \times \omega}[g_3^1(\varepsilon, \gamma, j)] \\ &\Leftrightarrow \exists m \in \omega \left[\left((k)_1 \right)_1 \neq 0 \text{ and } \mu(k) = m \text{ and } \forall j < m \ (\alpha, k) \in \rho^{2^\omega \times \omega}[g_3^1(\varepsilon, \gamma, j)] \right] \\ &\Leftrightarrow \exists m \in \omega \ (\alpha, k) \in \rho^{2^\omega \times \omega}[g_4^1(\varepsilon, \gamma, m)]. \end{aligned}$$

We also define a partial function $g_4^0 : (\omega^\omega)^2 \times \omega \rightarrow \omega^\omega$. It is defined relatively to g_3^0 exactly like g_4^1 was defined relatively to g_3^1 . It will satisfy the previous properties of g_4^1 if $\xi = 2$.

- By Lemma 3.6.(b), we get the existence of a partial function $\varphi_2^1: (\omega^\omega)^2 \rightarrow \omega^\omega$, recursive on its domain, such that $\varphi_2^1(\varepsilon, \gamma)$ is defined and in BC if $(\xi, \gamma, \varepsilon) \in Q^+$, and

$$g(\alpha) \in N(\omega^\omega, k) \Leftrightarrow (\alpha, k) \in \rho^{2^\omega \times \omega}[\varphi_2^1(\varepsilon, \gamma)]$$

if $(\xi, \gamma, \varepsilon, \alpha) \in Q^{++}$. We also define a partial function $\varphi_2^0: (\omega^\omega)^2 \rightarrow \omega^\omega$. It is defined relatively to g_4^0 exactly like φ_2^1 was defined relatively to g_4^1 . It will satisfy the previous properties of φ_2^1 if $\xi = 2$.

• Now we can define a partial function $\varphi^a: (\omega^\omega)^2 \rightarrow (\omega^\omega)^3$ by

$$\varphi^a(\varepsilon, \gamma) := \begin{cases} [F_0^a(\gamma), F_1^a(\gamma), F_2^a(\gamma)] & \text{if } \gamma(0) = 0, \\ [\varphi_0(\varepsilon, \gamma), \varphi_1(\varepsilon, \gamma), \varphi_2^a(\varepsilon, \gamma)] & \text{if } \gamma(0) \neq 0. \end{cases}$$

As φ^a is recursive on its domain, by Kleene's Recursion Theorem there is $\varepsilon^a \in \Sigma_1^0$ such that

$$\{\varepsilon^a\}^{\omega^\omega, (\omega^\omega)^3}(\gamma) = \varphi^a(\varepsilon^a, \gamma)$$

if $\varphi^a(\varepsilon^a, \gamma)$ is defined. We define a partial function $F^a: \omega^\omega \rightarrow (\omega^\omega)^3$ by $F^a(\gamma) := \{\varepsilon^a\}^{\omega^\omega, (\omega^\omega)^3}(\gamma)$, so that F^a is recursive on its domain. We already checked that $F^a(\gamma)$ is suitable if $\gamma \in BC_1$.

So assume that $2 \leq \xi < \omega_1$, and $\gamma \in BC_\xi$ codes $B := \neg \rho^{2^\omega}(\gamma) \in \Pi_\xi^0$. We will prove that $F^a(\gamma)$ is defined and fullfills the required properties by induction on ξ .

Note that $\{\varepsilon^a\}_2^{\omega^\omega, (\omega^\omega)^3}(\delta)$ is defined and in BC for each $\delta \in \bigcup_{1 \leq \eta < \xi} BC_\eta$, by induction assumption. This implies that $(\xi, \gamma, \varepsilon^a) \in Q^+$, $\varphi_2^a(\varepsilon^a, \gamma)$ and $F^a(\gamma)$ are defined, and

$$F^a(\gamma) = \{\varepsilon^a\}^{\omega^\omega, (\omega^\omega)^3}(\gamma) = \varphi^a(\varepsilon^a, \gamma) = [\varphi_0(\varepsilon^a, \gamma), \varphi_1(\varepsilon^a, \gamma), \varphi_2^a(\varepsilon^a, \gamma)].$$

(1) Note that $F_0^a(\gamma) \in BC_1$ since $F_0^a(\gamma) = \varphi_0(\varepsilon^a, \gamma) = 0 \smallfrown \psi_0(\varepsilon^a, \gamma)$. Moreover, with the previous notation, we get $\delta \notin \rho^{\omega^\omega}[F_0^a(\gamma)] \Leftrightarrow (\varepsilon^a, \gamma, \delta) \notin R \Leftrightarrow \delta \in C$, by induction assumption.

(2) We have $F_1^a(\gamma) = \varphi_1(\varepsilon^a, \gamma)$, so that, by induction assumption, and for each $\delta \in C$,

$$\begin{aligned} \{F_1^a(\gamma)\}^{\omega^\omega, 2^\omega}(\delta) &= \psi_1(\varepsilon^a, \gamma, \delta) \\ &= \left\{ \{\varepsilon^a\}_1^{\omega^\omega, (\omega^\omega)^3}(\gamma_{0,(\delta)_0(0)}) \right\}^{\omega^\omega, 2^\omega} \left([(\delta)_0]^* \right) \\ &= \left\{ F_1^a(\gamma_{0,(\delta)_0(0)}) \right\}^{\omega^\omega, 2^\omega} \left([(\delta)_0]^* \right) \\ &= f_{0,(\delta)_0(0)} \left([(\delta)_0]^* \right) = f(\delta). \end{aligned}$$

Clearly, f is continuous. If $\delta \in C$ and $i \in \omega$, then $f(\delta) = f_{i,(\delta)_i(0)} \left([(\delta)_i]^* \right) \in B_{i,(\delta)_i(0)}$, thus $f(\delta) \in B$. Let $\delta, \delta' \in C$ such that $\alpha := f(\delta) = f(\delta')$. Then $\alpha = f_{i,(\delta)_i(0)} \left([(\delta)_i]^* \right) \in B_{i,(\delta)_i(0)}$, so that α is in $B_{i,(\delta)_i(0)}$ and $B_{i,(\delta')_i(0)}$. This shows that $(\delta)_i(0) = (\delta')_i(0)$. Thus $[(\delta)_i]^* = [(\delta')_i]^*$ since $f_{i,(\delta)_i(0)}$ is one-to-one, $(\delta)_i = (\delta')_i$, and $\delta = \delta'$. This shows that f is one-to-one. If $\alpha \in B$ and $i \in \omega$, then there is a unique integer j_i with $\alpha \in B_{i,j_i}$. There is $\delta^i \in C_{i,j_i}$ with $\alpha = f_{i,j_i}(\delta^i)$. Put $(\delta)_i := j_i \smallfrown \delta^i$. Then $\delta \in C$ and $\alpha = f(\delta)$. This shows that f is onto.

(3) We have $F_2^a(\gamma) = \varphi_2^a(\varepsilon^a, \gamma)$.

- If $\xi = 2$, then $\eta_i = 1$ for each i , $\gamma_{i,j} \in BC_1$ for each (i, j) . Thus

◦ $g_1^0(\varepsilon^a, \gamma, j, l) \in BC_1$, by induction assumption, since $\{\varepsilon^a\}_2^{\omega, (\omega^\omega)^3}(\gamma_{(j)0}, l) \in BC_1$. This implies that $g_1^1(\varepsilon^1, \gamma, j, l) \in BC_2$.

◦ $g_2^1(\varepsilon^1, \gamma, j) \in BC_2$ and $g_2^0(\varepsilon^a, \gamma, j) \in BC_1$.

◦ $g_3^1(\varepsilon^1, \gamma, j) \in BC_2$ and $g_3^0(\varepsilon^a, \gamma, j) \in BC_1$.

◦ $g_4^1(\varepsilon^1, \gamma, m) \in BC_2$ and $g_4^0(\varepsilon^a, \gamma, m) \in BC_1$.

◦ $\varphi_2^1(\varepsilon^1, \gamma) \in BC_2 = BC_{\xi^-}$ and $\varphi_2^0(\varepsilon^a, \gamma) \in BC_1$.

- If $\xi \geq 3$, then

◦ $g_1^1(\varepsilon^1, \gamma, j, l) \in BC_{\max(2, \eta_{(j)0})}$, by induction assumption. Indeed, if $\gamma_{(j)0}, l \in BC_1$, then $g_1^0(\varepsilon^1, \gamma, j, l)$ is in BC_1 and $g_1^1(\varepsilon^1, \gamma, j, l) \in BC_2$. If $\gamma_{(j)0}, l \in BC^*$, then $\gamma_{(j)0}, l \in BC_{\eta_{(j)0}}$, and $g_1^0(\varepsilon^1, \gamma, j, l), g_1^1(\varepsilon^1, \gamma, j, l)$ too.

◦ $g_2^1(\varepsilon^1, \gamma, j) \in BC_{\max(2, \eta_{(j)0})}$.

◦ $g_3^1(\varepsilon^1, \gamma, j) \in BC_{\max(2, \eta_{(j)0})}$.

◦ $g_4^1(\varepsilon^1, \gamma, m) \in BC_{\max_{j < m}(2, \eta_{(j)0})} \subseteq BC_{\xi-1}$.

◦ $\varphi_2^1(\varepsilon^1, \gamma) \in BC_{\xi-1} = BC_{\xi^-}$.

Thus $F_2^0(\gamma) = \varphi_2^0(\varepsilon^0, \gamma) \in BC_1$ if $\gamma \in BC_2$, and $F_2^1(\gamma) = \varphi_2^1(\varepsilon^1, \gamma) \in BC_{\xi^-}$. And $\rho^{2^\omega \times \omega}[F_2^a(\gamma)]$ computes g on B . If $\alpha \in B$ and $i \in \omega$, then $\left[\left(g(\alpha) \right)_i \right]^* = g_{i, h(\alpha)(i)}(\alpha) \in C_{i, h(\alpha)(i)}$ since $\alpha \in B_{i, h(\alpha)(i)}$. Thus

$$f_{i, h(\alpha)(i)}[g_{i, h(\alpha)(i)}(\alpha)] = \alpha$$

and $g(\alpha) \in C$ since $\left(g(\alpha) \right)_i(0) = h(\alpha)(i)$. Moreover,

$$f[g(\alpha)] = f_{0, (g(\alpha))_0(0)} \left(\left[\left(g(\alpha) \right)_0 \right]^* \right) = \alpha.$$

If $\delta \in C$ and $i \in \omega$, then $\left(g[f(\delta)] \right)_i = h[f(\delta)](i) \frown g_{i, h[f(\delta)](i)}[f(\delta)] = (\delta)_i(0) \frown [(\delta)_i]^* = (\delta)_i$. Therefore $g[f(\delta)] = \delta$. This shows that g coincides with f^{-1} . \square

Proof of Theorem 1.5. Let $\gamma \in \Sigma_1^0 \cap BC_{\xi+1}$ with $B = \neg \rho^{2^\omega}(\gamma)$. By Theorem 4.1,

• If $\xi = 1$, then $F^0(\gamma) \in \Sigma_1^0$, $C \in \Pi_1^0$, f is a partial recursive function on C , and g is a partial Σ_ξ^0 -recursive function on B .

• If $\xi \geq 2$, then $F^1(\gamma) \in \Sigma_1^0$ and the same conclusion holds. \square

We also have a Δ_1^1 version of Theorem 1.5:

Theorem 4.2 *Let $\xi \geq 1$ be a countable ordinal, and $B \in \Pi_{\xi+1}^0(2^\omega) \cap \Delta_1^1$. Then there is $C \in \Pi_1^0 \cap \Delta_1^1(\omega^\omega)$, a Δ_1^1 -recursive function $f : \omega^\omega \rightarrow 2^\omega$, and a Δ_1^1 -recursive function $g : 2^\omega \rightarrow \omega^\omega$, such that $f|_C$ defines a continuous bijection from C onto B , $g|_B$ is Σ_ξ^0 -measurable, and $g|_B$ coincides with $(f|_C)^{-1}$.*

Proof. We set $\Sigma_\xi^0(\Delta_1^1)(X) = \{\rho^X(\gamma) \mid \gamma \in \Delta_1^1 \cap BC_\xi\}$ if $1 \leq \xi < \omega_1$. In [Lou80], it is essentially proved that $\Sigma_\xi^0(\Delta_1^1) = \Sigma_\xi^0 \cap \Delta_1^1$. Actually, A. Louveau does not use the coding for Borel sets that we use here, but he proves this specific result, with this coding, in [Lou??]. So let $\gamma \in \Delta_1^1 \cap BC_{\xi+1}$ with $B = \neg\rho^{2^\omega}(\gamma)$. By Theorem 4.1,

- If $\xi = 1$, then $F^0(\gamma) \in \Delta_1^1$, $C \in \Pi_1^0 \cap \Delta_1^1$, f and g are partial Δ_1^1 functions on Δ_1^1 sets, and can be extended to total Δ_1^1 maps.
- If $\xi \geq 2$, then $F^1(\gamma) \in \Delta_1^1$ and the same conclusion holds. □

5 Proof of Theorem 1.3.

The proof of Theorem 1.3.(2) is essentially identical to that of Theorem 1.3.(1), so it is enough to prove Theorem 1.3.(1) to get Theorem 1.3. In the sequel we will assume that $\xi < \omega_1^{CK}$, except where indicated. Let us indicate the specifications of the proof of Theorem 1.2 that we need. Theorem 3.14 gives $B \in \Gamma(2^\omega) \setminus \check{\Gamma}$. As $B \in \Pi_{\xi+1}^0$, Theorem 1.5 gives C , f and g . Here again, the dictionary A will be made of two pieces: we will have $A = \mu \cup \pi$ if $\xi \geq 3$.

Notation. Recall that $Q := \{(s, t) \in 2^{<\omega} \times 2^{<\omega} \mid |s| = |t|\}$. We will sometimes view Q as $\tilde{Q} \in \Delta_1^0(\omega)$:

$$\tilde{Q} := \left\{ m \in \omega \mid \text{Seq}(m) \text{ and } \forall i < \text{lh}(m) \left[\text{Seq}[(m)_i] \text{ and } \text{lh}[(m)_i] = 2 \text{ and } \forall j \in 2 \left((m)_i \right)_j < 2 \right] \right\}.$$

Implicitly, we have used the bijection $I : Q \rightarrow \tilde{Q}$ defined by

$$I(s, t) := \left\langle \langle s(0), t(0) \rangle, \dots, \langle s(|s|-1), t(|s|-1) \rangle \right\rangle.$$

Note that the map $\chi : \omega \rightarrow \omega$ defined by $\chi(r) := I(q_r)$ is a recursive injection with range \tilde{Q} . We define a recursive map $M : \omega \rightarrow \omega$ by $M(j) := M_j := \Sigma_{i < j} 4^{i+1}$.

Lemma 5.1 *The sets μ^0 , μ^1 and μ can be coded by recursive subsets of ω .*

Proof. We define a recursive map $\text{Exp} : \omega^2 \rightarrow \omega$ coding the finite sequence k^j :

$$\text{Exp}(k, j) := c \iff \text{Seq}(c) \text{ and } \text{lh}(c) = j \text{ and } \forall i < j (c)_i = k.$$

Using Exp , it is easy to build a recursive map $f : \omega^5 \rightarrow \omega$ such that $f(N, l, m, P, R)$ codes the sequence $2^N \hat{\cap} [\hat{\cap}_{i \leq l+1} m_i 2^{P_i} 3 2^{R_i}]$. Then we just have to use bounded quantifiers. □

Now we show that μ^∞ is “simple”.

Lemma 5.2 *The set μ^∞ is $\Pi_2^0(4^\omega)$.*

Proof. We have

$$\gamma \in \mu^\infty \Leftrightarrow \exists i \in 2 \ \forall j \in \omega \ \exists k \in \omega \ \exists t \in (\mu^i)^{<\omega} \ |t| \geq j \ \text{and} \ \gamma \upharpoonright k = \bigwedge_{l < |t|} t(l).$$

This shows that $\mu^\infty \in \Pi_2^0(4^\omega)$, by Lemma 5.1, since t can be coded by an integer, and the restriction and concatenation maps are recursive. \square

Notation. We define a partial function $c: 2^\omega \times \omega \rightarrow Q$ on $B \times \omega$ by $c(\alpha, l) := [g(\alpha), \alpha] \upharpoonright l$.

Lemma 5.3 *The set $E := \{(N, \alpha) \in \omega \times 2^\omega \mid \alpha \in E_N\}$ is in Γ .*

Proof. The map $h: \omega \times 2^\omega \rightarrow 2^\omega$ defined by $h(N, \alpha) := q_N^1 \alpha$ is clearly recursive. From this we deduce that E is in Γ , using Lemmas 3.3 and 3.5. \square

Notation. Now we code the maps $\varphi_{N,j}$. We set $\text{Dom} := \{(N, j, \gamma) \in \omega^2 \times 4^\omega \mid N \leq M_j \ \text{and} \ \gamma \in K_{N,j}\}$. We define a partial function $\tilde{\varphi}: \omega^2 \times 4^\omega \rightarrow 2^\omega$ as follows: $\tilde{\varphi}(N, j, \gamma)$ is defined if $(N, j, \gamma) \in \text{Dom}$, and its coordinates are the coordinates of γ in 2, in the same order as in γ (we forget the **2**'s and the **3**'s).

In the next lemma we consider the set expressing the fact that “ π^∞ will look like B on $K_{N,j}$ ”.

Lemma 5.4 *The set $\mathcal{F} := \{(N, j, \gamma) \in \text{Dom} \mid \tilde{\varphi}(N, j, \gamma) \in E_N\}$ is in Γ if $\xi \geq 2$.*

Proof. We define a map $\psi: 2^\omega \times \omega^2 \rightarrow 4^\omega$ by

$$\psi(\alpha, N, j) := \begin{cases} \mathbf{2}^N \alpha(0) \frown [\bigwedge_{k \in \omega} \mathbf{2}^{M(j+k+1)} \mathbf{3} \mathbf{2}^{M(j+k+1)} \alpha(k+1)] & \text{if } N \leq M(j), \\ 0^\infty & \text{if } N > M(j). \end{cases}$$

It is easy to see that ψ is recursive. If $N \leq M_j$ and $\gamma \in 4^\omega$, then $\gamma \in K_{N,j}$ is equivalent to

$$\forall i \in \omega \ [\psi(0^\infty, N, j)(i) = 0 \ \text{and} \ \gamma(i) \in 2] \ \text{or} \ [\psi(0^\infty, N, j)(i) \neq 0 \ \text{and} \ \gamma(i) = \psi(0^\infty, N, j)(i)].$$

This shows that $\text{Dom} \in \Pi_1^0$. Then $\tilde{\varphi}$ is clearly recursive on Dom . This shows that \mathcal{F} is in Γ if $\xi \geq 2$, by Lemmas 3.3, 3.5 and 5.3. \square

Now we describe the elements of $A^\infty \setminus \mu^\infty$.

Notation. Recall that $P_{t,S,j} := \{ \gamma \in 4^\omega \mid t \mathbf{2}^S \prec \gamma \ \text{and} \ \gamma - t \mathbf{2}^S \in K_{0,j} \}$. Note that the relation defined by “ $\gamma \in P_{t,S,j}$ ” is Π_1^0 in γ, t, S, j . Let (t, S, j) be suitable and $N \leq \min(M_j, S)$ ($N = S$ if $t = \emptyset$). Note that $(N, j, \gamma - t \mathbf{2}^{S-N}) \in \mathcal{F}$ means that $\gamma - t \mathbf{2}^{S-N} \in \pi^\infty \cap K_{N,j}$. This implies that

$$A_{t,S,j,N} = \{ \gamma \in P_{t,S,j} \mid (N, j, \gamma - t \mathbf{2}^{S-N}) \in \mathcal{F} \}.$$

Lemma 5.5 *The set of $(\gamma, t, S, j, N) \in 4^\omega \times (\{\emptyset\} \cup \mu) \times \omega^3$ such that (t, S, j) is suitable, $N \leq \min[M(j), S]$, $N = S$ if $t = \emptyset$ and $\gamma \in A_{t,S,j,N}$ can be coded by a set in $\Gamma(4^\omega \times \omega^4)$ if $\xi \geq 2$.*

Proof. Apply Lemmas 3.3, 3.5 and 5.4. \square

Let us specify a few facts about the definition of π .

Notation. As C is Π_1^0 and f is recursive on C , the graph $\text{Gr}(f)$ of f is a Π_1^0 subset of $\omega^\omega \times 2^\omega$. As the identity from 2^ω , viewed as a subset of ω^ω , into 2^ω is a partial recursive function on 2^ω (see the proof of Theorem 4.1), we can also say that $\text{Gr}(f)$ is a Π_1^0 subset of $\omega^\omega \times \omega^\omega$, by Lemma 3.3. By 4A.1 in [Mos80], there is $R \in \Delta_1^0(\omega^2)$ such that $\alpha = f(\beta) \Leftrightarrow \forall k \in \omega \ (\overline{\beta[k], \overline{\alpha[k]}}) \in R$ (recall that \bar{t} is defined at the beginning of section 3).

- We set $Q_f := \{(t, s) \in Q \mid (\bar{t}, \bar{s}) \in R \text{ and } t \neq \emptyset \text{ and } t(|t| - 1) = 1\}$. Note that Q_f can easily be coded by a recursive subset of ω .

- The definition of π is the same as the one in section 2. Here again, π can easily be coded by a recursive subset of ω .

Proof of Theorem 1.3.(1). We refer to the proof of Theorem 1.2. We put $A := \mu \cup \pi$, so that A can be coded by a Δ_1^0 subset of ω . We will prove that $A^\infty \in \Gamma \setminus \check{\Gamma}$.

- Here again we have $\varphi_{N,j}[\pi^\infty \cap K_{N,j}] = E_N$ if $N \leq M_j$. If $\gamma \in \pi^\infty \cap K_{N,j}$, then the only thing to notice is that $[\overline{\beta[k], \overline{(q_N^1 \alpha)[k]}}] \in R$ for each $k \in \omega$.

- We also have

$$A^\infty = \mu^\infty \cup \bigcup_{(t,S,j) \text{ suitable}} \bigcup_{\substack{N \leq \min(M_j, S) \\ N = S \text{ if } t = \emptyset}} A_{t,S,j,N}.$$

As Γ is uniformly closed under finite unions, the set of $(\gamma, t, S, j) \in 4^\omega \times (\{\emptyset\} \cup \mu) \times \omega^2$ such that (t, S, j) is suitable and $\gamma \in A_{t,S,j}$ can be coded by a set in $\Gamma(4^\omega \times \omega^3)$ if $\xi \geq 2$, by Lemma 5.5.

- By Lemmas 3.5, 3.6 and 5.2, we get $A^\infty \in \Gamma(4^\omega)$ if $\xi \geq 3$ and $\Gamma = \Sigma_\xi^0$.

- If $\xi \geq 3$ and $\Gamma = \Pi_\xi^0$, then we can write

$$A^\infty = \mu^\infty \setminus \left(\bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j} \right) \cup \bigcup_{(t,S,j) \text{ suitable}} A_{t,S,j} \cap P_{t,S,j}.$$

Thus

$$\neg A^\infty = \neg \left[\mu^\infty \cup \left(\bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j} \right) \right] \cup \bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j} \setminus A_{t,S,j}.$$

Here $\neg \left[\mu^\infty \cup \left(\bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j} \right) \right] \in \Delta_3^0(4^\omega) \subseteq \check{\Gamma}(4^\omega)$. By Lemma 3.6, $\bigcup_{(t,S,j) \text{ suitable}} P_{t,S,j} \setminus A_{t,S,j}$ is in $\check{\Gamma}(4^\omega)$, and by Lemma 3.5 $\neg A^\infty$ is in $\check{\Gamma}(4^\omega)$. Thus $A^\infty \in \Gamma(4^\omega)$.

- If $1 \leq \xi \leq 2$, then we argue as in the proof of Theorem 1.2. □

6 On the complexity of some sets of dictionaries.

The proof of Theorem 1.3 has the following consequence on the complexity of the sets Σ_ξ and Π_ξ defined in the introduction. Recall that if $1 \leq \xi < \omega_1$, then

$$\Sigma_\xi := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Sigma_\xi^0\} \quad \text{and} \quad \Pi_\xi := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Pi_\xi^0\}.$$

Notation. We set

$$\Sigma'_\xi := \{\gamma \in BC \mid \rho^{2^\omega}(\gamma) \in \Sigma_\xi^0\} \quad \text{and} \quad \Pi'_\xi := \{\gamma \in BC \mid \rho^{2^\omega}(\gamma) \in \Pi_\xi^0\}.$$

Corollary 6.1 *Let $3 \leq \xi < \omega_1$. Then there is $\varphi : \omega^\omega \rightarrow 2^{2^{<\omega}}$ continuous with $\Sigma'_\xi = BC \cap \varphi^{-1}(\Sigma_\xi)$ and $\Pi'_\xi = BC \cap \varphi^{-1}(\Pi_\xi)$.*

So Σ_ξ (resp., Π_ξ) is more complicated than the set of Borel codes for Σ_ξ^0 (resp., Π_ξ^0) sets, on BC , if $\xi \geq 3$.

Proof. Theorem 4.1 gives a partial function F^1 . Recall that $F^1_1(\gamma)$ codes a continuous bijection defined on a closed subset of ω^ω if $\gamma \in BC$. We now express the fact that its graph is a closed subset of $\omega^\omega \times \omega^\omega$ (see the notation after Lemma 5.5). In Theorem 4.1, the complement of $\rho^{2^\omega}(\gamma)$ is involved. This leads us to use the map u_- given by Lemma 3.1. There is $P \in \Pi_1^0[(\omega^\omega)^3]$ such that

$$(\gamma, \beta, \alpha) \in P \Leftrightarrow \alpha \in 2^\omega \quad \text{and} \quad \beta \notin \rho^{\omega^\omega} \left(F^1_0[u_-(\gamma)] \right) \quad \text{and} \quad \alpha = \{F^1_1[u_-(\gamma)]\}^{\omega^\omega, 2^\omega}(\beta)$$

if $\gamma \in BC$. By 4A.1 in [Mos80] there is $\tilde{R} \in \Delta_1^0(\omega^\omega \times \omega^2)$ such that

$$(\gamma, \beta, \alpha) \in P \Leftrightarrow \forall k \in \omega \quad [\gamma, \overline{\beta[k]}, \overline{\alpha[k]}] \in \tilde{R}$$

(see the notation after Lemma 5.5).

We say that $(t, s) \in \tilde{Q}_f$ if $(t, s) \in Q$, $[\gamma, \bar{t}, \bar{s}] \in \tilde{R}$, $t \neq \emptyset$ and $t(|t| - 1) = 1$ (we use again the definition of Q_f after Lemma 5.5, but here it is uniform in γ). Now we define π as we did in section 2, with “ $q_{p_i} \in \tilde{Q}_f$ ” instead of “ $q_{p_i} \in Q_f$ ”. After a coding of $4^{<\omega}$ with ω , we can define a recursive map $\tilde{\varphi} : \omega^\omega \rightarrow 2^\omega$ coding $\mu \cup \pi \subseteq 4^{<\omega}$ (we will identify $\tilde{\varphi}(\gamma)$ with $\mu \cup \pi$, identifying ω with $4^{<\omega}$; the notation $\tilde{\varphi}$ instead of φ is for ω in the range of $\tilde{\varphi}$ instead of $2^{<\omega}$ in the range of φ).

Now let $\gamma \in BC$. Then $u_-(\gamma) \in BC$, $F^1[u_-(\gamma)]$ is defined, $f : \neg \rho^{\omega^\omega} \left(F^1_0[u_-(\gamma)] \right) \rightarrow \rho^{2^\omega}(\gamma)$ is a bijection. The proof of Theorem 1.3.(1) shows that $[\tilde{\varphi}(\gamma)]^\infty$ is Σ_ξ^0 (resp., Π_ξ^0) if $\rho^{2^\omega}(\gamma)$ is Σ_ξ^0 (resp., Π_ξ^0), when $\xi \geq 3$. It also shows that $\varphi_{0,0} \left([\tilde{\varphi}(\gamma)]^\infty \cap K_{0,0} \right) = \rho^{2^\omega}(\gamma)$, so that $\rho^{2^\omega}(\gamma)$ is Σ_ξ^0 (resp., Π_ξ^0) if $[\tilde{\varphi}(\gamma)]^\infty$ is Σ_ξ^0 (resp., Π_ξ^0), when $\xi \geq 3$. \square

Corollary 6.2 *Let $B \in \Delta_1^1[(2^\omega)^2]$ and $3 \leq \xi < \omega_1$. Then there is $\psi : 2^\omega \rightarrow 2^{2^{<\omega}}$ continuous such that*

- (a) $\Sigma_\xi^B := \{\alpha \in 2^\omega \mid B_\alpha \in \Sigma_\xi^0\} = \psi^{-1}(\Sigma_\xi)$.
- (b) $\Pi_\xi^B := \{\alpha \in 2^\omega \mid B_\alpha \in \Pi_\xi^0\} = \psi^{-1}(\Pi_\xi)$.

Proof. (a) Let $\gamma_0 \in BC$ such that $B = \rho^{(2^\omega)^2}(\gamma_0)$. By Lemma 3.2, we get $\rho^{(2^\omega)^2}(\gamma_0)_\alpha = \rho^{2^\omega}[u_s^{2^\omega}(\gamma_0, \alpha)]$ for each $\alpha \in 2^\omega$. So we just have to set $\psi(\alpha) := \varphi[u_s^{2^\omega}(\gamma_0, \alpha)]$, using Corollary 6.1.

(b) The proof is similar. \square

Theorem 6.3 (Saint Raymond) *Let $1 \leq \xi < \omega_1$. Then there is $B \in \Delta_1^1[(2^\omega)^2]$ such that Σ_ξ^B is Π_1^1 -complete. Similarly, there is $B \in \Delta_1^1[(2^\omega)^2]$ such that Π_ξ^B is Π_1^1 -complete.*

Proof. Let $P \subseteq 2^\omega$ be a Π_1^1 -complete set, $G \in \Pi_2^0[(2^\omega)^2]$ such that $\neg P$ is the first projection of G , X in $\Delta_1^1(2^\omega) \setminus \Sigma_\xi^0$, and $B := \{(\alpha, \beta) \in (2^\omega)^2 \mid [\alpha, (\beta)_0, (\beta)_1] \in G \times X\}$. Then B is clearly Borel. If $\alpha \in P$, then $B_\alpha = \emptyset \in \Sigma_\xi^0$, so $\alpha \in \Sigma_\xi^B$. If $\alpha \notin P$, let $\beta_0 \in 2^\omega$ such that $(\alpha, \beta_0) \in G$, and $f : 2^\omega \rightarrow 2^\omega$ defined by $f(\gamma) := \langle \beta_0, \gamma \rangle$. Then $B_\alpha = \{\beta \in 2^\omega \mid [\alpha, (\beta)_0, (\beta)_1] \in G \times X\} \notin \Sigma_\xi^0$ since $X = f^{-1}(B_\alpha) \notin \Sigma_\xi^0$. Thus $\alpha \notin \Sigma_\xi^B$. We proved that $\Sigma_\xi^B = P$ is Π_1^1 -complete. We argue similarly for Π_ξ^B . \square

Remarks. (a) We actually proved that if $\xi \geq 3$ and $P \in \Pi_1^1(2^\omega)$, then there is $M \in \Pi_\xi^0[(2^\omega)^2]$ such that $P = \Sigma_\xi^M$. Similarly, there is $A \in \Sigma_\xi^0[(2^\omega)^2]$ such that $P = \Pi_\xi^A$.

(b) This proof also shows that if $P \in \Pi_2^1(2^\omega)$, then there is $M \in \Pi_1^1[(2^\omega)^2]$ such that $P = \Sigma_\xi^M$. Similarly, there is $A \in \Pi_1^1[(2^\omega)^2]$ such that $P = \Pi_\xi^A$.

(c) This proof also shows that if $P \in \Pi_2^1(2^\omega)$, then there is $C \in \Pi_1^1[(2^\omega)^2]$ such that $P = \{\alpha \in 2^\omega \mid C_\alpha \in \Delta_1^1\}$.

Corollary 6.4 *Let $3 \leq \xi < \omega_1$. Then Σ_ξ and Π_ξ are Π_1^1 -hard (and also $\Sigma_2^1(2^{2^{<\omega}}) \setminus \Sigma_1^1$).*

Proof. We just have to apply Theorem 6.3 and Corollary 6.2. \square

Remark. Recall that if X is a recursively presented Polish space and $\beta \in 2^\omega$, then

$$\Sigma_1^1(\beta)(X) := \{Q_\beta \mid Q \in \Sigma_1^1(2^\omega \times X)\},$$

$\Pi_1^1(\beta) := \check{\Sigma}_1^1(\beta)$ and $\Delta_1^1(\beta) := \Sigma_1^1(\beta) \cap \Pi_1^1(\beta)$. In [Lec05], the following sets are introduced:

$$\Sigma_\xi := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Sigma_\xi^0 \cap \Delta_1^1(A)\},$$

$$\Pi_\xi := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Pi_\xi^0 \cap \Delta_1^1(A)\}.$$

It is proved in [Lec05] that they are $\Pi_1^1 \setminus \Delta_1^1$ if $\xi \geq 2$. Under the axiom of Σ_1^1 -determinacy, this implies that they are Π_1^1 -complete. Here we can say more: they are Π_1^1 -complete if $\xi \geq 3$, without axiom of determinacy. Indeed, fix a Π_1^1 -complete set $\Pi \subseteq 2^\omega$. The proof of Theorem 6.3 gives $B \in \Delta_1^1[(2^\omega)^2]$ such that $B_\alpha = \emptyset$ if $\alpha \in \Pi$, and $B_\alpha \notin \Sigma_\xi^0$ if $\alpha \notin \Pi$. Now the proof of Corollary 6.2 gives γ_0 . If $\alpha \in \Pi$, then $\rho^{2^\omega}[u_s^{2^\omega}(\gamma_0, \alpha)] = \emptyset$, and the proof of Theorem 1.3.(1) shows that $[\psi(\alpha)]^\infty = (\varphi[u_s^{2^\omega}(\gamma_0, \alpha)])^\infty = \mu^\infty \in \Pi_2^0 \subseteq \Delta_1^1$. Thus $\psi(\alpha) \in \Sigma_\xi$ if $\alpha \in \Pi$. If $\alpha \notin \Pi$, then $\psi(\alpha) \notin \Sigma_\xi$, thus $\psi(\alpha) \notin \Sigma_\xi$. Therefore $\Pi = \psi^{-1}(\Sigma_\xi)$ and Σ_ξ is Π_1^1 -hard. As Σ_ξ is Π_1^1 , it is Π_1^1 -complete. We argue similarly for Π_ξ .

Definition 6.5 *Let Γ be a class, and $\mathcal{U}_\Gamma^{2^\omega} \subseteq (2^\omega)^2$ universal for $\Gamma(2^\omega)$. We say that $\mathcal{U}_\Gamma^{2^\omega}$ is a good universal for Γ if for each set $\mathcal{U}_\Gamma^{(2^\omega)^2} \subseteq (2^\omega)^3$ which is universal for $\Gamma[(2^\omega)^2]$, there is $S : (2^\omega)^2 \rightarrow 2^\omega$ continuous such that $[S(\alpha, \beta), \gamma] \in \mathcal{U}_\Gamma^{2^\omega} \Leftrightarrow (\alpha, \beta, \gamma) \in \mathcal{U}_\Gamma^{(2^\omega)^2}$ for each $(\alpha, \beta, \gamma) \in (2^\omega)^3$.*

Proposition 6.6 *Let $1 \leq \xi < \omega_1$. Then there are good universals for Σ_ξ^0 , Π_ξ^0 , Σ_1^1 and Π_1^1 .*

Proof. Let Γ be one of the classes of the statement, and $\mathcal{V}_{\Gamma}^{(2^\omega)^2}$ universal for $\Gamma[(2^\omega)^2]$. We define, for $\alpha \in 2^\omega$ and $\varepsilon \in 2$, $(\alpha)_\varepsilon \in 2^\omega$ by $(\alpha)_\varepsilon(n) := \alpha(2n + \varepsilon)$. We set

$$\mathcal{U}_{\Gamma}^{2^\omega} := \{(\alpha, \beta) \in (2^\omega)^2 \mid [(\alpha)_0, (\alpha)_1, \beta] \in \mathcal{V}_{\Gamma}^{(2^\omega)^2}\}.$$

It is clear that $\mathcal{U}_{\Gamma}^{2^\omega} \in \Gamma$, so that $\{(\mathcal{U}_{\Gamma}^{2^\omega})_\alpha \mid \alpha \in 2^\omega\} \subseteq \Gamma(2^\omega)$. Conversely, let $A \in \Gamma(2^\omega)$. Then the set $E := \{(\gamma, \beta) \in (2^\omega)^2 \mid \beta \in A\} \in \Gamma$, so there is $\alpha \in 2^\omega$ such that $E = (\mathcal{V}_{\Gamma}^{(2^\omega)^2})_\alpha$. We define $\langle \cdot, \cdot \rangle : (2^\omega)^2 \rightarrow 2^\omega$ by $\langle \alpha, \beta \rangle(2n) := \alpha(n)$ and $\langle \alpha, \beta \rangle(2n+1) := \beta(n)$. We get $A = (\mathcal{U}_{\Gamma}^{2^\omega})_{\langle \alpha, 0^\infty \rangle}$. We proved that $\mathcal{U}_{\Gamma}^{2^\omega}$ is universal for $\Gamma(2^\omega)$.

Now let $\mathcal{U}_{\Gamma}^{(2^\omega)^2}$ be universal for $\Gamma[(2^\omega)^2]$, and

$$F := \{(\beta, \gamma) \in (2^\omega)^2 \mid [(\beta)_0, (\beta)_1, \gamma] \in \mathcal{U}_{\Gamma}^{(2^\omega)^2}\}.$$

As $F \in \Gamma[(2^\omega)^2]$, there is $\alpha_0 \in 2^\omega$ such that $F = (\mathcal{V}_{\Gamma}^{(2^\omega)^2})_{\alpha_0}$. We get

$$\begin{aligned} (\alpha, \beta, \gamma) \in \mathcal{U}_{\Gamma}^{(2^\omega)^2} &\Leftrightarrow \langle \alpha, \beta \rangle, \gamma \in F \\ &\Leftrightarrow (\alpha_0, \langle \alpha, \beta \rangle, \gamma) \in \mathcal{V}_{\Gamma}^{(2^\omega)^2} \\ &\Leftrightarrow \langle \langle \alpha_0, \langle \alpha, \beta \rangle \rangle, \gamma \rangle \in \mathcal{U}_{\Gamma}^{2^\omega} \end{aligned}$$

So we just have to set $S(\alpha, \beta) := \langle \alpha_0, \langle \alpha, \beta \rangle \rangle$. □

Lemma 6.7 *We consider the good universal $\mathcal{U}_{\Sigma_1^1}^{2^\omega}$ for Σ_1^1 given by Proposition 6.6. Then there is a continuous map $c: 2^{2^{<\omega}} \rightarrow 2^\omega$ such that $A^\infty = (\mathcal{U}_{\Sigma_1^1}^{2^\omega})_{c(A)}$ for each $A \in \mathcal{P}(2^{<\omega}) \cong 2^{2^{<\omega}}$.*

Proof. Recall that $\mathcal{U}_{\Sigma_1^0}^{2^\omega} \subseteq (2^\omega)^2$ is universal for $\Sigma_1^0(2^\omega)$ and defined in the proof of Theorem 3.14 as follows:

$$(\gamma, \alpha) \in \mathcal{U}_{\Sigma_1^0}^{2^\omega} \Leftrightarrow \exists m \in \omega \ \gamma(m) = 0 \text{ and } \alpha \in N[2^\omega, m].$$

Similarly, we can define $\mathcal{U}_{\Sigma_1^0}^{2^\omega \times \omega^\omega} \subseteq (2^\omega)^2 \times \omega^\omega$, universal for $\Sigma_1^0(2^\omega \times \omega^\omega)$:

$$(\gamma, \alpha, \beta) \in \mathcal{U}_{\Sigma_1^0}^{2^\omega \times \omega^\omega} \Leftrightarrow \exists m \in \omega \ \gamma(m) = 0 \text{ and } (\alpha, \beta) \in N[2^\omega \times \omega^\omega, m].$$

Using this, we can define $\mathcal{V}_{\Sigma_1^1}^{2^\omega} \subseteq (2^\omega)^2$, universal for $\Sigma_1^1(2^\omega)$:

$$(\gamma, \alpha) \in \mathcal{V}_{\Sigma_1^1}^{2^\omega} \Leftrightarrow \exists \beta \in \omega^\omega \ (\gamma, \alpha, \beta) \notin \mathcal{U}_{\Sigma_1^0}^{2^\omega \times \omega^\omega}.$$

By [Lec05] there is a continuous map $\pi : 2^\omega \times \omega^\omega \times \omega \rightarrow 2^{<\omega}$ such that

$$\alpha \in A^\infty \Leftrightarrow \exists \beta \in \omega^\omega \ \forall n \in \omega \ [\beta(n+1) > 0 \text{ and } \pi(\alpha, \beta, n) \in A],$$

for each $\alpha \in 2^\omega$ and $A \subseteq 2^{<\omega}$. We define $R \in \Sigma_1^0(2^\omega \times \omega^\omega \times 2^{<\omega})$ by

$$(\alpha, \beta, A) \in R \Leftrightarrow \exists n \in \omega \ [\beta(n+1) = 0 \text{ or } \pi(\alpha, \beta, n) \notin A].$$

By 3C.5 in [Mos80], there is $R^* \subseteq \omega$ such that

$$(\alpha, \beta, A) \in R \Leftrightarrow \exists m \in \omega \left(\alpha \in N[2^\omega, (m)_1] \text{ and } \beta \in N[\omega^\omega, (m)_2] \text{ and } A \in N[2^{<\omega}, (m)_3] \text{ and } m \in R^* \right).$$

We define $d : 2^{<\omega} \rightarrow 2^\omega$ by $d(A)(m) = 0 \Leftrightarrow A \in N[2^{<\omega}, (m)_3]$ and $m \in R^*$. If $A \subseteq 2^{<\omega}$, then

$$\begin{aligned} \alpha \in (\mathcal{V}_{\Sigma_1^1}^{2^\omega})_{d(A)} &\Leftrightarrow \exists \beta \in \omega \ [d(A), \alpha, \beta] \notin \mathcal{U}_{\Sigma_1^1}^{2^\omega \times \omega^\omega} \\ &\Leftrightarrow \exists \beta \in \omega \ \neg \left(\exists m \in \omega \ d(A)(m) = 0 \text{ and } (\alpha, \beta) \in N[2^\omega \times \omega^\omega, m] \right) \\ &\Leftrightarrow \exists \beta \in \omega \ \neg \left(\exists m \in \omega \ d(A)(m) = 0 \text{ and } \alpha \in N[2^\omega, (m)_1] \text{ and } \beta \in N[\omega^\omega, (m)_2] \right) \\ &\Leftrightarrow \exists \beta \in \omega \ (\alpha, \beta, A) \notin R \\ &\Leftrightarrow \alpha \in A^\infty \end{aligned}$$

As $\mathcal{V}_{\Sigma_1^1}^{2^\omega} \in \Sigma_1^1[(2^\omega)^2]$, there is $\alpha_0 \in 2^\omega$ such that $\mathcal{V}_{\Sigma_1^1}^{2^\omega} = (\mathcal{U}_{\Sigma_1^1}^{(2^\omega)^2})_{\alpha_0}$. As $\mathcal{U}_{\Sigma_1^1}^{2^\omega}$ is a good universal, we get S continuous, and $(\mathcal{V}_{\Sigma_1^1}^{2^\omega})_{d(A)} = (\mathcal{U}_{\Sigma_1^1}^{2^\omega})_{S[\alpha_0, d(A)]}$. So we just have to set $c(A) := S[\alpha_0, d(A)]$. \square

Recall that $\mathcal{U}(\Gamma, \Gamma') := \{\alpha \in 2^\omega \mid (\mathcal{U}_\Gamma^{2^\omega})_\alpha \in \Gamma'\}$ and $\Delta := \{A \subseteq 2^{<\omega} \mid A^\infty \in \Delta_1^1\}$.

Corollary 6.8 *Let $3 \leq \xi < \omega_1$. We consider the good universals given by Proposition 6.6.*

(a) *The set $\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0)$ is Π_1^1 -complete, $\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0) \leq_W \Sigma_\xi <_W \mathcal{U}(\Sigma_1^1, \Sigma_\xi^0)$, and the set $\mathcal{U}(\Sigma_1^1, \Sigma_\xi^0)$ is Π_2^1 -hard and $\Sigma_3^1 \setminus \Sigma_2^1$.*

(b) *The set $\mathcal{U}(\Sigma_\xi^0, \Pi_\xi^0)$ is Π_1^1 -complete, $\mathcal{U}(\Sigma_\xi^0, \Pi_\xi^0) \leq_W \Pi_\xi <_W \mathcal{U}(\Sigma_1^1, \Pi_\xi^0)$, and the set $\mathcal{U}(\Sigma_1^1, \Pi_\xi^0)$ is Π_2^1 -hard and $\Sigma_3^1 \setminus \Sigma_2^1$.*

(c) *$\Delta <_W \mathcal{U}(\Sigma_1^1, \Delta_1^1)$, and the set $\mathcal{U}(\Sigma_1^1, \Delta_1^1)$ is Π_2^1 -hard and $\Sigma_3^1 \setminus \Sigma_2^1$. Moreover, the set $\mathcal{U}(\Sigma_1^1, \Delta_1^1)$ is Π_2^1 -complete.*

Proof. (a) By Theorem 6.3 and Remark (a) just after, there is $M \in \Pi_\xi^0[(2^\omega)^2]$ such that Σ_ξ^M is Π_1^1 -complete. Fix $\alpha_0 \in 2^\omega$ with $M = (\mathcal{U}_{\Pi_\xi^0}^{(2^\omega)^2})_{\alpha_0}$. We define $f : 2^\omega \rightarrow 2^\omega$ by $f(\alpha) := S(\alpha_0, \alpha)$, where S is provided by the fact that $\mathcal{U}_{\Pi_\xi^0}^{2^\omega}$ is a good universal. Then we get $\Sigma_\xi^M = f^{-1}(\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0))$, which proves that $\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0)$ is Π_1^1 -hard. By [Lou80] (or 35.H in [Kec95]), $\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0)$ is Π_1^1 , so it is Π_1^1 -complete.

By Corollary 6.2, we get $\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0) \leq_W \Sigma_\xi$ since

$$\mathcal{U}(\Pi_\xi^0, \Sigma_\xi^0) = \Sigma_\xi \overset{\mathcal{U}_{\Pi_\xi^0}^{2^\omega}}{\leq_W} \Sigma_\xi.$$

By Lemma 6.7 we get $\Sigma_\xi \leq_W \mathcal{U}(\Sigma_1^1, \Sigma_\xi^0)$. Remark (b) after Theorem 6.3 gives $\Sigma := \neg A \in \Sigma_1^1[(2^\omega)^2]$ such that Σ_ξ^Σ is Π_2^1 -complete. The beginning of the proof shows that $\mathcal{U}(\Sigma_1^1, \Sigma_\xi^0)$ is Π_2^1 -hard. In particular, $\mathcal{U}(\Sigma_1^1, \Sigma_\xi^0) \notin \Sigma_2^1$, and $\Sigma_\xi <_W \mathcal{U}(\Sigma_1^1, \Sigma_\xi^0)$ since $\Sigma_\xi \in \Sigma_2^1$. Finally, $\mathcal{U}(\Sigma_1^1, \Sigma_\xi^0)$ is Σ_3^1 since

$$\alpha \in \mathcal{U}(\Sigma_1^1, \Sigma_\xi^0) \Leftrightarrow \exists \beta \in 2^\omega (\mathcal{U}_{\Sigma_1^1}^{2^\omega})_\alpha = (\mathcal{U}_{\Sigma_\xi^0}^{2^\omega})_\beta.$$

(b) The proof is very similar to that of (a).

(c) The proof of the first sentence is very similar to that of (a), using Remark (c) after Theorem 6.3. This proof shows that $\mathcal{U}(\Sigma_1^1, \Delta_1^1)$ is Π_2^1 -hard. It remains to see that $\mathcal{U}(\Sigma_1^1, \Delta_1^1)$ is Π_2^1 . Recall the existence of Π_1^1 sets $W^{2^\omega} \subseteq \omega$, $C^{2^\omega} \subseteq \omega \times 2^\omega$ with $\Delta_1^1(2^\omega) = \{C_n^{2^\omega} \mid n \in W^{2^\omega}\}$ and

$$\{(n, \alpha) \in \omega \times 2^\omega \mid n \in W^{2^\omega} \text{ and } \alpha \notin C_n^{2^\omega}\} \in \Pi_1^1(\omega \times 2^\omega)$$

(see Theorem 3.3.1 in [HKL90]). This implies that

$$\alpha \in \mathcal{U}(\Sigma_1^1, \Delta_1^1) \Leftrightarrow \exists n \in W^{2^\omega} (\mathcal{U}_{\Sigma_1^1}^{2^\omega})_\alpha = C_n^{2^\omega}.$$

Thus $\mathcal{U}(\Sigma_1^1, \Delta_1^1)$ is Π_2^1 , and Π_2^1 -complete. □

Acknowledgements. We would like to thank J. Saint Raymond for allowing us to present here one of his results (see Theorem 6.3).

References

- [Arn83] A. Arnold, Topological Characterizations of Infinite Behaviours of Transition Systems, Automata, Languages and Programming, J. Diaz Ed., Lecture Notes in Computer Science, Volume 154, Springer, 1983, p. 28-38.
- [ABB96] J-M. Autebert, J. Berstel and L. Boasson, Context Free Languages and Pushdown Automata, in Handbook of Formal Languages, Vol 1, Springer Verlag 1996.
- [Dup01] J. Duparc, Wadge Hierarchy and Veblen Hierarchy: Part 1: Borel Sets of Finite Rank, Journal of Symbolic Logic, Vol. 66, no. 1, 2001, p. 56-86.
- [DF06] J. Duparc and O. Finkel, An ω -Power of a Context-Free Language Which Is Borel Above Δ_ω^0 , to appear in the Proceedings of the International Conference Foundations of the Formal Sciences V : Infinite Games, November 26th to 29th, 2004, Bonn, Germany, (S. Bold, B. Löwe, T. Räscher, J. van Benthem eds.), College Publications at King's College (Studies in Logic), 2007.
- [Fin01] O. Finkel, Topological Properties of Omega Context Free Languages, Theoretical Computer Science, Vol. 262 (1-2), July 2001, p. 669-697.
- [Fin03] O. Finkel, Borel Hierarchy and Omega Context Free Languages, Theoretical Computer Science, Vol. 290 (3), 2003, p. 1385-1405.

- [Fin04] O. Finkel, An omega-Power of a Finitary Language Which is a Borel Set of Infinite Rank, *Fundamenta Informaticae*, Volume 62 (3-4), 2004, p. 333-342.
- [Fin06] O. Finkel, Borel Ranks and Wadge Degrees of Omega Context Free Languages, *Mathematical Structures in Computer Science*, Volume 16 (5), 2006, p. 813-840.
- [FL07] O. Finkel and D. Lecomte, There Exist some ω -Powers of Any Borel Rank, in the Proceedings of the 16th EACSL Annual Conference on Computer Science and Logic, Lausanne, Switzerland, September 11-15, 2007, (J. Duparc and T. Henzinger eds), *Lecture Notes in Computer Science*, Volume 4646, Springer, 2007, p. 115-129.
- [HKL90] L. A. Harrington, A. S. Kechris, and A. Louveau, A Glimm-Effros dichotomy for Borel equivalence relations, *Journal of the American Mathematical Society*, Volume 3, 1990, p. 903-928.
- [HU69] J.E. Hopcroft and J.D. Ullman, *Formal Languages and their Relation to Automata*, Addison-Wesley Publishing Company, Reading, Massachusetts, 1969.
- [Kec95] A.S. Kechris, *Classical Descriptive Set Theory*, Springer-Verlag, 1995.
- [KMS89] A. S. Kechris, D. Marker, and R. L. Sami, Π_1^1 Borel Sets, *The Journal of Symbolic Logic*, Volume 54 (3), 1989, p. 915-920.
- [Kur66] K. Kuratowski, *Topology*, Vol. 1, Academic Press, New York 1966.
- [Lec01] D. Lecomte, Sur les Ensembles de Phrases Infinies Constructibles a Partir d'un Dictionnaire sur un Alphabet Fini, *Séminaire d'Initiation a l'Analyse*, Volume 1, Université Paris 6, année 2001-2002.
- [Lec05] D. Lecomte, Omega-Powers and Descriptive Set Theory, *Journal of Symbolic Logic*, Volume 70 (4), 2005, p. 1210-1232.
- [LT94] H. Lescow and W. Thomas, Logical Specifications of Infinite Computations, In: "A Decade of Concurrency" (J. W. de Bakker et al., eds), *Lecture Notes in Computer Science*, Volume 803, Springer, 1994, p. 583-621.
- [Lev79] A. Levy, *Basic Set Theory*, Springer-Verlag, 1979.
- [Lou80] A. Louveau, A Separation Theorem for Σ_1^1 Sets, *Transactions of the American Mathematical Society*, Volume 260, 1980, p. 363-378.
- [Lou??] A. Louveau, Handwritten Notes.
- [LStR88] A. Louveau and J. Saint Raymond, Les Propriétés de Réduction et de Norme pour les Classes de Boréliens, *Fundamenta Mathematicae*, Volume 131 (3), 1988, p. 223-243.
- [Mos80] Y. N. Moschovakis, *Descriptive Set Theory*, North-Holland, Amsterdam 1980.
- [Niw90] D. Niwinski, Problem on ω -Powers posed in the Proceedings of the 1990 Workshop "Logics and Recognizable Sets" (Univ. Kiel).

- [PP04] D. Perrin and J.-E. Pin, Infinite Words, Automata, Semigroups, Logic and Games, Volume 141 of Pure and Applied Mathematics, Elsevier, 2004.
- [Sim92] P. Simonnet, Automates et Théorie Descriptive, Ph. D. Thesis, Université Paris 7, March 1992.
- [Sta86] L. Staiger, Hierarchies of Recursive ω -Languages, Jour. Inform. Process. Cybernetics EIK 22 (1986) 5/6, 219-241.
- [Sta97a] L. Staiger, ω -Languages, Chapter of the Handbook of Formal Languages, Vol 3, edited by G. Rozenberg and A. Salomaa, Springer-Verlag, Berlin, 1997.
- [Sta97b] L. Staiger, On ω -Power Languages, in New Trends in Formal Languages, Control, Cooperation, and Combinatorics, Lecture Notes in Computer Science 1218, Springer-Verlag, Berlin 1997, 377-393.
- [Tho90] W. Thomas, Automata on Infinite Objects, in: J. Van Leeuwen, ed., Handbook of Theoretical Computer Science, Vol. B (Elsevier, Amsterdam, 1990), p. 133-191.
- [vEn] F. van Engelen, On Borel Ideals, Annals of Pure and Applied Logic, Volume 70 (2), 1994, p.177-203.