



HAL
open science

Aging and effective temperatures near a critical point

Sylvain Joubaud, Baptiste Percier, Artem Petrosyan, Sergio Ciliberto

► **To cite this version:**

Sylvain Joubaud, Baptiste Percier, Artem Petrosyan, Sergio Ciliberto. Aging and effective temperatures near a critical point. 2008. ensl-00327521v1

HAL Id: ensl-00327521

<https://ens-lyon.hal.science/ensl-00327521v1>

Preprint submitted on 8 Oct 2008 (v1), last revised 6 Mar 2009 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Aging and effective temperatures near a critical point

S. Joubaud, B. Percier, A. Petrosyan and S. Ciliberto

*Université de Lyon, Laboratoire de Physique, Ecole Normale Supérieure de Lyon,
CNRS , 46, Allée d'Italie, 69364 Lyon CEDEX 07, France*

(Dated: October 8, 2008)

The orientation fluctuations of the director of a liquid crystal(LC) are measured after a quench near the Fréedericksz transition, which is a second order transition driven by an electric field. We report experimental evidence that, because of the critical slowing down, the LC presents, after the quench, several properties of an aging system, such as power law scaling versus time of correlation and response functions. During this slow relaxation, a well defined effective temperature, much larger than the heat bath temperature, can be measured using the fluctuation dissipation relation.

PACS numbers: 05.40.-a, 05.70.Jk, 02.50.-r, 64.60.-i

The characterization of the thermodynamic properties of out of equilibrium and of slow relaxing systems is an important problem of great current interest which is studied both theoretically and experimentally. One of the main questions which has been investigated in the recent years is how a temperature can be defined for these out of equilibrium systems, mainly in the context of aging glasses, whose relaxation is nonexponential and extremely slow. Indeed when a glass is quenched from above to below the glass transition temperature its physical properties depend on the waiting time t_w spent after the quench at the low temperature T . For these materials it has been proposed that a "good definition" (in the thermodynamic sense) of an effective temperature, T_{eff} can be obtained by the so called fluctuation dissipation relation (FDR) which is an extension of the fluctuation dissipation theorem (FDT) for an out of equilibrium system [1]. This T_{eff} is a function of the age of the system and of the observational time scale. To be more precise, let us consider an observable θ , its conjugated variable Γ and the integrated response, $\chi(t, t_w)$, of $\theta(t)$, at time t , to a Heaviside perturbation given to Γ at a time $t_w < t$. This integrated response is related to the correlation function $C(t, t_w)$ of the thermal fluctuations $\delta\theta$ of θ by the FDR in the following way:

$$\chi(t, t_w) = \frac{1}{k_B T_{\text{eff}}(t, t_w)} [C(t, t) - C(t, t_w)] \quad (1)$$

where k_B is the Boltzmann constant. Of course at equilibrium $T_{\text{eff}} = T$ and both $\chi(t, t_w)$ and $C(t, t_w)$ depend only on $t - t_w$. For several models of aging dynamics it has been shown that the T_{eff} defined from eq. 1 is characterized by only two temperatures and a characteristic time, t^* , such that for $(t - t_w) > t^*$ then $T_{\text{eff}} > T$ whereas for $(t - t_w) < t^*$, $T_{\text{eff}} = T$ [1, 2]. However in experiments the results are less clear and the relevance for real materials of the T_{eff} as defined by eq. 1 is still an open question, which merits to be investigated [3, 4, 5]. The point is that in general it is very difficult to find simple theoretical models which can be directly compared with experiments. The recent theoretical works on aging at

critical point are very useful in this sense. Indeed it has been recently shown that a system which is quenched at the critical point of a second order phase transition shares several properties observed in the relaxation dynamics of an aging glass [6, 7, 8]. For this reason this kind of behavior has been named "aging at critical point". The theoretical predictions on this critical systems are very useful because they can be a real benchmark to compare theory with experiment. The purpose of this letter is to report on experimental results on a critical system, which is, on a first approximation, described by a Ginzburg-Landau equation. Thus many results can be directly compared to theory but at the same time as all new experimental measurements they open new theoretical perspectives. From a general point of view the main result of our investigation is that after a quench at the critical point the system presents a violation of the fluctuation dissipation theorem and that for long correlation times a well defined T_{eff} can be defined from eq. 1.

The system of our interest is the Fréedericksz transition of a liquid crystal (LC), submitted to an electric field \vec{E} [9, 10]. In this system, we measure the variable ζ which is the spatially averaged alignment of the LC molecules, whose local direction of alignment is defined by the unit vector \vec{n} . Let us first recall the general properties of the Fréedericksz transition. The system under consideration is a LC confined between two parallel glass plates at a distance $L = 9 \mu\text{m}$. The inner surfaces of the confining plates have transparent Indium-Tin-Oxyde (ITO) electrodes, used to apply the electric field. Furthermore the plate surfaces, are coated by a thin layer of polymer mechanically rubbed in one direction. This surface treatment causes the alignment of the LC molecules in a unique direction parallel to the surface (planar alignment), *i.e.* all the molecules have the same director parallel to x -axis. The cell is next filled by a LC having a positive dielectric anisotropy (p-pentylcyanobiphenyl, 5CB, produced by Merck). The LC is submitted to an electric field perpendicular to the plates, by applying a voltage V between the ITO electrodes, *i.e.* $E = V/L$. In order to avoid electrodynamic effects of the

motion of the ions invariably present in the liquid crystal, we apply an AC voltage at a frequency $f_V = 1$ kHz ($V = \sqrt{2}V_0 \cos(2\pi f_V t)$) [9, 10]. More details on the experimental set-up can be found in ref.[11, 12]. When V_0 exceeds a threshold value V_c the planar state becomes unstable and the LC molecules, except those anchored to the glass surfaces, try to align parallel to the field, *i.e.* the director, away from the confining plates, acquires a component parallel to the applied electric field (z -axis). This is the Fréedericksz transition whose properties are those of a second order phase transition [9, 10]. For V_0 close to V_c the motion of the director is characterized by its angular displacement θ in xz -plane (θ is the angle between the x axis and \vec{n}), whose space-time dependence has the following form : $\theta = \theta_0(x, y, t) \sin\left(\frac{\pi z}{L}\right)$ [9, 10, 13]. If θ_0 remains small then its dynamics is described by a Ginzburg-Landau equation and one expects mean-field critical phenomena [9, 10, 13], in which θ_0 is the order parameter and $\epsilon = \frac{V_0^2}{V_c^2} - 1$ is the reduced control parameter. We define the variable ζ as the spatially averaged alignment of the LC molecules and more precisely :

$$\zeta = \frac{2}{L} \frac{1}{\mathcal{A}} \iint_{\mathcal{A}} dx dy \int_0^L (1 - n_x^2) dz \quad (2)$$

where $\mathcal{A} = \pi D_0^2/4$ is the area, in the (x, y) plane, of the measuring region of diameter D_0 , which is about 2 mm in our case. If θ_0 remains small, ζ takes a simple form in terms of θ_0 ($n_x = \cos(\theta)$) : $\zeta = 1/\mathcal{A} \iint_{\mathcal{A}} dx dy \theta_0^2$. The variable ζ is measured using the anisotropic properties of the LC, *i.e.* the cell is a birefringent plate whose local optical axis is parallel to \vec{n} . This optical anisotropy can be precisely estimated by measuring the phase shift Φ between two linearly polarized beams which cross the cell, one polarized along x -axis (ordinary ray) and the other along the y -axis (extraordinary ray). It can be shown that $\zeta = a + b\Phi$ where a and b are known coefficients (see ref. [11, 12] for details). The phase Φ , measured by an interferometer [14], is acquired with a resolution of 24 bits at a sampling rate of 16384 Hz and then low pass filtered at 500 Hz in order to suppress the AC voltage at f_V . The instrumental noise of the apparatus is three orders of magnitude smaller than the amplitude $\delta\Phi$ of the fluctuations of Φ induced by the thermal fluctuations of ζ . It has been shown that ζ is characterized by a mean value $\langle \zeta \rangle \propto \epsilon$, by a divergent relaxation time $\tau = \tau_0/\epsilon$ and by fluctuations which have a Lorentzian spectrum [12, 15].

In this letter, we consider the dynamic of ζ as a function of time after a quench from $\epsilon = \epsilon_1 > 0.3$ to $\epsilon = \epsilon_0 \ll 0.1$. The fact that the control parameter is the electric field allows one to have extremely fast quenches, typically 2 ms. Furthermore the typical relaxation time τ of ζ is about 10s, at $\epsilon_0 \simeq 0$. This means that we can follow the out of equilibrium dynamics for about 4 order of magnitude in time. This relaxation dynamics, which lasts only a few seconds, allows us to repeat the experi-




FIG. 1: a) Logarithm of the mean relaxation of the system, $\log\langle\zeta(t)\rangle - \langle\zeta(\infty)\rangle$, as a function of time after a quench (\circ) from $\epsilon_1 = 0.3$ to $\epsilon_0 = 0.01$. The continuous line represent the theoretical response based on eq. 3 with $\langle\zeta\rangle \simeq \psi_0^2$. Dashed line is an exponential relaxation. b) Logarithm of the mean relaxation of the system as a function of the logarithm of time. The continuous line represent the theoretical relaxation.

ment several times and to perform an ensemble average (indicated by $\langle . \rangle$) of the measured quantities. We describe here a specific case of a quench from $\epsilon_1 = 0.3$ to $\epsilon_0 = 0.01$ (more details for other quenches will be given in a longer reports). The typical mean value $\langle \zeta(t) \rangle$ of ζ after this quench is plotted in fig. 1 as a function of time t ($t = 0$ is the time when the quench has been performed). This mean dynamics of ζ is obtained by repeating the same quench several times (7000 times). The behavior of $\langle \zeta \rangle$ remains constant for a certain time and then slowly relaxes (see fig. 1b). Above a characteristic time which is about 0.2 s in our case, the relaxation becomes exponential. This behavior can be modelled by the equation of momentum [13], which neglecting the dependence on (x, y) of θ_0 is:

$$\frac{\gamma \mathcal{A} L}{2} \frac{d\theta_0}{dt} = B [2 \epsilon \theta_0 - (\kappa + \epsilon + 1) \theta_0^3] + \eta \quad (3)$$

where γ is a viscosity coefficient, $B = \mathcal{A}\pi^2 K_1/4L$, $\kappa = (K_3 - K_1)/K_1$ and K_1, K_2 and K_3 are the three elastic constants of the LC. η is a thermal noise delta-correlated in time, such that: $\langle \eta(t) \rangle = 0$ and $\langle \eta(t_1)\eta(t_2) \rangle = k_B T \gamma \mathcal{A} L \delta(t_2 - t_1)$. We decompose the dynamics of θ_0 in a mean dynamics after the quench and its fluctuations, *i.e.* $\theta_0 = \psi_0(t) + \delta\theta_0$, where $\psi_0(t) = \langle \theta_0(t) \rangle$. The dynamics of ψ_0 obtained from the analytical solution of eq.3 is shown in fig.1 as a continuous line, which perfectly agrees with the measured dynamics. The experimental data and the solution of eq.3 have two well distinguish limits : for $t \gg \tau \equiv \tau_0/(2\epsilon)$, the relaxation is exponential ($\tau \simeq 0.22s$ at $\epsilon_0 = 0.01$) ; for $t \ll \tau \equiv \tau_0/(2\epsilon)$, the dynamics of ψ_0 is almost algebraic : $\psi_0(t) = \left[(\kappa + 1) \left(\frac{t}{\tau_0} + \frac{1}{(\kappa+1)\psi_0(0)} \right) \right]^{-1/2}$. This behavior is identical to the relaxation at the critical point [7]. Thus, the system should present aging phenomena in the limit $t \ll \tau$.

In our experimental system, we want to measure both the correlation function of the thermal fluctuations $\delta\theta_0$ of θ_0 , $C_\theta(t, t_w) \equiv \langle \delta\theta_0(t)\delta\theta_0(t_w) \rangle$, and the integrated re-




FIG. 2: a) Correlations functions $C_\theta(t, t_w)$ as a function of $t - t_w$ for different fixed $t = 0.2$ (\circ), 0.26 (\square), 0.28 (\diamond) and 0.31 (\times) after the same quench of fig. 1. Inset: The correlation functions have a simple master curve obtained by plotting $C_\theta(t, t_w)/C_\theta(t, t_w)$ versus $(t - t_w)/t$.

sponse function $\chi(t, t_w)$ of θ_0 , at time t to a perturbation, given at time t_w , to its conjugated external field, with $t_w < t$. In equilibrium $\chi(t, t_w)$ and $C_\theta(t, t_w)$ depends only on the difference $t - t_w$ which is not the case here because the system is out of equilibrium.

To compute $\chi(t, t_w)$ and $C_\theta(t, t_w)$ in our experiment one has to consider that the measured variable is ζ . As the area of the measuring beam is much larger than the correlation length, the global variable measured by the interferometer is $\zeta = \frac{1}{A} \iint \theta_0^2 dx dy \simeq \psi_0^2 + 2\psi_0\delta\theta_0$. Thus the mean value of ζ is $\langle \zeta(t) \rangle = \psi_0^2(t)$ and the fluctuations of $\delta\zeta$ of ζ can be related to the fluctuations of θ_0 : $\delta\zeta(t) = 2\psi_0(t)\delta\theta_0(t)$. The autocorrelation function of θ_0 is obtained using the values of $\psi_0(t)$ and $\psi_0(t_w)$, *i.e.* $C_\theta(t, t_w) = \langle \zeta(t)\zeta(t_w) \rangle / (4\psi_0(t)\psi_0(t_w))$. The $C_\theta(t, t_w)$, measured at various fixed t , are plotted as function of $t - t_w$ in fig.2. We see that a simple scaling of the correlation function can be obtained by plotting them as a function of $(t - t_w)/t$. Notice that such a kind of scaling is typical for aging systems(see for example [2, 16]) Thus in agreement with theoretical predictions our system presents aging properties during the slow relaxation after a quench. The response function is obtained by applying a small change of the voltage V_0 , which modifies the control parameter, *i.e.* $\epsilon = \epsilon_0 + \delta\epsilon$. First, we have to define the external torque conjugated to θ_0 . We separate $\theta_0(t)$ into the average part $\psi_0(t)$, solution of eq. 3 and a deviation $\Delta(t)$ due to $\delta\epsilon$: $\theta_0(t) = \psi_0(t) + \Delta(t)$. If the response is linear, the average value of Δ is proportional to $\delta\epsilon$. In this limit, the equation of momentum can be rewritten :

$$\frac{\gamma AL}{2} \frac{d\Delta}{dt} = B [2\epsilon_0 - (\kappa + \epsilon_0 + 1)3\psi_0^2] \Delta + 2B\delta\epsilon\psi_0(t) \left(1 - \frac{\psi_0^2(t)}{2}\right) + \eta + O(\Delta^2) \quad (4)$$

The external torque, and so the conjugated variable to θ_0 , is equal to :

$$\Gamma_{\text{ext}} = 2B\delta\epsilon\psi_0(t) \left(1 - \frac{\psi_0^2(t)}{2}\right) \quad (5)$$




FIG. 3: Parametric plot integrated response versus correlation for $t_w = 0.19$ s (\circ), 0.31 s (\square), 0.37 s (\diamond), 0.43 s ($+$), 0.56 s (\times). Continuous lines are linear fits. Dashed line represents the Fluctuation-Dissipation Theorem at equilibrium. Inset : value of the effective temperature, obtained by a linear fit of the FD plots, as a function of the age of the system.

We define the linear response function $R(t, t_w) = \langle \Delta \rangle / \Gamma_{\text{ext}}$ of θ_0 to Γ_{ext} using a Dirac delta function for $\delta\epsilon$ at an instant t_w , *i.e.* $\delta\epsilon(t) \propto \delta(t - t_w)$. In the experiment the Dirac delta function is approximated by a triangular function of amplitude $\delta\epsilon_0$ and duration $\tau_r \simeq 2\text{ms} \ll \tau$, specifically: $\delta\epsilon = \delta\epsilon_0(1 - 2|t - t_w|/\tau_r)$ for $|t - t_w| < \tau_r/2$ and $\delta\epsilon = 0$ for $|t - t_w| > \tau_r/2$, with $\tau_r/\tau \simeq 10^{-4}$. The measured quantity is the response $R_{\zeta, \delta\epsilon} = \langle \Delta\zeta \rangle / \delta\epsilon$, of $\zeta(t)$ to $\delta\epsilon(t_w)$. As $\Delta\zeta(t) = 2\psi_0(t)\Delta$ it follows that the linear response of θ_0 to Γ_{ext} is:

$$R(t, t_w) = \frac{R_{\zeta, \delta\epsilon}(t, t_w)}{4B\psi_0(t_w)\psi_0(t) \left(1 - \frac{\psi_0^2(t_w)}{2}\right)} \quad (6)$$

For getting the integrated response function χ , we apply the perturbation at different times after the quench. Then we measure in the experiment $R_{\zeta, \delta\epsilon}$ and the value of $\psi_0(t)$ as a function of time. These measured values are replaced in eq. 6 to compute $R(t, t_w)$. The integrated response χ is obtained by a numerical integration of $R(t, t_w)$: $\chi(t, t_w) = \int_t^{t_w} R(t, t') dt'$ The integrated response is related to the correlation by the FDR relations (see eq. 1). We study the FDR during an aging process by plotting the integrated response χ as a function of C_θ . Due to the non-equilibrium process, it is not equivalent if the parameter is the waiting time t_w or the observation time t .

First, t_w is kept fixed. We can see in fig. 3 the FD plot for four typical values of the age of the system t_w . χ is a linear function of C_θ as stated by eq. 1 with a slope $1/T_{\text{eff}}$ When t_w increases, the slope T_{eff} tends to the temperature of the room T as can be seen in fig. 3. The Fluctuation-Dissipation Theorem will be satisfied for large t_w . In the inset of figure 3, we have plotted the value of T_{eff} as a function of time t_w . We can see that T_{eff} decreases with t_w and tends to the ambient temperature when t_w tends to infinity.




FIG. 4: a) Parametric plot integrated response versus correlation for $t = 0.4$ s (\circ), 0.5 s (\square), 0.6 s (\diamond) and 0.7 s ($+$). Continuous lines are linear fits. Dark line represents the FDT at equilibrium. Inset : the characteristic time t^* is plotted as a function of t .

In a second time, t is kept constant, which is the original way proposed by Cugliandolo and Kurchan and in theoretical papers on the aging at critical point [1, 7]. The FDR plots can be seen in figure 4 for four characteristics value of t . When t_w is close to t (when the correlation function is close to 1), the FDT is satisfied. Below a value, C^* , which depends on t , the Fluctuation Dissipation Theorem is not satisfied. The integrated linear response function remains linear in C but the slope is almost three times the value at equilibrium. More precisely, if we define t^* the characteristic time associated to the value of C^* . For $C < C^*$, R remains linear in C and the slope is not related to the thermal bath temperature T but to a temperature T_{eff} , which is independent of t and almost twice T . This means that the slow dynamics is characterized by a well defined temperature. In the inset of figure 4, we have plotted the value of t^* as a function of t . We find that above a value of 0.22 s, t^* is linear with t . The value 0.22 s corresponds to the relaxation time at the system at ϵ_0 . As simple picture can explain this observation : we separate the time scale between an aging part ($0 < t_w < 0.22$) and an equilibrium part ($t > 0.22$ s). t^* is defined as the time between the equilibrium part and the aging one, thus it is equal to $t^* = t - 0.22$. This behavior matches very well the experimental result for $t > 0.22$ s. The dependence of T_{eff} on the initial and final value of the quench is an important feature which will be discussed in a longer report.

In conclusion we have shown that a LC quenched close to the critical point of the Fredericks transition presents aging features, such as a power law rescaling of correlation functions and the appearance of a well defined $T_{eff} > T$ (working at fixed t). What is very interesting here is that although we are not exactly at $\epsilon = 0$ we observe a large interval of time where the predicted aging at critical point can be observed. The results plotted in fig.4 are not in perfect agreement with those of ref. [7] for the mean field calculation of the aging at the

critical point. The reason of this discrepancy is probably related to the fact that eq. 3 is an approximation for the dynamics of our system for which the spatial behavior cannot be neglected. This opens the discussion for further theoretical and experimental development. Finally we want to stress that the results of fig.4 are in excellent quantitative agreement with the general predictions for the T_{eff} as defined by Cugliandolo and Kurchan. They show the existence of a very well defined effective temperature independent of t for the slow dynamics $t - t_w$ larger than a t^* . Notice that this result is in contrast with that of fig.3 where we see that the temperature is a decreasing function of t_w . This different behavior obtained by keeping fixed either t or t_w has been widely discussed in the theoretical literature, but it has never been so clearly observed in an experiment. This observation is actually very useful because in the study of FDR in polymers and colloids the procedure, which is often applied for several technical problems related to the measurement noise [3, 4] is the one with t_w fixed (fig.3). This of course may provoke several problems when comparing the experimental results with theory and our experimental results clearly point out this problem.

We acknowledge useful discussion with Peter Holdsworth. This work has been partially supported by ANR-05-BLAN-0105-01.

-
- [1] L. Cugliandolo, J. Kurchan and L. Peliti, *Phys. Rev. E*, **55**(4) 3898 (1997)
 - [2] L. Cugliandolo Cugliandolo L F, *Dynamics of glassy systems*, Lecture notes (Les Houches, July 2002)
 - [3] T.S. Grigera and N.E. Israeloff, *Phys. Rev. Lett.*, **83**, 5038(1999)
 - [4] L. Bellon, S. Ciliberto, C. Laroche *Europhys. Lett.*, **53**(4), 511 (2001); L. Buisson and S. Ciliberto *Physica D*, **204**(1-2) 1 (2005)
 - [5] D. Hérisson and M. Ocio, *Phys. Rev. Lett.*, **88**(25), 257202 (2002)
 - [6] L. Berthier, P.C.W. Holdsworth and M. Sellitto, *J. Phys. A: Math. Gen.*, **34**, pp.1805 – 1824 (2001)
 - [7] P. Calabrese, A. Gambassi and F. Krzakala, *J. Stat. Mech. : Theory and Experiment*, **P06016** (2006)
 - [8] S. Abriet, D. Karevski, *E.P.J. B*, **37**(1), 47-53 (2004)
 - [9] P.G. de Gennes and J. Prost, (Clarendon Press, 1974)
 - [10] P. Oswald and P. Pieranski, *Nematic and cholesteric liquid crystals* (Taylor & Francis, 2005)
 - [11] S. Joubaud, G. Huilard, A. Petrosyan and S. Ciliberto, to be published in *J. Stat. Mech.*
 - [12] S. Joubaud, A. Petrosyan, S.Ciliberto, N. Garnier, 2008 *Phys. Rev. Lett.* **100** 180601
 - [13] M. San Miguel, *Phys. Rev. A* **32**(6) 3811 (2005)
 - [14] L. Bellon, S. Ciliberto, H. Boubaker, L. Guyon, *Optics Communications* **207** 49-56 (2002).
 - [15] P. Galatola and M. Rajteri, 1994 *Phys. Rev.* **49** 623.
 - [16] E. Vincent, in *Ageing and the Glass Transition* edited by M. Henkel, M. Pleimling, R. Sanctuary (Springer-Verlag 2007).