



HAL
open science

Shallow Circuits with High-Powered Inputs

Pascal Koiran

► **To cite this version:**

| Pascal Koiran. Shallow Circuits with High-Powered Inputs. 2010. ensl-00477023v1

HAL Id: ensl-00477023

<https://ens-lyon.hal.science/ensl-00477023v1>

Preprint submitted on 27 Apr 2010 (v1), last revised 29 Jul 2010 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Shallow Circuits with High-Powered Inputs

Pascal Koiran

LIP*, École Normale Supérieure de Lyon, Université de Lyon

Department of Computer Science, University of Toronto[†]

`Pascal.Koiran@ens-lyon.fr`

April 27, 2010

Abstract

A polynomial identity testing algorithm must determine whether an input polynomial (given for instance by an arithmetic circuit) is identically equal to 0. In this paper, we show that a deterministic black-box identity testing algorithm for (high-degree) univariate polynomials would imply a lower bound on the arithmetic complexity of the permanent. The lower bounds that are known to follow from derandomization of (low-degree) multivariate identity testing are weaker.

To obtain our lower bound it would be sufficient to derandomize identity testing for polynomials of a very specific norm: sums of products of sparse polynomials with sparse coefficients. This observation leads to new versions of the Shub-Smale τ -conjecture on integer roots of univariate polynomials. In particular, we show that a lower bound for the permanent would follow if one could give a good enough bound on the number of real roots of sums of products of sparse polynomials (Descartes' rule of signs gives such a bound for sparse polynomials and products thereof).

*UMR 5668 ENS Lyon, CNRS, UCBL, INRIA.

[†]A part of this work was done during a visit to the Fields Institute.

1 Introduction

A polynomial identity testing algorithm must determine whether an input polynomial (given for instance by an arithmetic circuit) is identically equal to 0. If randomization is allowed, this problem can be solved efficiently thanks to the well-known Schwarz-Zippel lemma. Following Kabanets and Impagliazzo [11], it has become increasingly clear in recent years that efficient deterministic algorithms for polynomial identity testing would imply strong lower bounds (the connection between arithmetic circuit lower bounds and derandomization of polynomial identity testing was foreshadowed in a 30 years old paper by Heintz and Schnorr [10]). This approach to lower bounds was advocated in particular by Agrawal [1].

In this paper we show that an efficient black-box deterministic identity testing algorithm for univariate polynomials of a very specific form (namely, sums of products of sparse polynomials with sparse coefficients) would imply that the permanent does not belong to VP^0 . This is the class of polynomial families computable by constant-free arithmetic circuits of polynomial size and polynomially bounded formal degree. It plays roughly the same role for constant-free circuits as the class VP in Valiant's algebraic version of the P versus NP problem (in Valiant's original setting, arithmetic circuits can use arbitrary constants from the underlying field [8, 26]).

Compared to [1, 11], the main originality of the present paper is to show that lower bound for multivariate polynomials such as the permanent would follow from *univariate* identity testing algorithms. Most of the recent work on identity testing (surveyed in [2, 23]) has been focused on low-degree multivariate polynomials.¹ Nevertheless, we believe that the univariate approach is worth exploring for at least two reasons.

First, it would lead to stronger lower bounds. Indeed, we show that black-box derandomization of identity testing implies a lower bound for the permanent, whereas [1, Section 6.2] would only yield lower bounds for polynomials with coefficients computable in PSPACE (this complexity class was independently defined in [18], where it is called VPSPACE ; further results on this class and other space-bounded classes in Valiant's model can be found in [17, 22, 20]). The lower bound obtained from [11] would be even weaker, but could be obtained from a non-black-box identity testing algorithm.

A second, possibly even more important advantage of the univariate approach is that it leads to new (and hopefully more tractable) versions of Shub and Smale's τ -conjecture. According to the τ -conjecture, the number

¹Two exceptions are [7, 15].

of integer roots of a univariate polynomial $f \in \mathbb{Z}[X]$ should be bounded by a polynomial function of its arithmetic circuit size (the inputs to the circuit are the constant 1, or the variable X). It was shown by Bürgisser [9] that the τ -conjecture implies a lower bound for the permanent. Our main “hardness from derandomization” result can be viewed as an improvement of Bürgisser’s result. Indeed, it follows immediately from our result that to obtain a lower bound for the permanent, one just has to bound the number of integer roots for sums of products of sparse polynomials with sparse coefficients (rather than for arbitrary arithmetic circuits). Our strongest version of the τ -conjecture raises the intriguing possibility that tools from real analysis might be brought to bear on this problem (a bound on the number of real roots of a polynomial is a fortiori a bound on its number of integer roots). It is known that this approach cannot work for the original τ -conjecture because the number of real roots of a univariate polynomial can grow exponentially as a function of its arithmetic circuit size (Chebyshev polynomials provide such an example [25]). We conjecture that this behavior is not possible for sums of products of sparse polynomials.

1.1 Main Ideas

A hitting set \mathcal{H} for a set \mathcal{F} of polynomials is a (finite) set of points such that there exists for any non-identically zero polynomial $f \in \mathcal{F}$ at least one point $a \in \mathcal{H}$ such that $f(a) \neq 0$. Hitting sets are sometimes called “correct test sequences” [10]. It is well-known that deterministic constructions of hitting sets and black-box deterministic identity testing are two equivalent problems: any hitting set for \mathcal{H} yields an obvious black-box identity testing algorithm (declare that $f \equiv 0$ iff f evaluates to 0 on all the points of \mathcal{H}); conversely, assuming that \mathcal{F} contains the identically zero polynomial, the set of points queried by a black box algorithm on the input $f \equiv 0$ must be a hitting set for \mathcal{F} .

The connection between black-box identity testing and lower bounds is especially apparent for univariate polynomials [10]. Namely, let \mathcal{H} be a hitting set for \mathcal{F} . The polynomial

$$P = \prod_{a \in \mathcal{H}} (X - a) \tag{1}$$

cannot belong to \mathcal{F} since it is nonzero and vanishes on \mathcal{H} . The same remark applies to all nonzero multiples of P . If \mathcal{F} is viewed as some kind of “complexity class”, we have therefore obtained a lower bound against \mathcal{F} by exhibiting a polynomial P which does not belong to \mathcal{F} .

In the low-degree multivariate setting the polynomial which plays the same role is not given by such a simple formula as (1). Its coefficients can be obtained by solving an exponential size system of linear equations. This can be done in PSPACE, explaining why the lower bound in [1] would be for polynomials with coefficients computable in PSPACE. By contrast one can show that the coefficients in exponential-size products such as (1) are in the counting hierarchy, a subclass of PSPACE. This is the reason why we can obtain a lower bound for a polynomial in VNP (namely, the permanent) rather than in VPSPACE as in [1, Section 6.2].

It remains to explain why we only have to derandomize identity testing for sums of products of sparse polynomials in order to obtain a lower bound. This class of polynomials comes into the picture thanks to the recent depth reduction theorem of Agrawal and Vinay [3]: any multilinear polynomial in n variables which has an arithmetic circuit of size $2^{o(n)}$ also has a depth-4 arithmetic circuit of size $2^{o(n)}$. Sums of products of sparse polynomials are very far from being multilinear (they are univariate polynomials of possibly very high degree). They are nonetheless connected to depth-4 circuits by a simple transformation: if we replace the input variables in a depth-4 circuit by powers of a single variable X , we obtain a SPS polynomial $f(X)$.

At this point, we should stress that we do *not* claim that univariate arithmetic circuits can be efficiently converted into SPS polynomials (this would be a kind of high-degree analogue of Agrawal and Vinay's depth reduction theorem). On the contrary, we conjecture that such a transformation is in general impossible, and that Chebyshev polynomials provide a counterexample (because, as pointed out earlier, they have too many real roots). Nevertheless, to obtain our results we represent efficiently (in Theorem 6) certain exponential size products by sums of products of sparse polynomials. *This is possible only under the assumption that the permanent is easy.* This assumption (and the resulting representation) is of course very likely to be false, but there is no harm in making it since the ultimate goal is a proof by contradiction that the permanent is hard.

1.2 Organization of the Paper

In the next section we present our model of computation for the permanent (constant-free arithmetic circuits) as well as the corresponding complexity classes. We also recall some definitions and results about the counting hierarchy (as explained above, this class plays a crucial role in the derivation of a lower bound for the permanent). In Section 3 we define precisely the notion of sum of products of sparse polynomials with sparse coefficients, and ex-

plain the connection to depth-4 circuits. In Section 4 we present the notion of algebraic number generator. This is basically just a sequence of efficiently computable polynomials in $\mathbb{Z}[X]$. We wish to use them to construct hitting sets, by taking the sets of all roots of the polynomials in an initial segment of this sequence. In Section 5 we prove our main result: if a polynomial-size initial segment provides a hitting set against sums of products of sparse polynomials with sparse coefficients, then the permanent is not in VP^0 . In Section 6 we present three new versions of the τ -conjecture. A proof of any of these conjectures would yield a lower bound for the permanent. We conclude the paper with a few remarks on some tools that might be useful to attack these conjectures.

2 Preliminaries

2.1 Arithmetic Circuits

We recall that an arithmetic circuit contains addition, subtraction and multiplication gates. We usually assume that these gates have arity 2, except when dealing with constant-depth circuits as in e.g. Theorem 3. The input gates are labelled by variables or constants. A circuit where the only constants are from the set $\{0, -1, 1\}$ is said to be constant-free (in such a circuit one can even assume that -1 is the only constant, and that there are no subtraction gates). A constant-free circuit represents a polynomial in $\mathbb{Z}[X_1, \dots, X_n]$, where X_1, \dots, X_n are the variables labelling the input gates.

In this paper we investigate the complexity of computing the permanent polynomial with constant-free arithmetic circuits. This model of computation was systematically studied by Malod [21]. In particular, he defined a class VP^0 of polynomial families that are “easy to compute” by constant-free arithmetic circuits. First we need to recall the notion of formal degree:

- (i) The formal degree of an input gate is equal to 1.
- (ii) The formal degree of an addition or subtraction gate is the maximum of the formal degrees of its two incoming gates, and the formal degree of a multiplication gate is the sum of these two formal degrees.

Finally, the formal degree of a circuit is equal to the formal degree of its output gate. This is obviously an upper bound on the degree of the polynomial computed by the circuit.

Definition 1 *A sequence (f_n) of polynomials belongs to VP^0 if there exists a polynomial $p(n)$ and a sequence (C_n) of constant-free arithmetic circuits*

such that C_n computes f_n and is of size (number of gates) and formal degree at most $p(n)$.

The size constraint implies in particular that f_n depends on polynomially many variables. The constraint on the formal degree forbids the computation of polynomials of high degree such as e.g. X^{2^n} ; it also forbids the computation of large constants such as 2^{2^n} .

A central question in the constant-free setting is whether the permanent family belongs to VP^0 . A related question is whether $\tau(\text{PER}_n)$, the constant-free arithmetic circuit of the $n \times n$ permanent, is polynomially bounded in n . Obviously, if $\text{PER} \in \text{VP}^0$ then $\tau(\text{PER}_n)$ is polynomially bounded in n , but (as pointed out in e.g. [8]) it is not clear whether the converse holds true. In this paper we focus on the first question (see section 7 for further comments).

Another important complexity class in the constant-free setting is the class VNP^0 of easily definable families. It is obtained from VP^0 in the natural way:

Definition 2 *A sequence $(f_n(X_1, \dots, X_{u(n)}))$ belongs to VNP^0 if there exists a sequence $(g_n(X_1, \dots, X_{v(n)}))$ in VP^0 such that:*

$$f_n(X_1, \dots, X_{u(n)}) = \sum_{\bar{\epsilon} \in \{0,1\}^{v(n)-u(n)}} g_n(X_1, \dots, X_{u(n)}, \bar{\epsilon}).$$

For instance, the permanent family is in VNP^0 . If this family in fact belongs to VP^0 then the same is true of every VNP^0 family up to constant multiplicative factors. Indeed, we have the following result (Theorem 4.3 of [14]):

Theorem 1 *Assume that the permanent family is in VP^0 . For every family (f_n) in VNP^0 there exists a polynomially bounded function $p(n)$ such that the family $(2^{p(n)} f_n)$ is in VP^0 .*

The occurrence of the factor $2^{p(n)}$ in this theorem is due to the fact that the completeness proof of the permanent uses the constant $1/2$. As in [14] one could avoid this factor by working with the Hamiltonian polynomial instead of the permanent.

The next lemma is Valiant's criterion. The present formulation is basically that of [14, Th. 2.3] but this lemma essentially goes back to [26] (see also [8, Prop. 2.20]).

Theorem 2 (Valiant’s criterion) *Suppose that $n \mapsto p(n)$ is a polynomially bounded function, and that $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{Z}$ is such that the map $1^n 0j \mapsto f(j, n)$ is in the complexity class GapP/poly . Then the family (f_n) of multilinear polynomials defined by*

$$f_n(X_1, \dots, X_{p(n)}) = \sum_{j \in \{0,1\}^{p(n)}} f(j, n) X_1^{j_1} \dots X_{p(n)}^{j_{p(n)}} \quad (2)$$

is in VNP^0 . Here j_k denotes the bit of j of weight 2^{k-1} .

Note that we use a unary encoding for n but a binary encoding for j . We recall the definition of GapP/poly (and a few other boolean complexity classes) in Section 2.2. In this paper we only need to apply Valiant’s criterion to boolean-valued functions ($f(j, n) \in \{0, 1\}$ for all j and n) such that the map $1^n 0j \mapsto f(j, n)$ is in P/poly .

Finally, we will use a recent result of Agrawal and Vinay: low-degree multivariate polynomials which admit nontrivial arithmetic circuits also admit nontrivial arithmetic circuits of depth four. We will apply their depth reduction theorem to multilinear polynomials only. In this case their result (Corollary 2.5 in [3]) reads as follows:

Theorem 3 (Reduction to depth four) *A multilinear polynomial in n variables which has an arithmetic circuit of size $2^{o(n)}$ also has a depth 4 arithmetic circuit of size $2^{o(n)}$.*

The resulting depth 4 circuits are $\sum \prod \sum \prod$ arithmetic formulas: the output gate (at depth 4) and the gates at depth 2 are addition gates, and the other gates are multiplication gates.

2.2 The Counting Hierarchy

A connection between the counting hierarchy and algebraic complexity theory was discovered in [4]. This connection was further explored in [9] and [16]. For instance, it was shown in [9] that the polynomials $\prod_{i=0}^{2^n} (X - i)$ have polynomial-size circuits if the same is true for the permanent family.

We first recall the definition of the two counting classes $\sharp\text{P}$ and GapP .

Definition 3 *The class $\sharp\text{P}$ is the set of functions $f : \{0, 1\}^* \rightarrow \mathbb{N}$ such that there exist a language $A \in \text{P}$ and a polynomial $p(n)$ satisfying*

$$f(x) = \#\{y \in \{0, 1\}^{p(|x|)} : (x, y) \in A\}.$$

A function $f : \{0, 1\}^ \rightarrow \mathbb{Z}$ is in GapP if it is the difference of two $\sharp\text{P}$ functions.*

The counting hierarchy introduced in [27] is a class of languages rather than functions. It is defined via the majority operator \mathbf{C} as follows.

Definition 4 *If K is a complexity class, the class $\mathbf{C}.K$ is the set of languages A such that there exist a language $B \in K$ and a polynomial $p(n)$ satisfying*

$$x \in A \iff \#\{y \in \{0,1\}^{p(|x|)} : (x,y) \in B\} \geq 2^{p(|x|)-1}.$$

The i -th level $\mathbf{C}_i\mathbf{P}$ of the counting hierarchy is defined recursively by $\mathbf{C}_0\mathbf{P} = \mathbf{P}$ and $\mathbf{C}_{i+1}\mathbf{P} = \mathbf{C}.\mathbf{C}_i\mathbf{P}$. The counting hierarchy \mathbf{CH} is the union of the levels $\mathbf{C}_i\mathbf{P}$ for all $i \geq 0$.

The counting hierarchy contains all the polynomial hierarchy \mathbf{PH} and is contained in \mathbf{PSPACE} .

The arithmetic circuit classes defined in Section 2.1 are nonuniform. As a result, we will actually work with nonuniform versions of the counting classes defined above. We use the standard Karp-Lipton notation [12]:

Definition 5 *If K is a complexity class, the class K/poly is the set of languages A such that there exist a language $B \in K$, a polynomial $p(n)$ and a family $(a_n)_{n \geq 0}$ of words (the "advice") satisfying*

- for all $n \geq 0$, $|a_n| \leq p(n)$;
- for all word x , $x \in A \iff (x, a(|x|)) \in B$.

Note that the advice only depends on the size of x .

The next lemma [9, Lemmas 2.6 and 2.13] provides a first link between arithmetic complexity and the counting hierarchy.

Lemma 1 *If the permanent family is in \mathbf{VP}^0 then $\mathbf{CH}/\text{poly} = \mathbf{P}/\text{poly}$.*

In particular, Lemma 1 was used to show that large sums and products are computable in the counting hierarchy [9, Theorem 3.10].

In the remainder of this section we summarize some relevant results from [16].

Definition 6 *Let (f_n) be a family of polynomials in $\mathbb{Z}[X]$ such that the degree of f_n and the bitsize of its coefficients are smaller than $2^{p(n)}$ for some polynomial p .*

The coefficient sequence of (f_n) is the (double) sequence of integers $a(n, \alpha)$ defined by the relation

$$f_n(x) = \sum_{\alpha=0}^{2^{p(n)}-1} a(n, \alpha)x^\alpha.$$

The coefficient sequence is said to be definable in CH/poly if the language $\text{Bit}(a) = \{(1^n, \alpha, j, b); \text{ the } j\text{-th bit of } a(n, \alpha) \text{ is equal to } b\}$ is in CH/poly.

Note that in the above definition of $\text{Bit}(a)$, the input n is given in unary but α and j are in binary (this is the same convention as in [16]; by contrast, in [9] all inputs are in binary).

Definition 7 Let (f_n) be a family of polynomials as in Definition 6. We say that this family can be evaluated in CH/poly if the language

$$\{(1^n, i, j, b); 0 \leq i < 2^{p(n)} \text{ and the } j\text{-th bit of } f_n(i) \text{ is equal to } b\}$$

is in CH/poly.

The following result establishes a connection between these two definitions. It is stated (and proved) in the proof of the main theorem (Theorem 3.5) of [16].

Theorem 4 Let (f_n) be a family of polynomials as in Definition 6. If (f_n) can be evaluated in CH/poly at integer points, the coefficient sequence of (f_n) is definable in CH/poly.

In [16] we actually prove a multivariate version of this result, but the univariate case will be sufficient for our purposes.

3 Sums of Products of Sparse Polynomials

A sum of products of sparse polynomials is an expression of the form $\sum_i \prod_j f_{ij}$ where each $f_{ij} \in \mathbb{Z}[X]$ is a sparse univariate polynomial. Here “sparse” means as usual that we only represent the nonzero monomials of each f_{ij} . As a result one can represent concisely polynomials of very high degree. We define the size of such an expression as the sum of the number of monomials in all the f_{ij} . Note that this measure of size does not take into account the size of the coefficients of the f_{ij} , or their degrees. These relevant parameters are taken into account in the following definition.

Definition 8 We denote by $\text{SPS}_{s,e}$ the set of all polynomials in $\mathbb{Z}[X]$ which can be represented by an expression of the form $\sum_i \prod_j f_{ij}$ so that:

- The size of the expression as defined above is at most s .
- Each coefficient of each f_{ij} can be written as the difference of two nonnegative integers with at most s nonzero digits in their binary representations.
- These coefficients are of absolute value at most 2^e , and the f_{ij} are of degree at most e .

Remark 1 The polynomials f_{ij} in this definition can be thought of as a "sparse polynomial with sparse coefficients". The integer s serves as a sparsity parameter for the number of monomials as well as for the number of digits in their coefficients. A typical choice for these parameters is $s = 2^{O(n)}$ and $e = 2^{O(n)}$, where n represents an input size (see for instance Theorem 6 in Section 5).

We will show in Section 5 that constructing polynomial size hitting sets for sums of products of sparse polynomials implies the lower bound $\text{PER} \not\subseteq \text{VP}^0$. Here "polynomial size" means polynomial in $s + \log e$. It is quite natural to insist on a size bound which is polynomial in s and $\log e$: s is an arithmetic circuit size bound, and $\log e$ can also be interpreted as an arithmetic cost since each power x^α in an f_{ij} can be computed from x in $O(\log e)$ operations by repeated squaring. Likewise, we can write each coefficient of each f_{ij} as the difference of two nonnegative integers as in Definition 1, and each of the $\leq s$ powers of 2 occurring in a nonnegative integer can be computed from the constant 2 in $O(\log e)$ operations. Each coefficient can therefore be computed in $O(s \log e)$ operations. As a result, a polynomial in $\text{SPS}_{s,e}$ can be evaluated from the constant 1 and the variable X in a number of arithmetic operations which is polynomial in $s + \log e$.

Remark 2 The size of a SPS polynomial as we have defined it is essentially the size of a depth three arithmetic circuit (or more precisely of a depth three arithmetic formula) computing the polynomial. In this depth three formula each input gate carries a monomial; each addition gate at level 1 computes a f_{ij} ; each multiplication gate at level 2 computes a product of the form $\prod_j f_{ij}$; and the output gate at level 3 computes the final sum.

We can further refine this representation of SPS polynomials by arithmetic circuits. Namely, instead of viewing the monomial aX^β as an atomic object which is fed to an input gate, we can decompose it as a sum of terms

of the form $\pm 2^\alpha X^\beta$; and each term can be further decomposed as a product of factors of the form $\pm 2^{2^i}$ and X^{2^j} . The resulting object is a depth four circuit where each input gate carries an expression of the form $\pm 2^{2^i}$ or x^{2^j} (note the symmetry between variables and constants in this representation). This connection between depth four circuits and SPS polynomials plays a crucial role in our results.

4 Algebraic number generators

As explained in Section 1.2, we wish to construct hitting sets by taking the sets of all roots of the polynomials in an initial segment of an efficiently computable sequence of polynomials. The following definition makes the notion of “efficiently computable” precise (compare with the notion of *hitting set generator* in [1, Section 6.2]).

Definition 9 *An algebraic number generator is a sequence $(f_i)_{i \geq 1}$ of nonzero univariate polynomials $f_i(X) = \sum_\alpha a(\alpha, i)X^\alpha$ such that for some integer constant $c \geq 1$:*

1. *The exponents α range from 0 to i^c ;*
2. *$a(\alpha, i)$ is a sequence of integers of absolute value $\leq 2^{i^c}$;*
3. *The language $L(f) = \{(\alpha, i, j, b); \text{ the } j\text{-th bit of } a(\alpha, i) \text{ is equal to } b\}$ is in CH/poly.*

In the above definition we work with the complexity class CH/poly because this is the largest complexity class for which our proofs go through. As shown in the next example, the language $L(f)$ can often be located in a much smaller complexity class.

Example 1 *Each of the three sequences $f_i = x - i$, $(x^i - 1)$ or $x^i - 2^i x + i^2 + 1$ is an algebraic number generator. Notice that in these three examples we can compute the coefficients of the f_i in polynomial time rather than in CH/poly, i.e., there is no need for counting and the construction of the f_i is uniform.*

Theorem 5 *Let (f_i) be an algebraic number generator. From this sequence we define a family of univariate polynomials g_n by the formula:*

$$g_n(x) = \prod_{i=1}^{2^n} f_i(x).$$

The coefficient sequence $b(n, \alpha)$ of g_n , defined by $g_n(x) = \sum_\alpha b(n, \alpha)x^\alpha$, is definable in CH/poly.

Proof. The family (g_n) can be evaluated in CH/poly at integer points. This follows from the fact that integer sequences definable in CH/poly are stable under products and summations [9, Theorem 3.10]. The result then follows from Theorem 4. \square

We illustrate this result on two examples.

Example 2 For $f_i = x - i$ we have $g_n(x) = \prod_{i=1}^{2^n} (x - i)$. This is the Pochhammer-Wilkinson polynomial of order 2^n . It was shown in [9, proof of Main Theorem 1.2] that the coefficient sequence of Pochhammer-Wilkinson polynomials is definable in CH.

Example 3 For $f_i = x^i - 1$ we have $g_n(x) = \prod_{i=1}^{2^n} (x^i - 1)$. This product can be written as

$$g_n(x) = \prod_{\bar{\epsilon}} h_n(x, \bar{\epsilon}) \quad (3)$$

where the auxiliary family h_n is defined by:

$$h_n(x, \epsilon_1, \dots, \epsilon_n) = x \prod_{j=1}^n [(1 - \epsilon_j) + \epsilon_j x^{2^{j-1}}] - 1.$$

Note that the powers $x^{2^{j-1}}$ in the above formula can be computed efficiently by repeated squaring. The family (h_n) therefore belongs to the class VP_{nb}^0 of polynomials that can be evaluated in a polynomial number of arithmetic operations in the constant-free unbounded-degree model. It then follows from (3) that g_n belongs to the class $\text{V}\Pi\text{P}^0$ (by definition, the families of this class are obtained as in (3) from a VP_{nb}^0 family by taking an exponential-size product over a VP_{nb}^0 family). It is shown in [16, Theorem 3.7] that the class $\text{V}\Pi\text{P}^0$ would collapse to VP_{nb}^0 if VNP^0 collapses to VP^0 . The proof of this theorem is based on definability of coefficients in CH/poly for $\text{V}\Pi\text{P}^0$ families (in our particular example there is again no need for nonuniformity since the family (f_i) is uniform).

5 From a Hitting Set to a Lower Bound

In this section we prove our main result: constructing hitting sets for the class $\text{SPS}_{s,e}$ of sums of products of sparse polynomials with sparse coefficients implies a lower bound for the permanent (recall that the class $\text{SPS}_{s,e}$ is defined in Section 3).

We begin with a lemma showing that under the assumption $\text{PER} \in \text{VP}^0$, polynomials with coefficients definable in CH/poly can be efficiently represented by sums of products of sparse polynomials. This result is an adaptation of [16, Lemma 3.2], which was itself a scaled up version of [9, Th. 4.1(2)]. The main new ingredient is the depth reduction theorem of [3].

Lemma 2 *Let $g_n(x) = \sum_{\alpha} a(n, \alpha)x^{\alpha}$ where the integers α range from 0 to $2^{c \cdot n} - 1$, $a(n, \alpha)$ is a sequence of integers of absolute value $< 2^{2^{c \cdot n}}$ definable in CH/poly , and c is an integer constant (independent of n).*

If $\text{PER} \in \text{VP}^0$ there is a polynomially bounded function $p(n)$ such that $2^{p(n)}g_n \in \text{SPS}_{s,e}$ where $s = 2^{o(n)}$ and $e = 2^{O(n)}$.

Proof. Expand a in binary: $a(n, \alpha) = \sum_{i=0}^{2^{c \cdot n} - 1} a_i(n, \alpha)2^i$. Let h_n be the following multilinear polynomial:

$$h_n(x_1, x_2, \dots, x_{c \cdot n}, z_1, \dots, z_{c \cdot n}) = \sum_{i=0}^{2^{c \cdot n} - 1} \sum_{\alpha=0}^{2^{c \cdot n} - 1} a_i(n, \alpha) z_1^{i_1} \dots z_{c \cdot n}^{i_{c \cdot n}} x_1^{\alpha_1} x_2^{\alpha_2} \dots x_{c \cdot n}^{\alpha_{c \cdot n}}.$$

Then we have:

$$h_n(x^{2^0}, x^{2^1}, \dots, x^{2^{c \cdot n - 1}}, 2^{2^0}, 2^{2^1}, \dots, 2^{2^{c \cdot n - 1}}) = g_n(x). \quad (4)$$

Assume that the permanent family is in VP^0 . by Lemma 1 the nonuniform counting hierarchy collapses, therefore computing the i -th bit $a_i(n, \alpha)$ on input $(1^n, \alpha, i)$ is in GapP/poly (and even in P/poly). By Lemma 2, $(h_n) \in \text{VNP}^0$. By Theorem 1 there exists a polynomially bounded function $p(n)$ such that the family $h'_n = 2^{p(n)}h_n$ is in VP^0 . By reduction to depth 4 (Theorem 3) we can compute h'_n by a depth four circuit C_n of size $2^{o(n)}$. The inputs to C_n are the variables of h'_n and the constant -1 (which we do need to compute non-monotone polynomials). Finally, by virtue of (4) we can plug powers of 2 and powers of x in h'_n to represent $2^{p(n)}g_n$ by an expression $\sum_i \prod_j f_{ij}$ of the required form. In particular, as already pointed out in Remark 2, after this substitution of inputs the multiplication gates in the first layer of C_n will compute expression of the $\pm 2^{\alpha} x^{\beta}$, and the addition gates in the second layers will compute sparse polynomials with sparse coefficients. Note that the sparseness property for the coefficients of the f_{ij} relies on the fact that depth reduction does not require the introduction of large integer constants in the resulting depth 4 circuits. \square

Next we show that the product of the first 2^n polynomials of an algebraic number generator can be represented by a sum of products of sparse polynomials of subexponential size, assuming again that the permanent is in VP^0 . Namely:

Theorem 6 *Let (f_i) be an algebraic number generator and $g_n(x) = \prod_{i=1}^{2^n} f_i(x)$. If $\text{PER} \in \text{VP}^0$ there is a polynomially bounded function $p(n)$ such that $2^{p(n)}g_n \in \text{SPS}_{s,e}$ where $s = 2^{o(n)}$ and $e = 2^{O(n)}$. Here $\text{SPS}_{s,e}$ is the class of sums of products of sparse polynomials from Definition 8.*

Proof. We wish to apply Lemma 2 to the polynomial $g_n(x) = \prod_{i=1}^{2^n} f_i(X)$. Each polynomial f_i in this product is of degree less than 2^{cn} (except possibly f_n , which may be of degree up to 2^{cn}). Hence g_n is of degree less than $2^{(c+1)n}$. As to the coefficient size, we have $\|g_n\|_1 \leq \prod_i \|f_i\|_1$ where $\|\cdot\|_1$ denotes the sum of the absolute values of the coefficients of a polynomial. For each i we have $\|f_i\|_1 < (2^{cn} + 1) \cdot 2^{2^{cn}} \leq 2^{2^{(c+1)n}}$ so that $\|g_n\|_1 \leq 2^{2^{(c+2)n}}$. Finally, definability of coefficients in CH/poly is provided by Theorem 5. \square

We can finally prove our main result.

Theorem 7 (Lower Bound from Hitting Sets) *Let (f_i) be an algebraic number generator and H_m the set of all roots of the polynomials f_i for all $i \leq m$. If there exists a polynomial p such that $H_{p(s+\log e)}$ is a hitting set for $\text{SPS}_{s,e}$ then the permanent is not in VP^0 .*

Proof. Let $g_n(x) = \prod_{i=1}^{2^n} f_i(x)$ be the polynomial of Theorem 6. Assume by contradiction that:

- (i) There exists a polynomial p such that $H_{p(s+\log e)}$ is a hitting set for $\text{SPS}_{s,e}$.
- (ii) The permanent family is in VP^0 .

From our second assumption and Theorem 6 we know that $2^{p(n)}g_n$ is in $\text{SPS}_{s,e}$ for $s = 2^{o(n)}$, $e = 2^{O(n)}$ and some polynomially bounded function $p(n)$. Hence H_m is a hitting set for g_n for $m = 2^{o(n)}$ since $s + \log e = 2^{o(n)}$. By construction, g_n vanishes on H_{2^n} and $H_m \subseteq H_{2^n}$ since $m \leq 2^n$ if n is large enough. This is a contradiction since g_n vanishes on the hitting set H_m but is not identically 0. \square

Remark 3 *The hitting sets that we consider in this theorem are initial segments of a single "universal" sequence of algebraic numbers. This requirement is not really restrictive: we can always assume that hitting sets have*

this form, by taking the union of all hitting sets up to a given value of $s + \log e$ if necessary (and this transformation does not blow up the size of the hitting sets by too much).

6 Hitting Sets from Real Analysis ?

In this section we present our new versions of the τ -conjecture. Each of the three conjectures implies that the permanent is not in VP^0 .

Conjecture 1 (τ -conjecture for SPS polynomials) *There is a polynomial p such that any nonzero polynomial in $\text{SPS}_{s,e}$ has at most $p(s + \log e)$ integer roots.*

This conjecture implies that $\text{PER} \notin \text{VP}^0$ (apply Theorem 7 to the algebraic number generator $f_i(x) = x - i$). Conjecture 1 follows from the τ -conjecture of Shub and Smale on integer roots of polynomials [24, 25] since, as explained after Definition 8, polynomials in $\text{SPS}_{s,e}$ can be evaluated by constant-free arithmetic circuits of size polynomial in s and $\log e$. It was already shown in [9] that the τ -conjecture implies a lower bound for the permanent. The point of Conjecture 1 is that to obtain such a lower bound we no longer have to bound the number of integer roots of arbitrary arithmetic circuits: we need only do this for sums of products of sparse polynomials. This looks like a much more manageable class of circuits, but the question is of course still wide open. Another related benefit of SPS polynomials in this context is that techniques from real analysis might become applicable. Before explaining this in more detail we formulate a somewhat stronger conjecture. The idea is that the parameter e in Conjecture 1 as well as the sparsity hypothesis on the integer coefficients might be irrelevant. This leads to:

Conjecture 2 (τ -conjecture for SPS polynomials, strong form)

Consider a polynomial of the form

$$f(X) = \sum_{i=1}^k \prod_{j=1}^m f_{ij}(X),$$

where each $f_{ij} \in \mathbb{Z}[X]$ has at most t monomials. The number of integer roots of f is bounded by a polynomial function of kmt .

Note that the size of f as defined in Section 3 is bounded by kmt . Therefore, Conjecture 2 is indeed stronger than Conjecture 1. Finally, we formulate an even stronger conjecture.

Conjecture 3 (real τ -conjecture) Consider a polynomial of the form

$$f(X) = \sum_{i=1}^k \prod_{j=1}^m f_{ij}(X),$$

where each $f_{ij} \in \mathbb{R}[X]$ has at most t monomials. The number of real roots of f is bounded by a polynomial function of kmt .

One could also formulate a weak version of the real τ -conjecture where the parameters s and e would play the same role as in Conjecture 1.

At present there isn't a lot of evidence for or against Conjecture 3. We do know that the conjecture holds true when $k = 1$: by Descartes' rule each polynomial f_{1j} has at most $2t - 2$ nonzero real roots, so f has at most $2m(t - 1) + 1$ real roots. The case $k = 2$ already looks nontrivial. In the general case we can expand f as a sum of at most kt^m monomials, so we have at most $2kt^m - 1$ real roots. A refutation of the conjecture would be interesting from the point of view of real algebra and geometry as it would yield examples of "sparse like" polynomials with many real roots. Of course, a proof of the conjecture would be even more interesting as it would yield a lower bound for the permanent.

Finally we recall that if true, Conjecture 3 would be a property that is really specific to SPS polynomials: as explained in the introduction, it is known that for general arithmetic circuits, the number of real roots cannot be bounded by a polynomial function of the circuit size.

7 Final Remarks

We have shown that constructing hitting sets for sums of products of sparse polynomials with sparse coefficients will show that $\text{PER} \notin \text{VP}^0$. It should be possible to obtain a variation of this result where the conclusion is that $\tau(\text{PER}_n)$, the constant-free arithmetic circuit complexity of the permanent, is not polynomial in n . To obtain this stronger conclusion, a stronger hypothesis should be necessary. It seems natural to expect that the role played by sparse polynomials with sparse coefficients will now be played by sparse polynomials with coefficients of "small" τ -complexity (this is a larger class of polynomials since sparse coefficients are certainly of small τ -complexity).

Most importantly, one should try to prove or disprove the real τ -conjecture. A solution in the case $k = 2$ (a sum of two products of sparse polynomials) would already be quite interesting. We note that the search for good upper bounds on the number of solutions of sparse multivariate

systems is a topic of current interest in real algebraic geometry. The theory of fewnomials [13] provides finiteness results and sometimes quantitative estimates on the number of real roots in very general “sparse like” situations. The general estimates from [13], at least when applied in a straightforward manner, do not seem to yield any useful bound for our problem. Nevertheless, one can hope that the methods developed in [13] as well as in more recent work such as [5, 6, 19] will turn out to be useful.

References

- [1] M. Agrawal. Proving lower bounds via pseudo-random generators. In *Proc. FSTTCS 2005*. Invited paper.
- [2] M. Agrawal and R. Saptharishi. Classifying Polynomials and Identity Testing. *Current Trends in Science*, 2009.
- [3] M. Agrawal and V. Vinay. Arithmetic circuits: A chasm at depth four. In *Proc. 49th IEEE Symposium on Foundations of Computer Science*, pages 67–75, 2008.
- [4] E. Allender, P. Bürgisser, J. Kjeldgaard-Pedersen, and P. Bro-Miltersen. On the complexity of numerical analysis. *SIAM Journal on Computing*, 38(5):1987–2006, 2009. Conference version in CCC 2006.
- [5] B. Bertrand, F. Bihan, and F. Sottile. Polynomial systems with few real zeroes. *Mathematische Zeitschrift*, 253(2):361–385, 2006.
- [6] F. Bihan and F. Sottile. New fewnomial upper bounds from Gale dual polynomial systems. *Moscow Mathematical Journal*, 7(3), 2007.
- [7] M. Bläser, M. Hardt, R. J. Lipton, and N. K. Vishnoi. Deterministically testing sparse polynomial identities of unbounded degree. *Information Processing Letters*, 109(3):187–192, 2009.
- [8] P. Bürgisser. *Completeness and Reduction in Algebraic Complexity Theory*. Number 7 in Algorithms and Computation in Mathematics. Springer, 2000.
- [9] P. Bürgisser. On defining integers and proving arithmetic circuit lower bounds. *Computational Complexity*, 18:81–103, 2009. Conference version in STACS 2007.

- [10] J. Heintz and C.-P. Schnorr. Testing polynomials which are easy to compute. In *Logic and Algorithmic (an International Symposium held in honour of Ernst Specker)*, pages 237–254. Monographie n° 30 de L’Enseignement Mathématique, 1982. Preliminary version in *Proc. 12th ACM Symposium on Theory of Computing*, pages 262–272, 1980.
- [11] V. Kabanets and R. Impagliazzo. Derandomizing polynomial identity test means proving circuit lower bounds. *Computational Complexity*, 13(1-2):1–46, 2004.
- [12] R. Karp and R. Lipton. Turing machines that take advice. *L’Enseignement Mathématique*, 28:191–209, 1982.
- [13] A. G. Khovanskii. *Fewnomials*, volume 88 of *Translations of Mathematical Monographs*. American Mathematical Society, 1991.
- [14] P. Koiran. Valiant’s model and the cost of computing integers. *Computational Complexity*, 13:131–146, 2004.
- [15] P. Koiran. A hitting set construction, with application to arithmetic circuit lower bounds. <http://arxiv.org/abs/0907.5575>, 2009.
- [16] P. Koiran and S. Perifel. Interpolation in Valiant’s theory, 2007. Available from <http://perso.ens-lyon.fr/pascal.koiran/publications.html>. To appear in *Computational Complexity*.
- [17] P. Koiran and S. Perifel. VPSACE and a transfer theorem over the complex field. In *Proc. 32nd International Symposium on Mathematical Foundations of Computer Science*, volume 4162 of *Lecture Notes in Computer Science*, pages 359–370, 2007.
- [18] P. Koiran and S. Perifel. VPSACE and a transfer theorem over the reals. *Computational Complexity*, 18:551–575, 2009. Conference version in STACS 2007.
- [19] T.-Y. Li, J. Maurice Rojas, and X. Wang. Counting real connected components of trinomial curve intersections and m -nomial hypersurfaces. *Discrete and Computational Geometry*, 30(3):379–414, 2003.
- [20] M. Mahajan and R. Rao. Small-space analogues of Valiant’s classes. In *Proc. 17th International Symposium on Fundamentals of Computation Theory*, pages 250–261. Springer, 2009.

- [21] G. Malod. *Polynômes et coefficients*. PhD thesis, Université Claude Bernard - Lyon 1, 2003.
- [22] B. Poizat. A la recherche de la définition de la complexité d'espace pour le calcul des polynômes à la manière de Valiant. *Journal of Symbolic Logic*, 73(4):1179–1201, 2008.
- [23] N. Saxena. Progress on Polynomial Identity Testing. *Bull. EATCS*, 99:49–79, 2009.
- [24] M. Shub and S. Smale. On the intractability of Hilbert's Nullstellensatz and an algebraic version of “P=NP”. *Duke Mathematical Journal*, 81(1):47–54, 1995.
- [25] S. Smale. Mathematical problems for the next century. *Mathematical Intelligencer*, 20(2):7–15, 1998.
- [26] L. G. Valiant. Completeness classes in algebra. In *Proc. 11th ACM Symposium on Theory of Computing*, pages 249–261, 1979.
- [27] Klaus W. Wagner. The complexity of combinatorial problems with succinct input representation. *Acta Inf.*, 23(3):325–356, 1986.