

Extracting Herbrand trees in classical realizability using forcing

Derivations of some proof terms

Lionel RIEG

Contents

1 Foreword	2
2 Absoluteness of connectives	4
2.1 Conjunction	4
2.2 Disjunction	7
2.3 Classical existential quantifier (on multiple propositions)	9
3 Properties of the generic set	11
3.1 G is a filter	11
3.1.1 G is non-empty	11
3.1.2 G is a subset of C	11
3.1.3 G is upward closed	11
3.1.4 G is stable by product	12
3.2 G is generic	12
4 Finite relations and valuations	14
4.1 Bool and Bool'	14
4.2 Totality of mem	15
4.3 Totality of eval	16
4.4 Merging of subH	17
4.5 Monotonicity of test	18
4.6 Monotonicity of eval and subHtree	18
4.7 A proof of upward closure of FVal	20
4.8 A proof of $C[1]$	20
4.9 A proof of $\forall p^k q^k. C[pq] \Rightarrow C[p]$	21
5 Absoluteness of datatypes	21
5.1 Tree is absolute	21
5.2 subH is absolute	25
5.3 FVal is absolute	27
5.4 C is absolute	29
6 Proof of forcing the extra axiom (A)	30
6.1 A textual proof	30
6.2 The formal proof	31
6.3 Simplifying the combinators	33

$$\begin{aligned}
\perp &:= \forall Z^o. Z \\
\neg A &:= A \Rightarrow \perp \\
a =_r b &:= \forall Z^{r \rightarrow o}. Za \Rightarrow Zb \\
A \wedge \dots \wedge B &:= \forall Z^o. (A \Rightarrow \dots \Rightarrow B \Rightarrow Z) \Rightarrow Z \\
A \vee \dots \vee B &:= \forall Z^o. ((A \Rightarrow Z) \Rightarrow \dots \Rightarrow (B \Rightarrow Z) \Rightarrow Z) \Rightarrow Z \\
\exists x^r. A \& \dots \& B := \neg(\forall x^r. A \Rightarrow \dots \Rightarrow B \Rightarrow \perp)
\end{aligned}$$

Figure 1: Encoding of usual connectives into minimal logic

$$\begin{array}{c}
\dfrac{}{\mathcal{E}; \Gamma, x : A \vdash x : A} \quad \dfrac{\mathcal{E}; \Gamma \vdash t : A}{\mathcal{E}; \Gamma \vdash t : A'} \quad A \simeq_{\mathcal{E}} A' \\
\dfrac{\mathcal{E}; \Gamma, x : A \vdash t : B}{\mathcal{E}; \Gamma \vdash \lambda x. t : A \Rightarrow B} \\
\dfrac{\mathcal{E}; \Gamma \vdash t : A \Rightarrow B \quad \mathcal{E}; \Gamma \vdash u : A}{\mathcal{E}; \Gamma \vdash tu : B} \\
\dfrac{\mathcal{E}, M = N; \Gamma \vdash t : A}{\mathcal{E}; \Gamma \vdash t : M \doteq_r N \mapsto A} \quad \dfrac{\mathcal{E}; \Gamma \vdash t : M \doteq_r M \mapsto A}{\mathcal{E}; \Gamma \vdash t : A} \\
\dfrac{\mathcal{E}; \Gamma \vdash t : A}{\mathcal{E}; \Gamma \vdash t : \forall x^r. A} \quad x \notin FV(\Gamma) \quad \dfrac{\mathcal{E}; \Gamma \vdash t : \forall x^r. A}{\mathcal{E}; \Gamma \vdash t : A[N^r/x^r]} \\
\dfrac{}{\mathcal{E}; \Gamma \vdash \text{callcc} : ((A \Rightarrow B) \Rightarrow A) \Rightarrow A}
\end{array}$$

Figure 2: Deduction rules for $\text{PA}\omega^+$

1 Foreword

This appendix is not meant to be read but is rather given as a reference for the definitions and derivations of the proof terms whose existence is claimed in the paper.

The presentation of formal derivation we use is inspired from Fitch-style natural deduction, in the sense that we use a linear presentation and not proof trees. (This is necessary when proof-terms become big).

The bracket on the left side of proofs denote the scope of assumptions that can be of three kinds:

- abstractions, denoted by $[\lambda]$, that introduce proof variables (that thus modify both the term and its type);
- universal quantifications, denoted by $[\forall]$, that do not alter the proof term but only introduce term variables;
- equations, denoted by $[\text{Eq.}]$, that do not effect the proof term but introduce equations for the congruence $\simeq_{\mathcal{E}}$. (The subscript \mathcal{E} will never be written explicitly).

Whenever we write $[Y/Z^*r]$, it stands for the substitution $[(\lambda \vec{x}. Y \vec{x})/Z^*]$. Note that Y and Z will often be the same letter. As in the paper, \equiv stands for syntactic identity, \simeq stands for congruence and $=$ stand for Leibniz equality.

Let us first remind the reader of the definition of the connectives in second-order encodings, the full set of inference rules of the congruence $\simeq_{\mathcal{E}}$ and the deduction rules of system $\text{PA}\omega^+$.

Extending the proof system with new constructions

Notations

In order to ease the presentation of the proof and make it both more understandable and program-like, we will use the following abbreviations:

Reflexivity, symmetry, transitivity and base case

$$\frac{}{M \simeq_{\mathcal{E}} M} \quad \frac{(M = N) \in \mathcal{E}}{M \simeq_{\mathcal{E}} N} \quad \frac{M \simeq_{\mathcal{E}} N}{N \simeq_{\mathcal{E}} M} \quad \frac{M \simeq_{\mathcal{E}} N \quad N \simeq_{\mathcal{E}} P}{M \simeq_{\mathcal{E}} P}$$

Context closure

$$\frac{M \simeq_{\mathcal{E}} N}{\lambda x. M \simeq_{\mathcal{E}} \lambda x. N} \quad \frac{M \simeq_{\mathcal{E}} N \quad P \simeq_{\mathcal{E}} Q}{MP \simeq_{\mathcal{E}} NPQ} \quad \frac{A \simeq_{\mathcal{E}} B \quad C \simeq_{\mathcal{E}} D}{A \Rightarrow C \simeq_{\mathcal{E}} B \Rightarrow D}$$

$$\frac{A \simeq_{\mathcal{E}} B}{\forall x^{\tau}. A \simeq_{\mathcal{E}} \forall x^{\tau}. B} \quad \frac{M \simeq_{\mathcal{E}} N \quad P \simeq_{\mathcal{E}} Q \quad A \simeq_{\mathcal{E}, M=P} B}{M \doteq P \mapsto A \simeq_{\mathcal{E}} N \doteq Q \mapsto B}$$

$\beta\eta\iota$ -conversion

$$\frac{}{(\lambda x. M) N \simeq_{\mathcal{E}} M[N^{\tau}/x^{\tau}]} \quad \frac{}{\lambda x. M x \simeq_{\mathcal{E}} M} \quad x \notin FV(M)$$

$$\frac{\text{rec}_{\tau} MN 0 \simeq_{\mathcal{E}} M}{\text{rec}_{\tau} MN(S P) \simeq_{\mathcal{E}} NP (\text{rec}_{\tau} MN P)}$$

Semantically equivalent propositions

$$\frac{\forall x^{\tau} \forall y^{\sigma}. A \simeq_{\mathcal{E}} \forall y^{\sigma} \forall x^{\tau}. A}{A \Rightarrow \forall x^{\tau}. B \simeq_{\mathcal{E}} \forall x^{\tau}. A \Rightarrow B} \quad \frac{x \notin FV(A)}{\forall x^{\tau}. A \simeq_{\mathcal{E}} A} \quad x \notin FV(A)$$

$$\frac{\forall x^{\tau} M \doteq M \mapsto A \simeq_{\mathcal{E}} A}{M \doteq N \mapsto A \simeq_{\mathcal{E}} N \doteq M \mapsto A} \quad \frac{M \doteq N \mapsto A \simeq_{\mathcal{E}} N \doteq M \mapsto A}{M \doteq N \mapsto P \doteq Q \mapsto A \simeq_{\mathcal{E}} P \doteq Q \mapsto M \doteq N \mapsto A}$$

$$\frac{M \doteq N \mapsto P \doteq Q \mapsto A \simeq_{\mathcal{E}} P \doteq Q \mapsto M \doteq N \mapsto A}{A \Rightarrow M \doteq N \mapsto B \simeq_{\mathcal{E}} M \doteq N \mapsto A \Rightarrow B} \quad \frac{A \Rightarrow M \doteq N \mapsto B \simeq_{\mathcal{E}} M \doteq N \mapsto A \Rightarrow B}{\forall x^{\tau}. M \doteq N \mapsto A \simeq_{\mathcal{E}} M \doteq N \mapsto \forall x^{\tau}. A} \quad x \notin FV(M, N)$$

Figure 3: Inference rules for the relation $\simeq_{\mathcal{E}}$

$\lambda^* x. M$		$\gamma_1(\lambda x. M)$
$M @ N$		$(\gamma_3(M) N)$
$\langle a, b \rangle$		$\lambda f. f a b$
let $(x, y) := c$ in M		$c(\lambda xy. M)$
true		$\lambda xy. x$
false		$\lambda xy. y$
if b then f else g		$b f g$
true*		$\gamma_1(\lambda u. \gamma_1(\lambda v. \beta_3(\beta_4 u)))$
false*	becomes	$\gamma_1(\lambda u. \gamma_1(\lambda v. \beta_4 v))$
Leaf f		$\lambda x_L x_N. x_L f$
Node $a t_1 t_2$		$\lambda x_L x_N. x_N a(t_1 x_L x_N)(t_2 x_L x_N)$
match _{Tree} t with Leaf f => M Node $a t_1 t_2$ => N		$t(\lambda f. M(\lambda a t_1 t_2. N))$
\emptyset		$\lambda x_{\emptyset} x_+ x_- . x_{\emptyset}$
consT $a p$		$\lambda x_{\emptyset} x_+ x_- . x_+ a(p x_{\emptyset} x_+ x_-)$
consF $a p$		$\lambda x_{\emptyset} x_+ x_- . x_- a(p x_{\emptyset} x_+ x_-)$
match _{FVal} c with \emptyset => M consT $a p$ => N consF $a p$ => P		$c M(\lambda a p. N)(\lambda a p. P)$

Admissible derivation rules

Using the definitions of the previous constructions, we can prove the admissibility in $\text{PA}\omega^+$ of the inference rules given figure 4.

$$\begin{array}{c}
\frac{\mathcal{E}; \Gamma, x : q\mathbb{F}A \vdash M : pq\mathbb{F}B}{\mathcal{E}; \Gamma \vdash \lambda^*x. M : p\mathbb{F}A \Rightarrow B} \quad q \notin FV(\Gamma) \\
\frac{\mathcal{E}; \Gamma \vdash M : A \quad \mathcal{E}; \Gamma \vdash N : B}{\mathcal{E}; \Gamma \vdash \langle M, N \rangle : A \wedge B} \quad \frac{\mathcal{E}; \Gamma \vdash M : p\mathbb{F}A \Rightarrow B \quad \mathcal{E}; \Gamma \vdash N : p\mathbb{F}B}{\mathcal{E}; \Gamma \vdash M @ N : p\mathbb{F}B} \\
\frac{\mathcal{E}; \Gamma \vdash \text{true} : 1 \in \text{Bool} \quad \mathcal{E}; \Gamma \vdash \text{false} : 0 \in \text{Bool}}{\mathcal{E}; \Gamma \vdash \text{true}^* : p\mathbb{F}1 \in \text{Bool} \quad \mathcal{E}; \Gamma \vdash \text{false}^* : p\mathbb{F}0 \in \text{Bool}} \\
\frac{\mathcal{E}; \Gamma \vdash M : b \in \text{Bool} \quad \mathcal{E}, b = 1; \Gamma \vdash N : A \quad \mathcal{E}, b = 0; \Gamma \vdash P : A}{\mathcal{E}; \Gamma \vdash \text{if } M \text{ then } N \text{ else } P : A} \\
\frac{\mathcal{E}; \Gamma \vdash M : a \in \text{Atom} \quad \mathcal{E}; \Gamma \vdash N : p \in \text{FVal}}{\mathcal{E}; \Gamma \vdash \text{consF } MN : pa^0 \in \text{FVal}} \quad \frac{\mathcal{E}; \Gamma \vdash M : a \in \text{Atom} \quad \mathcal{E}; \Gamma \vdash N : p \in \text{FVal}}{\mathcal{E}; \Gamma \vdash \text{consT } MN : pa^1 \in \text{FVal}} \text{ memap } \approx_{\mathcal{E}} 0 \\
\frac{\mathcal{E}; \Gamma \vdash \emptyset : \emptyset \in \text{FVal}}{\mathcal{E}; \Gamma \vdash \text{match}_{\text{FVal}} M \text{ with } | \emptyset => N | \text{ consT } xy => P | \text{ consF } xy => Q : A} \quad \frac{\mathcal{E}, p = qa^1; \Gamma \vdash \lambda xy. P : a \in \text{Atom} \Rightarrow q \in \text{FVal} \Rightarrow A \quad \mathcal{E}, p = qa^0; \Gamma \vdash \lambda xy. Q : a \in \text{Atom} \Rightarrow q \in \text{FVal} \Rightarrow A}{\mathcal{E}; \Gamma \vdash p \doteq \emptyset \mapsto N : A}
\end{array}$$

Figure 4: Admissible inference rules for macros

2 Absoluteness of connectives

The derivations for the primitive connectives (false, implication, universal quantification, Leibniz equality) and for the datatype \mathbb{N} can be found in the appendix of [13].

2.1 Conjunction

We deal only with binary conjunctions but the extension to bigger arities is straightforward. Assuming that both propositions A and B are absolute, we want to show that $A \wedge B$ is absolute.

$$\begin{aligned}
& [[\lambda] z : p \models A \wedge B \simeq \forall(Z^*)^{\kappa \rightarrow o}. p \models (A \Rightarrow B \Rightarrow Z) \Rightarrow Z \\
& [\lambda] c : C[p] \\
& [[\forall] Z^o \\
& [\lambda] x : A \Rightarrow B \Rightarrow Z \\
& z : (p \models A \Rightarrow B \Rightarrow Z) \Rightarrow Z [Z/Z^* r] \\
& \gamma_3 z : (p \models A \Rightarrow B \Rightarrow Z) [Z/Z^* r] \Rightarrow (p \models Z) [Z/Z^* r] \\
& [[\forall] q_A^\kappa \\
& [\lambda] u : q_A \models A \\
& \xi_A u : C[q_A] \Rightarrow A \\
& [[\forall] q_B^\kappa \\
& [\lambda] v : q_B \models B \\
& \xi_B v : C[q_B] \Rightarrow B \\
& [[\forall] r^\kappa \\
& [\lambda] c : C[pq_A q_B r] \\
& \alpha_2(\alpha_1^2 c) : C[q_A] \\
& \xi_A u(\alpha_2(\alpha_1^2 c)) : A \\
& x(\xi_A u(\alpha_2(\alpha_1^2 c))) : B \Rightarrow Z \\
& \alpha_2(\alpha_1 c) : C[q_B] \\
& \xi_B v(\alpha_2(\alpha_1 c)) : B \\
& x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c))) : Z \\
& \lambda c. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c))) : \forall r^\kappa. C[pq_A q_B r] \Rightarrow Z \equiv (pq_A q_B \models Z) [Z/Z^* r] \\
& \lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c))) : \forall q_B^\kappa. (q_B \models B) \Rightarrow (pq_A q_B \models Z) [Z/Z^* r] \\
& \gamma_1(\lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c)))) : (pq_A \models B \Rightarrow Z) [Z/Z^* r] \\
& \lambda u. \gamma_1(\lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c)))) : \forall q_A^\kappa. (q_A \models A) \Rightarrow (pq_A \models B \Rightarrow Z) [Z/Z^* r] \\
& \gamma_1(\lambda u. \gamma_1(\lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c)))) : (p \models A \Rightarrow B \Rightarrow Z) [Z/Z^* r] \\
& \gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c)))))) : (p \models Z) [Z/Z^* r] \equiv \forall r^\kappa. C[pr] \Rightarrow Z \\
& \alpha_7 c : C[p1] \\
& \gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c)))))(\alpha_7 c) : Z \\
& \lambda x. \gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c)))))(\alpha_7 c) : \forall Z^o. (A \Rightarrow B \Rightarrow Z) \Rightarrow Z \equiv A \wedge B \\
& \lambda z cx. \gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda vc. x(\xi_A u(\alpha_2(\alpha_1^2 c))) (\xi_B v(\alpha_2(\alpha_1 c)))))(\alpha_7 c) : (p \models A \wedge B) \Rightarrow (C[p] \Rightarrow A \wedge B)
\end{aligned}$$

$$\begin{aligned}
& [\lambda] z : C[p] \Rightarrow A \wedge B \\
& [\forall] (Z^*)^{\kappa \rightarrow o} \\
& [\forall] q^\kappa \\
& [\lambda] x : q \Vdash A \Rightarrow B \Rightarrow Z \\
& \gamma_3 x : (q \Vdash A) \Rightarrow (q \Vdash B \Rightarrow Z) \\
& [\forall] r^\kappa \\
& [\lambda] c : C[pqr] \\
& \alpha_1^2 c : C[p] \\
& z(\alpha_1^2 c) : A \wedge B \\
& z(\alpha_1^2 c) : (A \Rightarrow B \Rightarrow Z^* r) \Rightarrow Z^* r \\
& [\lambda] u : A \\
& [\lambda] v : B \\
& \lambda_{_} u : C[q] \Rightarrow A \\
& \xi'_A (\lambda_{_} u) : q \Vdash A \\
& \gamma_3 x (\xi'_A (\lambda_{_} u)) : q \Vdash B \Rightarrow Z \\
& \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) : (q \Vdash B) \Rightarrow (q \Vdash Z) \\
& \lambda_{_} v : C[q] \Rightarrow B \\
& \xi'_B (\lambda_{_} v) : q \Vdash B \\
& \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) : q \Vdash Z \equiv \forall r^\kappa. C[qr] \Rightarrow Z^* r \\
& \alpha_{10} c : C[qr] \\
& \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c) : Z^* r \\
& \lambda u v. \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c) : A \Rightarrow B \Rightarrow Z^* r \\
& z(\alpha_1^2 c) (\lambda u v. \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c)) : Z^* r \\
& \lambda c. z(\alpha_1^2 c) (\lambda u v. \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c)) : \forall r^\kappa. C[pqr] \Rightarrow Z^* r \equiv pq \Vdash Z \\
& \lambda x c. z(\alpha_1^2 c) (\lambda u v. \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c)) : \forall q^\kappa. (q \Vdash A \Rightarrow B \Rightarrow Z) \Rightarrow (pq \Vdash Z) \\
& \gamma_1 (\lambda x c. z(\alpha_1^2 c) (\lambda u v. \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c))) : p \Vdash (A \Rightarrow B \Rightarrow Z) \Rightarrow Z \\
& \gamma_1 (\lambda x c. z(\alpha_1^2 c) (\lambda u v. \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c))) : \forall Z^*. p \Vdash (A \Rightarrow B \Rightarrow Z) \Rightarrow Z \simeq p \Vdash A \wedge B \\
& \lambda z. \gamma_1 (\lambda x c. z(\alpha_1^2 c) (\lambda u v. \gamma_3 (\gamma_3 x (\xi'_A (\lambda_{_} u))) (\xi'_B (\lambda_{_} v)) (\alpha_{10} c))) : (C[p] \Rightarrow A \wedge B) \Rightarrow (p \Vdash A \wedge B)
\end{aligned}$$

Abstracting over the proof terms ξ_A , ξ'_A , ξ_B and ξ'_B , we get two combinators ξ_\wedge and ξ'_\wedge which can be used to prove absoluteness of more complex propositions.

2.2 Disjunction

Again, we deal only with the binary case but the extension to bigger arities is straightforward.

$$\begin{aligned}
& [\lambda] z : p \Vdash A \vee B \\
& [\lambda] c : C[p] \\
& \boxed{[\forall] Z^o} \\
& [\lambda] x : A \Rightarrow Z \\
& [\lambda] y : B \Rightarrow Z \\
& z : (p \Vdash (A \Rightarrow Z) \Rightarrow (B \Rightarrow Z) \Rightarrow Z)[Z/Z^* r] \\
& \gamma_3 z : (p \Vdash A \Rightarrow Z)[Z/Z^* r] \Rightarrow (p \Vdash (B \Rightarrow Z) \Rightarrow Z)[Z/Z^* r] \\
& \boxed{[\forall] q^\kappa} \\
& [\lambda] u : q \Vdash A \\
& \xi_A u : C[q] \Rightarrow A \\
& \boxed{[\forall] r^\kappa} \\
& [\lambda] c' : C[pqr] \\
& \alpha_1 c' : C[pq] \\
& \alpha_2(\alpha_1 c') : C[q] \\
& \xi_A u(\alpha_2(\alpha_1 c')) : A \\
& [x(\xi_A u(\alpha_2(\alpha_1 c')))] : Z \\
& \lambda c'. x(\xi_A u(\alpha_2(\alpha_1 c'))) : \forall r^\kappa. C[pqr] \Rightarrow Z \equiv (pq \Vdash Z)[Z/Z^* r] \\
& \lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c'))) : \forall q^\kappa. (q \Vdash A) \Rightarrow (pq \Vdash Z)[Z/Z^* r] \\
& \gamma_1(\lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c')))) : (p \Vdash A \Rightarrow Z)[Z/Z^* r] \\
& \text{idem} \\
& \gamma_1(\lambda u c'. x(\xi_B u(\alpha_2(\alpha_1 c')))) : (p \Vdash B \Rightarrow Z)[Z/Z^* r] \\
& \gamma_3 z(\gamma_1(\lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c'))))) : (p \Vdash (B \Rightarrow Z) \Rightarrow Z)[Z/Z^* r] \\
& \gamma_3(\gamma_3 z(\gamma_1(\lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c')))))) : (p \Vdash B \Rightarrow Z)[Z/Z^* r] \Rightarrow (p \Vdash Z)[Z/Z^* r] \\
& \gamma_3(\gamma_3 z(\gamma_1(\lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c'))))))(\gamma_1(\lambda u c'. x(\xi_B u(\alpha_2(\alpha_1 c'))))) : (p \Vdash Z)[Z/Z^* r] \equiv \forall r^\kappa. C[pr] \Rightarrow Z \\
& \alpha_7 c : C[p1] \\
& \gamma_3(\gamma_3 z(\gamma_1(\lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c'))))))(\gamma_1(\lambda u c'. x(\xi_B u(\alpha_2(\alpha_1 c')))))(\alpha_7 c) : Z \\
& \lambda x y. \gamma_3 z(\gamma_3 z(\gamma_1(\lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c'))))))(\gamma_1(\lambda u c'. x(\xi_B u(\alpha_2(\alpha_1 c')))))(\alpha_7 c) : \forall Z^o. (A \Rightarrow Z) \Rightarrow (B \Rightarrow Z) \Rightarrow Z \equiv A \vee B \\
& \lambda z c x y. \gamma_3 z(\gamma_3 z(\gamma_1(\lambda u c'. x(\xi_A u(\alpha_2(\alpha_1 c'))))))(\gamma_1(\lambda u c'. x(\xi_B u(\alpha_2(\alpha_1 c')))))(\alpha_7 c) : (p \Vdash A \vee B) \Rightarrow (C[p] \Rightarrow A \vee B)
\end{aligned}$$

$$\begin{aligned}
& [\lambda] z : C[p] \Rightarrow A \vee B \\
& [\forall] (Z^*)^{\kappa \rightarrow o} \\
& [\forall] q_A^\kappa \\
& [\lambda] x : q_A \Vdash A \Rightarrow Z \\
& \beta_3 x : (q_A q_B \Vdash A \Rightarrow Z) \\
& \gamma_3 (\beta_3 x) : (q_A q_B \Vdash A) \Rightarrow (q_A q_B \Vdash Z) \\
& [\forall] q_B^\kappa \\
& [\lambda] y : q_B \Vdash B \Rightarrow Z \\
& \beta_4 y : (q_A q_B \Vdash B \Rightarrow Z) \\
& \gamma_3 (\beta_4 y) : (q_A q_B \Vdash B) \Rightarrow (q_A q_B \Vdash Z) \\
& [\forall] r^\kappa \\
& [\lambda] c : C[pq_A q_B r] \\
& \alpha_1^3 c : C[p] \\
& z(\alpha_1^3 c) : A \vee B \\
& z(\alpha_1^3 c) : (A \Rightarrow (q_A q_B \Vdash Z)) \Rightarrow (B \Rightarrow (q_A q_B \Vdash Z)) \Rightarrow (q_A q_B \Vdash Z) \\
& [\lambda] u : A \\
& \lambda_{_} u : C[q_A q_B] \Rightarrow A \\
& \xi'_A (\lambda_{_} u) : q_A q_B \Vdash A \\
& \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u)) : q_A q_B \Vdash Z \\
& \lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u)) : A \Rightarrow (q_A q_B \Vdash Z) \\
& z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) : (B \Rightarrow (q_A q_B \Vdash Z)) \Rightarrow (q_A q_B \Vdash Z) \\
& [\lambda] v : B \\
& \lambda_{_} v : C[q_A q_B] \Rightarrow B \\
& \xi'_B (\lambda_{_} u) : q_A q_B \Vdash B \\
& \gamma_3 (\beta_4 x) (\xi'_B (\lambda_{_} v)) : q_A q_B \Vdash Z \\
& \lambda v. \gamma_3 (\beta_4 x) (\xi'_B (\lambda_{_} v)) : B \Rightarrow (q_A q_B \Vdash Z) \\
& z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) : q_A q_B \Vdash Z \equiv \forall r^\kappa. C[q_A q_B r] \Rightarrow Z^* r \\
& \alpha_{16} c : C[q_A q_B r] \\
& z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c) : Z^* r \\
& \lambda c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c) : \forall r^\kappa. C[pq_A q_B r] \Rightarrow Z^* r \equiv pq_A q_B \Vdash Z \\
& \lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c) : \forall q_B^\kappa. (q_B \Vdash B \Rightarrow Z) \Rightarrow (pq_A q_B \Vdash Z) \\
& \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c)) : pq_A \Vdash (B \Rightarrow Z) \Rightarrow Z \\
& \lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c)) : \forall q_A^\kappa (q_A \Vdash A \Rightarrow Z) \Rightarrow (pq_A \Vdash (B \Rightarrow Z) \Rightarrow Z) \\
& \gamma_1 (\lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c))) : p \Vdash (A \Rightarrow Z) \Rightarrow (B \Rightarrow Z) \Rightarrow Z \\
& \gamma_1 (\lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c))) \\
& \quad : \forall Z^*. p \Vdash (A \Rightarrow Z) \Rightarrow (B \Rightarrow Z) \Rightarrow Z \simeq p \Vdash A \vee B \\
& \lambda z. \gamma_1 (\lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_A (\lambda_{_} u))) (\lambda v. \gamma_3 (\beta_4 y) (\xi'_B (\lambda_{_} v))) (\alpha_{16} c))) : (C[p] \Rightarrow A \vee B) \Rightarrow (p \Vdash A \vee B)
\end{aligned}$$

Abstracting over the proof terms ξ_A , ξ'_A , ξ_B and ξ'_B , we get two combinators ξ_V and ξ'_V which can be used to prove absoluteness of more complex propositions.

2.3 Classical existential quantifier (on multiple propositions)

Remember that $\exists x^\tau. A_1 \& \dots \& A_k$ is defined as $\neg(\forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp)$.

Here, we assume that τ is a T -sort, that is we have $\tau^* = \tau$. Otherwise, the result does not hold.

$$\begin{aligned}
& [\lambda] z : p \Vdash \exists x^\tau. A_1 \& \dots \& A_k \equiv p \Vdash (\forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp) \Rightarrow \perp \\
& [\lambda] c : C[p] \\
& [\lambda] x : \forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp \\
& \gamma_3 z : (p \Vdash \forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp) \Rightarrow (p \Vdash \perp) \\
& [\forall] x^\tau \\
& \quad [\forall] q_1^k \\
& \quad [\lambda] x_1 : q_1 \Vdash A_1 \\
& \quad \xi_1 x_1 : C[q_1] \Rightarrow A_1 \\
& \quad \vdots \\
& \quad [\forall] q_k^k \\
& \quad [\lambda] x_k : q_k \Vdash A_k \\
& \quad \xi_k x_k : C[q_k] \Rightarrow A_k \\
& \quad [\lambda] c' : C[pq_1 \dots q_k] \\
& \quad \alpha_2(\alpha_1^{k-1} c') : C[q_1] \\
& \quad \xi_1 x_1(\alpha_2(\alpha_1^{k-1} c')) : A_1 \\
& \quad \vdots \\
& \quad \alpha_2 c' : C[q_k] \\
& \quad \xi_k x_k(\alpha_2 c') : A_k \\
& \quad x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2 c')) : \perp \\
& \quad \lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c'))) : C[pq_1 \dots q_k] \Rightarrow \perp \\
& \quad \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))) : pq_1 \dots q_k \Vdash \perp \\
& \quad \lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))) : \forall q_k^k (q_k \Vdash A_k) \Rightarrow (pq_1 \dots q_k \Vdash \perp) \\
& \quad \gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c'))))) : pq_1 \dots q_{k-1} \Vdash A_k \Rightarrow \perp \\
& \quad \vdots \\
& \quad \lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots) : \forall q_1^k (q_1 \Vdash A_1) \Rightarrow (pq_1 \Vdash A_2 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp) \\
& \quad \gamma_1(\lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots) : p \Vdash A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp \\
& \quad \gamma_1(\lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots) \\
& \quad \quad : \forall x^\tau. p \Vdash A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp \simeq p \Vdash \forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp \\
& \quad \gamma_3 z(\gamma_1(\lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots)) : p \Vdash \perp \\
& \quad \xi_\perp(\gamma_3 z(\gamma_1(\lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots)))) : C[p] \Rightarrow \perp \\
& \quad \xi_\perp(\gamma_3 z(\gamma_1(\lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots)))) c : \perp \\
& \quad \lambda z c x. \xi_\perp(\gamma_3 z(\gamma_1(\lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots)))) c \\
& \quad : (p \Vdash \exists x^\tau. A_1 \& \dots \& A_k) \Rightarrow C[p] \Rightarrow (\forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp) \Rightarrow \perp \equiv (p \Vdash \exists x^\tau. A_1 \& \dots \& A_k) \Rightarrow (C[p] \Rightarrow \exists x^\tau. A_1 \& \dots \& A_k)
\end{aligned}$$

Abstracting over the k proof term ξ_1, \dots, ξ_k , we get the proof term

$$\xi_{\exists_k} := \lambda \xi_1 \dots \xi_k \lambda z c x. \xi_\perp(\gamma_3 z(\gamma_1(\lambda x_1. \gamma_1 \dots (\gamma_1(\lambda x_k. \xi'_\perp(\lambda c'. x(\xi_1 x_1(\alpha_2(\alpha_1^{k-1} c'))) \dots (\xi_k x_k(\alpha_2(\alpha_1 c')))))) \dots)))) c$$

which can be used as a combinator to prove the absoluteness of the proposition $\exists x^\tau. A_1 \& \dots \& A_k$ from the proof terms ξ_1, \dots, ξ_k of absoluteness of A_1, \dots, A_k .

$[\lambda] z : C[p] \Rightarrow \exists x^\tau. A_1 \& \dots \& A_k$
$[\forall] q^k$
$[\lambda] x : q \Vdash \forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp \simeq \forall x^\tau. q \Vdash A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp$
$[\lambda] c : C[pq]$
$\alpha_1 c : C[p]$
$z(\alpha_1 c) : \exists x^\tau. A_1 \& \dots \& A_k \equiv (\forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp) \Rightarrow \perp$
$[\forall] x^\tau$
$[\lambda] y_1 : A_1$
\vdots
$[\lambda] y_k : A_k$
$\lambda_{_} y_1 : C[pq] \Rightarrow A_1$
$\xi'_1(\lambda_{_} y_1) : pq \Vdash A_1$
$\beta_4 x : \forall x^\tau. pq \Vdash A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp$
$\gamma_3(\beta_4 x) : \forall x^\tau. (pq \Vdash A_1) \Rightarrow (pq \Vdash A_2 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp)$
$\gamma_3 x(\xi'_1(\lambda_{_} y_1)) : pq \Vdash A_2 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp$
\vdots
$\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots) : pq \Vdash A_k \Rightarrow \perp$
$\lambda_{_} y_k : C[pq] \Rightarrow A_k$
$\xi'_k(\lambda_{_} y_k) : pq \Vdash A_k$
$\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots)) : (pq \Vdash A_k) \Rightarrow (pq \Vdash \perp)$
$\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)) : pq \Vdash \perp$
$\xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k))) : C[pq] \Rightarrow \perp$
$\xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c : \perp$
$\lambda y_1 \dots y_k. \xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c : \forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp$
$\lambda z(\alpha_1 c)(\lambda y_1 \dots y_k. \xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c) : \perp$
$\lambda c. \lambda z(\alpha_1 c)(\lambda y_1 \dots y_k. \xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c) : C[pq] \Rightarrow \perp$
$\lambda \xi'_\perp(\lambda c. \lambda z(\alpha_1 c)(\lambda y_1 \dots y_k. \xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c)) : pq \Vdash \perp$
$\lambda x. \xi'_\perp(\lambda c. \lambda z(\alpha_1 c)(\lambda y_1 \dots y_k. \xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c))$
$: \forall q^k. (q \Vdash \forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp) \Rightarrow (pq \Vdash \perp)$
$\gamma_1(\lambda x. \xi'_\perp(\lambda c. \lambda z(\alpha_1 c)(\lambda y_1 \dots y_k. \xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c)))$
$: p \Vdash (\forall x^\tau. A_1 \Rightarrow \dots \Rightarrow A_k \Rightarrow \perp) \Rightarrow \perp \equiv p \Vdash \exists x^\tau. A_1 \& \dots \& A_k$
$\lambda z. \gamma_1(\lambda x. \xi'_\perp(\lambda c. \lambda z(\alpha_1 c)(\lambda y_1 \dots y_k. \xi_\perp(\gamma_3(\gamma_3(\dots(\gamma_3(\beta_4 x)(\xi'_1(\lambda_{_} y_1)))\dots))(\xi'_k(\lambda_{_} y_k)))c)))$
$: (C[p] \Rightarrow \exists x^\tau. A_1 \& \dots \& A_k) \Rightarrow (p \Vdash \exists x^\tau. A_1 \& \dots \& A_k)$

Again, we build a combinator ξ'_{\exists_k} to use in bigger propositions.

3 Properties of the generic set

We recall that by definition, $(p \in G)^* q := C[pq]$ so that $q \Vdash p \in G \equiv \forall r^\kappa. C[qr] \Rightarrow C[pr] \equiv q \leq p$.

3.1 G is a filter

Saying that G is a filter amounts to proving four properties:

- G is non empty: $1 \in G$
- G is a subset of $\forall C$:
- G is upward-closed:
- G is closed under product:

3.1.1 G is non-empty

We want to prove $1 \Vdash 1 \in G \equiv 1 \leq 1 \equiv \forall r^\kappa. C[1r] \Rightarrow C[1r]$.

$$\begin{array}{l} [\forall] r^\kappa \\ [\lambda] x : C[1r] \\ x : C[1r] \\ \lambda x. x : \forall r^\kappa. C[1r] \Rightarrow C[1r] \end{array}$$

3.1.2 G is a subset of C

We want to prove that for all p^κ , $1 \Vdash p \in G \Rightarrow C[p]$.

$$\begin{array}{l} [\forall] q^\kappa \\ [\lambda] x : q \Vdash p \in G \equiv q \leq p \equiv \forall r^\kappa. C[qr] \Rightarrow C[pr] \\ [\forall] r^\kappa \\ [\lambda] c : C[1q] \\ \alpha_3 c : C[q1] \\ x(\alpha_3 c) : C[p1] \\ \alpha_1(x(\alpha_3 c)) : C[p] \\ \alpha_1 \circ x \circ \alpha_3 \equiv \lambda c. \alpha_1(x(\alpha_3 c)) : C[1q] \Rightarrow C[p] \\ \xi'_C(\alpha_1 \circ x \circ \alpha_3) : 1q \Vdash C[p] \\ \lambda x. \xi'_C(\alpha_1 \circ x \circ \alpha_3) : \forall q^\kappa. (q \Vdash p \in G) \Rightarrow (1q \Vdash C[p]) \\ \gamma_1(\lambda x. \xi'_C(\alpha_1 \circ x \circ \alpha_3)) : 1 \Vdash p \in G \Rightarrow C[p] \end{array}$$

3.1.3 G is upward closed

We want to prove that for all p^κ and q^κ , $1 \Vdash pq \in G \Rightarrow p \in G$.

$$\begin{array}{l} [\forall] r^\kappa \\ [\lambda] x : r \Vdash pq \in G \equiv r \leq pq \equiv \forall s^\kappa. C[rs] \Rightarrow C[pqs] \\ [\forall] s^\kappa \\ [\lambda] c : C[1rs] \\ \alpha_{10} c : C[rs] \\ x(\alpha_{10} c) : C[pqs] \\ \alpha_9(x(\alpha_{10} c)) : C[ps] \\ \alpha_9 \circ x \circ \alpha_{10} \equiv \lambda c. \alpha_9(x(\alpha_{10} c)) : \forall s^\kappa. C[1rs] \Rightarrow C[ps] \equiv 1r \leq p \equiv 1r \Vdash p \in G \\ \lambda x. \alpha_9 \circ x \circ \alpha_{10} : \forall r^\kappa. r \Vdash pq \in G \Rightarrow 1r \Vdash p \in G \\ \gamma_1(\lambda x. \alpha_9 \circ x \circ \alpha_{10}) : 1 \Vdash pq \in G \Rightarrow p \in G \end{array}$$

3.1.4 G is stable by product

We want to prove that for all p^κ and q^κ , $1 \Vdash p \in G \Rightarrow q \in G \Rightarrow pq \in G$.

$$\begin{aligned}
& [\forall] r_p^\kappa \\
& [\lambda] x : r_p \Vdash p \in G \equiv \forall s^\kappa. C[r_p s] \Rightarrow C[ps] \\
& [\forall] r_q^\kappa \\
& [\lambda] y : r_q \Vdash q \in G \equiv \forall s^\kappa. C[r_q s] \Rightarrow C[qs] \\
& [\forall] s^\kappa \\
& [\lambda] c : C[1r_p r_q s] \\
& \alpha_5 c : C[1r_p(r_q s)] \\
& \alpha_5^2 c : C[1(r_p(r_q s))] \\
& \alpha_2(\alpha_5^2 c) : C[r_p(r_q s)] \\
& x(\alpha_2(\alpha_5^2 c)) : C[p(r_q s)] \\
& \alpha_{12}(x(\alpha_2(\alpha_5^2 c))) : C[r_q(sp)] \\
& y(\alpha_{12}(x(\alpha_2(\alpha_5^2 c)))) : C[q(sp)] \\
& \alpha_{13}(y(\alpha_{12}(x(\alpha_2(\alpha_5^2 c))))) : C[pqs] \\
& \alpha_{13} \circ y \circ \alpha_{12} \circ x \circ \alpha_2 \circ \alpha_5^2 \equiv \lambda c. \alpha_{13}(y(\alpha_{12}(x(\alpha_2(\alpha_5^2 c))))) : \forall s^\kappa. C[1r_p r_q s] \Rightarrow C[pqs] \equiv 1r_p r_q \Vdash pq \in G \\
& \lambda y. \alpha_{13} \circ y \circ \alpha_{12} \circ x \circ \alpha_2 \circ \alpha_5^2 : \forall r_q^\kappa. r_q \Vdash q \in G \Rightarrow 1r_p r_q \Vdash pq \in G \\
& \gamma_1(\lambda y. \alpha_{13} \circ y \circ \alpha_{12} \circ x \circ \alpha_2 \circ \alpha_5^2) : 1r_p \Vdash q \in G \Rightarrow pq \in G \\
& \lambda x. \gamma_1(\lambda y. \alpha_{13} \circ y \circ \alpha_{12} \circ x \circ \alpha_2 \circ \alpha_5^2) : \forall r_p^\kappa. r_p \Vdash p \in G \Rightarrow 1r_p \Vdash q \in G \Rightarrow pq \in G \\
& \gamma_1(\lambda x. \gamma_1(\lambda y. \alpha_{13} \circ y \circ \alpha_{12} \circ x \circ \alpha_2 \circ \alpha_5^2)) : 1 \Vdash p \in G \Rightarrow q \in G \Rightarrow pq \in G
\end{aligned}$$

3.2 G is generic

We want to prove for all absolute set $D^{\kappa \rightarrow o}$:

$$1 \Vdash (\forall p^\kappa. C[p] \Rightarrow \exists q^\kappa. C[pq] \& pq \in D) \Rightarrow \exists p^\kappa. p \in G \& p \in D.$$

Recall that with the proof terms of absoluteness of existential quantification (with two formulas), we can get proof terms for the absoluteness of the formula $\exists x. A \& B$:

- for all x and p , $\xi_{\exists_2} \xi_A \xi_B : (p \Vdash \exists x. A \& B) \Rightarrow (C[p] \Rightarrow \exists x. A \& B)$
- for all x and p , $\xi'_{\exists_2} \xi'_A \xi'_B : (C[p] \Rightarrow \exists x. A \& B) \Rightarrow (p \Vdash \exists x. A \& B)$

Because D is absolute, we have for all p^κ and r^κ :

$$\begin{aligned}
\xi_D & : (r \Vdash p \in D) \Rightarrow (C[r] \Rightarrow p \in D) \\
\xi'_D & : (C[r] \Rightarrow p \in D) \Rightarrow (r \Vdash p \in D)
\end{aligned}$$

We will use the fact that if $D^{\kappa \rightarrow o}$ is absolute, then for all $p^\kappa, q \mapsto pq \in D$ is also absolute and the proof terms are the same. This comes from the fact that the deduction rules for \forall are computationally transparent.

$\boxed{[\forall] r_1^\kappa}$	
$[\lambda] x : r_1 \models \forall p^\kappa. C[p] \Rightarrow \exists q^\kappa. C[pq] \& pq \in D$	
$\boxed{[\forall] r_2^\kappa}$	
$[\lambda] y : r_2 \models \forall p^\kappa. p \in G \Rightarrow p \in D \Rightarrow \perp \simeq \forall p^\kappa. r_2 \models p \in G \Rightarrow p \in D \Rightarrow \perp$	
$[\lambda] c : C[1r_1r_2]$	
$x : r_1 \models C[1r_1r_2] \Rightarrow \exists q^\kappa. C[1r_1r_2q] \& 1r_1r_2q \in D$	
$\gamma_3 x : (r_1 \models C[1r_1r_2]) \Rightarrow (r_1 \models \exists q^\kappa. C[1r_1r_2q] \& 1r_1r_2q \in D)$	
$\lambda_{_}. c : C[r_1] \Rightarrow C[1r_1r_2]$	
$\xi'_C (\lambda_{_}. c) : r_1 \models C[1r_1r_2]$	
$\gamma_3 x (\xi'_C (\lambda_{_}. c)) : r_1 \models \exists q^\kappa. C[1r_1r_2q] \& 1r_1r_2q \in D$	
$\xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) : C[r_1] \Rightarrow \exists q^\kappa. C[1r_1r_2q] \& 1r_1r_2q \in D$	
$\alpha_1 c : C[1r_1]$	
$\alpha_2 (\alpha_1 c) : C[r_1]$	
$\xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) : \exists q^\kappa. C[1r_1r_2q] \& 1r_1r_2q \in D \equiv \neg(\forall q^\kappa. C[1r_1r_2q] \Rightarrow 1r_1r_2q \in D \Rightarrow \perp)$	
$\boxed{[\forall] q^\kappa}$	
$[\lambda] c' : C[1r_1r_2q]$	
$[\lambda] d : 1r_1r_2q \in D$	
$y : r_2 \models 1r_1r_2q \in G \Rightarrow 1r_1r_2q \in D \Rightarrow \perp$	
$\beta_4 y : 1r_1r_2 \models 1r_1r_2q \in G \Rightarrow 1r_1r_2q \in D \Rightarrow \perp$	
$\beta_3 (\beta_4 y) : 1r_1r_2q \models 1r_1r_2q \in G \Rightarrow 1r_1r_2q \in D \Rightarrow \perp$	
$\gamma_3 (\beta_3 (\beta_4 y)) : (1r_1r_2q \models 1r_1r_2q \in G) \Rightarrow (1r_1r_2q \models 1r_1r_2q \in D \Rightarrow \perp)$	
$\lambda x. x : 1r_1r_2q \leq 1r_1r_2q \equiv 1r_1r_2q \models 1r_1r_2q \in G$	
$\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x) : 1r_1r_2q \models 1r_1r_2q \in D \Rightarrow \perp$	
$\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) : (1r_1r_2q \models 1r_1r_2q \in D) \Rightarrow (1r_1r_2q \models \perp)$	
$\lambda_{_}. d : C[1r_1r_2q] \Rightarrow 1r_1r_2q \in D$	
$\xi'_D (\lambda_{_}. d) : 1r_1r_2q \models 1r_1r_2q \in D$	
$\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d)) : 1r_1r_2q \models \perp$	
$\xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) : C[1r_1r_2q] \Rightarrow \perp$	
$\xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c' : \perp$	
$\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c' : \forall q^\kappa. C[1r_1r_2q] \Rightarrow 1r_1r_2q \in D \Rightarrow \perp$	
$\xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) (\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c')) : \perp$	
$\lambda c. \xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) (\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c')) : C[1r_1r_2] \Rightarrow \perp$	
$\xi'_\perp (\lambda c. \xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) (\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c'))) : 1r_1r_2 \models \perp$	
$\lambda y. \xi'_\perp (\lambda c. \xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) (\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c'))) : 1r_1r_2 \models \perp$	
$: \forall r_2^\kappa. (r_2 \models \forall p^\kappa. p \in G \Rightarrow p \in D \Rightarrow \perp) \Rightarrow (1r_1r_2 \models \perp)$	
$\gamma_1 (\lambda y. \xi'_\perp (\lambda c. \xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) (\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c'))))$	
$: 1r_1 \models (\forall p^\kappa. p \in G \Rightarrow p \in D \Rightarrow \perp) \Rightarrow \perp \equiv 1r_1 \models \exists p^\kappa. p \in G \& p \in G$	
$\lambda x. \gamma_1 (\lambda y. \xi'_\perp (\lambda c. \xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) (\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c'))))$	
$: \forall r_1^\kappa. (r_1 \models \forall p^\kappa. C[p] \Rightarrow \exists q^\kappa. C[pq] \& pq \in D) \Rightarrow (1r_1 \models \exists p^\kappa. p \in G \& p \in G)$	
$\gamma_1 (\lambda x. \gamma_1 (\lambda y. \xi'_\perp (\lambda c. \xi_{\exists_2} \xi_C \xi_D (\gamma_3 x (\xi'_C (\lambda_{_}. c))) (\alpha_2 (\alpha_1 c)) (\lambda c' d. \xi_\perp (\gamma_3 (\gamma_3 (\beta_3 (\beta_4 y)) (\lambda x. x)) (\xi'_D (\lambda_{_}. d))) c')))))$	
$: 1 \models (\forall p^\kappa. C[p] \Rightarrow \exists q^\kappa. C[pq] \& pq \in D) \Rightarrow \exists p^\kappa. p \in G \& p \in G$	

4 Finite relations and valuations

4.1 Bool and Bool'

Let $b \in \text{Bool}' := \forall Z^o. (b \doteq 1 \mapsto Z) \Rightarrow (b \doteq 0 \mapsto Z) \Rightarrow Z$.

We show the equivalence with $b \in \text{Bool} := \forall Z^{t \rightarrow o}. Z 1 \Rightarrow Z 0 \Rightarrow Z b$.

Since the denotation in classical realizability are the same, realizers are also the same and from a computational point of view, we can consider Bool and Bool' to be equal. Nevertheless, they are not exactly the same from the point of view of proofs.

$$\begin{array}{l} \boxed{[\lambda] z : b \in \text{Bool}' \\ \quad \boxed{[\forall] Z^{t \rightarrow o} \\ \quad \boxed{[\lambda] x : Z 1 \\ \quad \boxed{[\lambda] y : Z 0 \\ \quad z : (b \doteq 1 \mapsto Z b) \Rightarrow (b \doteq 0 \mapsto Z b) \Rightarrow Z b \\ \quad x : b \doteq 1 \mapsto Z 1 \simeq b \doteq 1 \mapsto Z b \\ \quad y : b \doteq 0 \mapsto Z 0 \simeq b \doteq 0 \mapsto Z b \\ \quad zxy : Z b \\ \quad \boxed{\lambda xy. zxy : \forall Z^{t \rightarrow o}. Z 1 \Rightarrow Z 0 \Rightarrow Z b \equiv b \in \text{Bool}} \\ \quad \boxed{\lambda zxy. zxy : b \in \text{Bool}' \Rightarrow b \in \text{Bool}}} } } } } \end{array}$$

If we had subtyping in $\text{PA}\omega^+$, the conversion would be the identity and not an η -expansion thereof.

$$\begin{array}{l} \boxed{[\lambda] z : b \in \text{Bool} \\ \quad \boxed{[\forall] Z^o \\ \quad \boxed{\text{We set } Z x := x \doteq b \mapsto Z \text{ in } z \\ \quad z : (1 \doteq b \mapsto Z) \Rightarrow (0 \doteq b \mapsto Z) \Rightarrow (b \doteq b \mapsto Z) \simeq (b \doteq 1 \mapsto Z) \Rightarrow (b \doteq 0 \mapsto Z) \Rightarrow Z \\ \quad z : b \in \text{Bool}' \\ \quad \lambda z. z : b \in \text{Bool} \Rightarrow b \in \text{Bool}'}}} } } \end{array}$$

Yet, it is enough to have the deduction rule we want for the if – then – else construction:

$$\frac{\mathcal{E}; \Gamma \vdash M : b \in \text{Bool}}{\mathcal{E}; \Gamma \vdash M : (b \doteq 1 \mapsto A) \Rightarrow (b \doteq 0 \mapsto A) \Rightarrow (b \doteq b \mapsto A)} \quad \frac{\mathcal{E}, b = 1; \Gamma \vdash N : A}{\mathcal{E}; \Gamma \vdash N : b \doteq 1 \mapsto A} \quad \frac{\mathcal{E}, b = 0; \Gamma \vdash P : A}{\mathcal{E}; \Gamma \vdash P : b \doteq 0 \mapsto A}$$

$$\frac{\mathcal{E}; \Gamma \vdash M : (b \doteq 1 \mapsto A) \Rightarrow (b \doteq 0 \mapsto A) \Rightarrow A \quad \mathcal{E}; \Gamma \vdash MN : (b \doteq 0 \mapsto A) \Rightarrow A}{\mathcal{E}; \Gamma \vdash MN : (b \doteq 0 \mapsto A) \Rightarrow A} \quad \frac{}{\mathcal{E}; \Gamma \vdash MNP : A}$$

4.2 Totality of mem

We want to prove that the function $\text{mem } a\ p := \text{test } p\ a\ 1 \parallel \text{test } p\ a\ 0$ is total from the proof Tot_{test} of totality of test. That is, we assume

$$\text{Tot}_{\text{test}} : \forall p \in \text{FVal}. \forall a \in \text{Atom}. \forall b \in \text{Bool}. \text{test } p\ a\ b \in \text{Bool}$$

and we want to prove

$$\text{Tot}_{\text{mem}} : \forall p \in \text{FVal}. \forall a \in \text{Atom}. \text{mem } a\ p \in \text{Bool}$$

for a suitable proof term Tot_{mem} .

$$\begin{aligned}
 & [\forall] p^\kappa \\
 & [\lambda] x : p \in \text{FVal} \\
 & [\forall] a^t \\
 & [\lambda] y : a \in \text{Atom} \\
 & [\forall] Z^{t \rightarrow o} \\
 & [\lambda] u : Z 1 \\
 & [\lambda] v : Z 0 \\
 & \text{Tot}_{\text{test}}\ x\ y : \forall b \in \text{Bool}. \text{test } a\ b\ p \in \text{Bool} \\
 & \text{true} : 1 \in \text{Bool} \\
 & \text{Tot}_{\text{test}}\ x\ y\ \text{true} : \text{test } a\ 1\ p \in \text{Bool} \equiv \forall Y^o. Y 1 \Rightarrow Y 0 \Rightarrow Y(\text{test } a\ 1\ p) \\
 & \text{Tot}_{\text{test}}\ x\ y\ \text{true} : Z(1 \parallel \text{test } a\ 0\ p) \Rightarrow Z(0 \parallel \text{test } a\ 0\ p) \Rightarrow Z(\text{test } a\ 1\ p \parallel \text{test } a\ 0\ p) \simeq Z 1 \Rightarrow Z(\text{test } a\ 0\ p) \Rightarrow Z(\text{test } a\ 1\ p \parallel \text{test } a\ 0\ p) \\
 & \text{Tot}_{\text{test}}\ x\ y\ \text{true}\ u : Z(\text{test } a\ 0\ p) \Rightarrow Z(\text{test } a\ 1\ p \parallel \text{test } a\ 0\ p) \\
 & \text{false} : 0 \in \text{Bool} \\
 & \text{Tot}_{\text{test}}\ x\ y\ \text{false} : \text{test } a\ 0\ p \in \text{Bool} \equiv \forall Y^o. Y 1 \Rightarrow Y 0 \Rightarrow Y(\text{test } a\ 0\ p) \\
 & \text{Tot}_{\text{test}}\ x\ y\ \text{false} : Z 1 \Rightarrow Z 0 \Rightarrow Z(\text{test } a\ 0\ p) \\
 & \text{Tot}_{\text{test}}\ x\ y\ \text{false}\ u\ v : Z(\text{test } a\ 0\ p) \\
 & \text{Tot}_{\text{test}}\ x\ y\ \text{true}\ u(\text{Tot}_{\text{test}}\ x\ y\ \text{false}\ u\ v) : Z(\text{test } a\ 1\ p \parallel \text{test } a\ 0\ p) \simeq Z(\text{mem } a\ p) \\
 & \lambda u v. \text{Tot}_{\text{test}}\ x\ y\ \text{true}\ u(\text{Tot}_{\text{test}}\ x\ y\ \text{false}\ u\ v) : \forall Z. Z 1 \Rightarrow Z 0 \Rightarrow Z(\text{mem } a\ p) \equiv \text{mem } a\ p \in \text{Bool} \\
 & \lambda x y u v. \text{Tot}_{\text{test}}\ x\ y\ \text{true}\ u(\text{Tot}_{\text{test}}\ x\ y\ \text{false}\ u\ v) : \forall p \in \text{FVal}. \forall a \in \text{Atom}. \text{mem } a\ p \in \text{Bool}
 \end{aligned}$$

4.3 Totality of eval

We want to find a proof term Tot_{eval} such that

$$\text{Tot}_{\text{eval}} : \forall p \in \text{FVal}. \forall c \in \text{Comp}. \forall b \in \text{Bool}. \text{eval } c \ b \ p \in \text{Bool}$$

$$\begin{aligned}
& [\forall] p^t \\
& [\lambda] x_p : p \in \text{FVal} \\
& [\forall] c^t \\
& [\lambda] x_c : c \in \text{Comp} \\
& [\forall] b^t \\
& [\lambda] y : b \in \text{Bool} \\
& \text{false} : 0 \in \text{Bool} \simeq \text{eval } \perp \ 1 \ p \in \text{Bool} \\
& \text{true} : 1 \in \text{Bool} \simeq \text{eval } \perp \ 0 \ p \in \text{Bool} \\
& \lambda y. \text{false true} : \text{eval } \perp \ b \ p \in \text{Bool} \\
& \lambda y. \text{y false true} : \forall b \in \text{Bool}. \text{eval } \perp \ b \ p \in \text{Bool} \\
& [\forall] a^t \\
& [\lambda] z : a \in \text{Atom} \\
& [\forall] b^t \\
& [\lambda] y : b \in \text{Bool} \\
& \text{Tot}_{\text{test}} x_p z y : \text{test } p \ a \ b \in \text{Bool} \simeq \text{eval } a \ b \ p \in \text{Bool} \\
& \lambda z y. \text{Tot}_{\text{test}} x_p z y : \forall b \in \text{Bool}. \forall a \in \text{Atom}. \text{eval } a \ b \ p \in \text{Bool} \\
& [\forall] c_1^t \\
& [\forall] c_2^t \\
& [\lambda] y_1 : \forall b \in \text{Bool}. \text{eval } c_1 \ b \ p \in \text{Bool} \\
& [\lambda] y_2 : \forall b \in \text{Bool}. \text{eval } c_2 \ b \ p \in \text{Bool} \\
& [\forall] b^t \\
& [\lambda] x : b \in \text{Bool} \\
& y_2 \ x : \text{eval } c_2 \ b \ p \in \text{Bool} \\
& \lambda u v. x v u : \text{not } b \in \text{Bool} \\
& y_1 (\lambda u v. x v u) : \text{eval } c_1 (\text{not } b) \ p \in \text{Bool} \\
& [\forall] Z^{t \rightarrow o} \\
& [\lambda] z_1 : Z 1 \\
& [\lambda] z_2 : Z 0 \\
& y_1 (\lambda u v. x v u) z_1 z_2 : Z (\text{eval } c_1 (\text{not } b) \ p) \simeq Z (\text{eval } c_1 (\text{not } b) \ p \parallel 0) \\
& z_1 : Z 1 \simeq Z (\text{eval } c_1 (\text{not } b) \ p \parallel 1) \\
& y_2 \ x z_1 (y_1 (\lambda u v. x v u) z_1 z_2) : Z (\text{eval } c_1 (\text{not } b) \ p \parallel \text{eval } c_2 \ b \ p) \simeq Z (\text{eval } (c_1 \Rightarrow c_2) \ b \ p) \\
& \lambda z_1 z_2. y_2 \ x z_1 (y_1 (\lambda u v. x v u) z_1 z_2) : \text{eval } (c_1 \Rightarrow c_2) \ b \ p \in \text{Bool} \\
& \lambda x z_1 z_2. y_2 \ x z_1 (y_1 (\lambda u v. x v u) z_1 z_2) : \forall b \in \text{Bool}. \text{eval } (c_1 \Rightarrow c_2) \ b \ p \in \text{Bool} \\
& \lambda y_1 y_2 x z_1 z_2. y_2 \ x z_1 (y_1 (\lambda u v. x v u) z_1 z_2) \\
& \quad : \forall c_1^t c_2^t. (\forall b \in \text{Bool}. \text{eval } c_1 \ b \ p \in \text{Bool}) \Rightarrow (\forall b \in \text{Bool}. \text{eval } c_2 \ b \ p \in \text{Bool}) \Rightarrow (\forall b \in \text{Bool}. \text{eval } (c_1 \Rightarrow c_2) \ b \ p \in \text{Bool}) \\
& x_c (\lambda y. \text{y false true}) (\lambda z y. \text{Tot}_{\text{test}} x_p z y) (y_1 y_2 x z_1 z_2. y_2 \ x z_1 (y_1 (\lambda u v. x v u) z_1 z_2)) : \forall b \in \text{Bool}. \text{eval } c \ b \ p \in \text{Bool} \\
& \lambda x_p x_c. x_c (\lambda y. \text{y false true}) (\lambda z y. \text{Tot}_{\text{test}} x_p z y) (y_1 y_2 x z_1 z_2. y_2 \ x z_1 (y_1 (\lambda u v. x v u) z_1 z_2)) \\
& \quad : \forall p \in \text{FVal}. \forall c \in \text{Comp}. \forall b \in \text{Bool}. \text{eval } c \ b \ p \in \text{Bool}
\end{aligned}$$

The proof for $\text{Tot}_{\text{subHtree}}$ is similar to this one and is omitted here.

4.4 Merging of subH

We want to prove the following result

$$\forall p. \forall a \in \text{Atom}. \text{mem } a \neq 0 \mapsto \text{subH}(pa^1) \Rightarrow \text{subH}(pa^0) \Rightarrow \text{subH } p$$

```

[ [V] p'
[ [V] at
[λ] xa : a ∈ Atom
[Eq.] mem a p = 0
[λ] x : subH(pa1)
[λ] y : subH(pa0)
[Let] x1 : t1 ∈ Tree
[Let] x2 : subHtree(pa1) t1 = 1
[Let] y1 : t2 ∈ Tree
[Let] y2 : subHtree(pa0) t2 = 1
Node xa x1 y1 : Node a t1 t2 ∈ Tree
[ [V] Zo
[λ] u : Z(subHtree p (Node a t1 t2)) ≈ Z( subHtree(pa1) t1 && subHtree(pa0) t2)
x2 : Z( subHtree(pa1) t1 && subHtree(pa0) t2) ⇒ Z(1 && subHtree(pa0) t2)
x2 u : Z(1 && subHtree(pa0) t2) ≈ Z( subHtree(pa0) t2)
y2 : Z( subHtree(pa0) t2) ⇒ Z 1
y2 (x2 u) : Z 1
y2 o x2 ≡ λu. y2 (x2 u) : ∀Zo. Z(subHtree p (Node a t1 t2)) ⇒ Z 1 ≡ subHtree p (Node a t1 t2) = 1
⟨Node xa x1 y1, y2 o x2⟩ : subH p
let (y1, y2) := y in ⟨Node xa x1 y1, y2 o x2⟩ : subH p
let (x1, x2) := x in let (y1, y2) := y in ⟨Node xa x1 y1, y2 o x2⟩ : subH p
λxaxy. let (x1, x2) := x in let (y1, y2) := y in ⟨Node xa x1 y1, y2 o x2⟩
: ∀p'. ∀a ∈ Atom. mem a p ≠ 0 ⇒ subH(pa1) ⇒ subH(pa0) ⇒ subH p

```

We let

$$\text{merge} := \lambda x_a xy. \text{let } (x_1, x_2) := x \text{ in let } (y_1, y_2) := y \text{ in } \langle \text{Node } x_a x_1 y_1, y_2 o x_2 \rangle .$$

4.5 Monotonicity of test

We want to prove that

$$\forall p^t \forall q^t \forall a^t \forall b^t. \text{test } p \ a \ b = 1 \Rightarrow \text{test}(p \cup q) \ a \ b = 1$$

$$\begin{aligned} & [\forall] p^t \\ & [\forall] q^t \\ & [\forall] a^t \\ & [\forall] b^t \\ & [\lambda] x : \text{test } p \ a \ b = 1 \\ & [\forall] Z^{t \rightarrow o} \\ & [\lambda] y : Z(\text{test}(p \cup q) \ a \ b) \simeq \text{test } a \ b \ p \parallel \text{test } a \ b \ q \\ & [xy : Z(1 \parallel \text{test } a \ b \ q) \simeq Z \ 1] \\ & [\lambda y. \ xy : \forall Z^{t \rightarrow o}. Z(\text{test}(p \cup q) \ a \ b) \Rightarrow Z \ 1 \equiv \text{test}(p \cup q) \ a \ b = 1] \\ & [\lambda xy. \ xy : \forall p^t \forall q^t \forall a^t \forall b^t. \text{test } a \ b \ p = 1 \Rightarrow \text{test}(p \cup q) \ a \ b = 1] \end{aligned}$$

and we write $\text{Mon}_{\text{test}} := \lambda xy. \ xy$.

4.6 Monotonicity of eval and subHtree

We want to prove that

$$\text{Mon}_{\text{eval}} : \forall p^t. \forall c \in \text{Comp}. \forall b \in \text{Bool}. \text{eval } c \ b \ p = 1 \Rightarrow \text{eval } c \ b \ (p \cup q) = 1$$

and

$$\text{Mon}_{\text{subHtree}} : \forall p^t. \forall t \in \text{Tree}. \forall b \in \text{Bool}. \text{subHtree } p \ t = 1 \Rightarrow \text{subHtree}(p \cup q) \ t = 1$$

for suitable proof terms Mon_{eval} and $\text{Mon}_{\text{subHtree}}$. Since both proofs follow the same pattern, we will focus only on the first one.

Induction principle on quantifier-free formulas To prove that eval is monotonic, we will need to use the *induction principle* of quantifier-free proposition. It is very similar to the relativization predicate $c \in \text{Comp}$ except that recursive arguments are guarded by relativization. More precisely, it is a proof term ind_{Comp} of the formula

$$\begin{aligned} c \in \text{Comp} \Rightarrow \forall Z^{t \rightarrow o}. Z \Downarrow \Rightarrow \\ (\forall a \in \text{Atom}. Z \ a) \Rightarrow \\ (\forall \underline{c_1 \in \text{Comp}}. \forall \underline{c_2 \in \text{Comp}}. Z \ c_1 \Rightarrow Z \ c_2 \Rightarrow Z(c_1 \Rightarrow c_2)) \Rightarrow \\ Z \ c \end{aligned}$$

(The differences with Comp are underlined.)

It can be easily built using the proof term of $c \in \text{Comp}$ on the proposition $P \ c := c \in \text{Comp} \Rightarrow Z \ c$.

$$\begin{aligned}
& [\forall] p^t \\
& [\forall] c^t \\
& [\lambda] x_c : c \in \text{Comp} \\
& [\forall] b^t \\
& [\lambda] x_b : b \in \text{Bool} \\
& [\lambda] y : \text{eval } \perp\!\!\!\perp b p = 1 \\
& y : \text{eval } \perp\!\!\!\perp b p = 1 \simeq \text{not } b = 1 \simeq \text{eval } \perp\!\!\!\perp b (p \cup q) = 1 \\
& \lambda x_b y. y : \forall b \in \text{Bool}. \text{eval } \perp\!\!\!\perp b p = 1 \Rightarrow \text{eval } \perp\!\!\!\perp b (p \cup q) = 1 \\
& [\forall] b^t \\
& [\lambda] x_b : b \in \text{Bool} \\
& [\forall] p^t \\
& [\forall] a^t \\
& [\lambda] u : a \in \text{Atom} \\
& [\lambda] v : \text{eval } a b p = 1 \simeq \text{test } a b p = 1 \\
& \text{Mon}_{\text{test}} v : \text{test}(p \cup q) a b = 1 \simeq \text{eval } a b (p \cup q) = 1 \\
& \lambda x_b u v. \text{Mon}_{\text{test}} v : \forall b \in \text{Bool}. \forall a \in \text{Atom}. \text{eval } a b p = 1 \Rightarrow \text{eval } a b (p \cup q) = 1 \\
& [\forall] c_1^t \\
& [\lambda] x_1 : c_1 \in \text{Comp} \\
& [\forall] c_2^t \\
& [\lambda] x_2 : c_2 \in \text{Comp} \\
& [\lambda] u : \forall b \in \text{Bool}. \forall p^t. \text{eval } c_1 b p = 1 \Rightarrow \text{eval } c_1 b (p \cup q) = 1 \\
& [\lambda] v : \forall b \in \text{Bool}. \forall p^t. \text{eval } c_2 b p = 1 \Rightarrow \text{eval } c_2 b (p \cup q) = 1 \\
& [\forall] b^t \\
& [\lambda] x_b : b \in \text{Bool} \\
& [\lambda] w : \text{eval}(c_1 \Rightarrow c_2) b p = 1 \simeq \text{eval } c_1 (\text{not } b) p \parallel \text{eval } c_2 b p = 1 \\
& \lambda u v. x_n v u : \text{not } b \in \text{Bool} \\
& \text{Tot}_{\text{eval}} x_1 (\lambda u v. x_n v u) : \text{eval } c_1 (\text{not } b) p \in \text{Bool} \\
& \text{I Tot}_{\text{eval}} x_1 (\lambda u v. x_n v u) : \text{eval } c_1 (\text{not } b) p \in \text{Bool}' \\
& [\text{Eq.}] \text{ eval } c_1 (\text{not } b) p = 1 \\
& u x_b : \text{eval } c_1 (\text{not } b) p = 1 \Rightarrow \text{eval } c_1 (\text{not } b) (p \cup q) = 1 \simeq 1 = 1 \Rightarrow \text{eval } c_1 (\text{not } b) (p \cup q) = 1 \\
& u x_b \text{ I} : \text{eval } c_1 (\text{not } b) (p \cup q) = 1 \\
& [\forall] Z^{t \rightarrow o} \\
& [\lambda] z : Z(\text{eval}(c_1 \Rightarrow c_2) b (p \cup q)) \simeq Z(\text{eval } c_1 (\text{not } b) (p \cup q) \parallel \text{eval } c_1 b (p \cup q)) \\
& [u x_b \text{ I} z : Z(1 \parallel \text{eval } c_1 b (p \cup q)) \simeq Z 1 \\
& \lambda z. u x_b \text{ I} z : \text{eval}(c_1 \Rightarrow c_2) b (p \cup q) = 1 \\
& \lambda z. u x_b \text{ I} z : \text{eval } c_1 (\text{not } b) p \doteq 1 \mapsto \text{eval}(c_1 \Rightarrow c_2) b (p \cup q) = 1 \\
& [\text{Eq.}] \text{ eval } c_1 (\text{not } b) p = 0 \\
& w : \text{eval } c_1 (\text{not } b) p \parallel \text{eval } c_2 b p = 1 \simeq 0 \parallel \text{eval } c_2 b p = 1 \simeq \text{eval } c_2 b p = 1 \\
& v x_b w : \text{eval } c_2 b (p \cup q) = 1 \\
& [\forall] Z^{t \rightarrow o} \\
& [\lambda] z : Z(\text{eval}(c_1 \Rightarrow c_2) b (p \cup q)) \simeq Z(\text{eval } c_1 (\text{not } b) (p \cup q) \parallel \text{eval } c_2 b (p \cup q)) \\
& [v x_b w z : Z(\text{eval } c_1 (\text{not } b) (p \cup q) \parallel 1) \simeq Z 1 \\
& \lambda z. v x_b w z : \text{eval}(c_1 \Rightarrow c_2) b (p \cup q) = 1 \\
& \lambda z. v x_b w z : \text{eval } c_1 (\text{not } b) p \doteq 0 \mapsto \text{eval}(c_1 \Rightarrow c_2) b (p \cup q) = 1 \\
& \text{I Tot}_{\text{eval}} x_1 (\lambda u v. x_n v u) (\lambda z. u x_b \text{ I} z) (\lambda z. v x_b w z) : \text{eval}(c_1 \Rightarrow c_2) b (p \cup q) = 1 \\
& \lambda x_1 x_2 u v x_b w. \text{I Tot}_{\text{eval}} x_1 (\lambda u v. x_n v u) (\lambda z. u x_b \text{ I} z) (\lambda z. v x_b w z) : \dots \\
& \text{ind}_{\text{Comp}} x_c (\lambda x_b y. y) (\lambda x_b u v. \text{Mon}_{\text{test}} v) (\lambda x_1 x_2 u v x_b w. \text{I Tot}_{\text{eval}} x_1 (\lambda u v. x_n v u) (\lambda z. u x_b \text{ I} z) (\lambda z. v x_b w z)) \\
& : \forall b \in \text{Bool}. \text{eval } c b p = 1 \Rightarrow \text{eval } c b (p \cup q) = 1 \\
& \lambda x_c. \text{ind}_{\text{Comp}} x_c (\lambda x_b y. y) (\lambda x_b u v. \text{Mon}_{\text{test}} v) (\lambda x_1 x_2 u v x_b w. \text{I Tot}_{\text{eval}} x_1 (\lambda u v. x_n v u) (\lambda z. u x_b \text{ I} z) (\lambda z. v x_b w z)) \\
& : \forall c \in \text{Comp}. \forall b \in \text{Bool}. \text{eval } c b p = 1 \Rightarrow \text{eval } c b (p \cup q) = 1
\end{aligned}$$

4.7 A proof of upward closure of FVal

We need to enrich the interface for finite relations over $\text{Atom} \times \text{Bool}$ with an intersection operation $_ \cap _$ with the following properties: for all a, b, p and q ,

$$p \cap \emptyset \simeq \emptyset \quad p \cap (p \cup q) \simeq p \quad p \cap (qa^b) \simeq_{\text{mem } a \text{ } p=1} (p \cap q)a^b \quad p \cap (qa^b) \simeq_{\text{mem } a \text{ } p=0} p \cap q$$

These congruences could be derived from more primitive ones like the one we have for \cup but these are exactly the ones we need to prove the upward closure of FVal.

We let

$$T^b := \forall r^\kappa. \forall a \in \text{Atom}. \text{ mem } a \text{ } r = 0 \mapsto Z(p \cap r) \Rightarrow Z(p \cap (ra^b))$$

$[\forall] p^\kappa$	
$[\forall] q^\kappa$	
$[\lambda] z : pq \in \text{FVal}$	
$[\forall] Z^o$	
$[\lambda] e_0 : Z \emptyset$	
$[\lambda] e_+ : \forall r^\kappa. \forall a \in \text{Atom}. \text{ mem } a \text{ } r = 0 \mapsto Zr \Rightarrow Z(ra^1)$	
$[\lambda] e_- : \forall r^\kappa. \forall a \in \text{Atom}. \text{ mem } a \text{ } r = 0 \mapsto Zr \Rightarrow Z(ra^0)$	
$\delta_{20} e_0 : Z(p \cap \emptyset)$	
$z(\delta_{20} e_0) : T^1 \Rightarrow T^0 \Rightarrow Z(p \cap (p \cup q))$	
$[\forall] r^\kappa$	
$[\forall] a'$	
$[\lambda] x_a : a \in \text{Atom}$	
$[\text{Eq.}] \text{ mem } a \text{ } r = 0$	
$[\lambda] y : Z(p \cap r)$	
$e_+ : \text{mem } a \text{ } p = 0 \mapsto Z(p \cap r) \Rightarrow Z((p \cap r)a^1) \simeq Z(p \cap r) \Rightarrow Z((p \cap r)a^1)$	
$e_+ y : Z((p \cap r)a^1)$	
$[\text{Eq.}] \text{ mem } ar = 1$	
$\delta_{22} : \text{mem } a \text{ } r \doteq 1 \mapsto (p \cap r)a^1 = p \cap (ra^1) \simeq (p \cap r)a^1 = p \cap (ra^1) \equiv \forall Z. Z((p \cap r)a^1) \Rightarrow Z(p \cap (ra^1))$	
$\delta_{22}(e_+ y) : Z(p \cap (ra^1))$	
$\delta_{22}(e_+ y) : \text{mem } ar \doteq 1 \mapsto Z(p \cap (ra^1))$	
$[\text{Eq.}] \text{ mem } ar = 0$	
$\delta_{23} : \text{mem } a \text{ } r \doteq 0 \mapsto p \cap r = p \cap (ra^1) \simeq p \cap r = p \cap (ra^1) \equiv \forall Z. Z(p \cap r) \Rightarrow Z(p \cap (ra^1))$	
$\delta_{23} y : Z(p \cap (ra^1))$	
$\delta_{23} y : \text{mem } ar \doteq 0 \mapsto Z(p \cap (ra^1))$	
$\delta_{10} x_a : \text{mem } a \text{ } p \in \text{Bool}$	
$I(\delta_{10} x_a) : \text{mem } a \text{ } p \in \text{Bool}' \equiv \forall Z^o. (\text{mem } a \text{ } p \doteq 1 \mapsto Z) \Rightarrow (\text{mem } a \text{ } p \doteq 0 \mapsto Z) \Rightarrow Z$	
$I(\delta_{10} x_a)(\delta_{22}(e_+ y))(\delta_{23} y) : Z(p \cap (ra^1))$	
$\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_+ y))(\delta_{23} y) : \forall r. \forall a \in \text{Atom}. \text{ mem } a \text{ } r = 0 \mapsto Z(p \cap r) \Rightarrow Z(p \cap (ra^1)) \equiv T^1$	
<i>idem</i>	
$\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_- y))(\delta_{23} y) : \forall r. \forall a \in \text{Atom}. \text{ mem } a \text{ } r = 0 \mapsto Z(p \cap r) \Rightarrow Z(p \cap (ra^0)) \equiv T^0$	
$z(\delta_{20} e_0)(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_+ y))(\delta_{23} y))(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_- y))(\delta_{23} y)) : Z(p \cap (p \cup q))$	
$\delta_{21}(z(\delta_{20} e_0)(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_+ y))(\delta_{23} y))(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_- y))(\delta_{23} y))) : Zp$	
$\lambda e_0 e_+ e_- . \delta_{21}(z(\delta_{20} e_0)(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_+ y))(\delta_{23} y))(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_- y))(\delta_{23} y))) : p \in \text{FVal}$	
$\lambda e_0 e_+ e_- . \delta_{21}(z(\delta_{20} e_0)(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_+ y))(\delta_{23} y))(\lambda x_a y. I(\delta_{10} x_a)(\delta_{22}(e_- y))(\delta_{23} y))) : \forall p \forall q. pq \in \text{FVal} \Rightarrow p \in \text{FVal}$	

4.8 A proof of $C[1]$

$[\forall] Z^o$	
$[\lambda] x : Z 1$	
$[\lambda] y : \forall p^\kappa. \forall a \in \text{Atom}. \text{ mem } a \text{ } p = 0 \mapsto Zp \Rightarrow Z(pa^1)$	
$[\lambda] z : \forall p^\kappa. \forall a \in \text{Atom}. \text{ mem } a \text{ } p = 0 \mapsto Zp \Rightarrow Z(pa^0)$	
$x : Z 1$	
$\lambda xyz. z : \forall Z^o. Z 1 \Rightarrow (\dots) \Rightarrow (\dots) \Rightarrow Z 1 \equiv 1 \in \text{FVal}$	
$\langle \lambda xyz. z, I \rangle : 1 \in \text{FVal} \wedge (\text{subH } 1 \Rightarrow \text{subH } 1) \equiv C[1]$	

4.9 A proof of $\forall p^\kappa q^\kappa. C[pq] \Rightarrow C[p]$

Recall that we have the following proof terms:

$$\begin{aligned} \text{Mon}_{\text{subHtree}} &: \forall p^t q^t. \text{subHtree } p t = 1 \Rightarrow \text{subHtree}(p \cup q) t = 1 \\ \text{Up}_{\text{FVal}} &: \forall p^t q^t. (p \cup q) \in \text{FVal} \Rightarrow p \in \text{FVal} \end{aligned}$$

$$\begin{aligned} &[\forall] p^\kappa \\ &[\forall] p^\kappa \\ &[\lambda] c : C[pq] := pq \in \text{FVal} \wedge \text{subH}(pq) \Rightarrow \text{subH } \emptyset \\ &[\text{Let}] c_1 : pq \in \text{FVal} \\ &[\text{Let}] c_2 : \text{subH}(pq) \Rightarrow \text{subH } \emptyset \\ &\text{Up}_{\text{FVal}} c_1 : p \in \text{FVal} \\ &[\lambda] x : \text{subH } p \\ &[\text{Let}] x_1 : t \in \text{Tree} \\ &[\text{Let}] x_2 : \text{subHtree } p t = 1 \\ &\text{Mon}_{\text{subHtree}} x_2 : \text{subHtree}(p \cup q) t = 1 \\ &\langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle : \text{subH } (pq) \\ &\langle c_2, \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle \rangle : \text{subH } \emptyset \\ &\text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle : \text{subH } \emptyset \\ &\lambda x. \text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle : \text{subH } p \Rightarrow \text{subH } \emptyset \\ &\langle \text{Up}_{\text{FVal}} c_1, \lambda x. \text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle \rangle : C[p] \\ &\text{let } (c_1, c_2) := c \text{ in } \langle \text{Up}_{\text{FVal}} c_1, \lambda x. \text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle \rangle : C[p] \\ &\lambda c. \text{let } (c_1, c_2) := c \text{ in } \langle \text{Up}_{\text{FVal}} c_1, \lambda x. \text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle \rangle : C[pq] \Rightarrow C[p] \end{aligned}$$

5 Absoluteness of datatypes

5.1 Tree is absolute

The ideas are the same as for \mathbb{N} . We recall the definition of $t \in \text{Tree}$:

$$\begin{aligned} t \in \text{Tree} &:= \forall Z^{t \rightarrow o}. (\forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) \Rightarrow \\ &(\forall t'_1 \forall t'_2. \forall a \in \text{Atom}. Z t'_1 \Rightarrow Z t'_2 \Rightarrow Z(\text{Node } a t'_1 t'_2)) \Rightarrow \\ &Z t \end{aligned}$$

To ease writing, we set

$$\text{closZ} := \forall t'_1 \forall t'_2. \forall a \in \text{Atom}. Z t'_1 \Rightarrow Z t'_2 \Rightarrow Z(\text{Node } a t'_1 t'_2)$$

Because of its length, the proof is split in two parts. The first one is the main proof and the second one is meant to be plugged into it.

$[\lambda] z : p \Vdash t \in \text{Tree}$
$[\lambda] c : C[p]$
$[\forall] Z^{\iota \rightarrow o}$
$[\lambda] x : \forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})$
$[\lambda] y : \text{closZ}$
$z : (p \Vdash (\forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) \Rightarrow \text{closZ} \Rightarrow Z t) [Z/Z^* r]$
$\gamma_3 z : (p \Vdash \forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) [Z/Z^* r] \Rightarrow (p \Vdash \text{closZ} \Rightarrow Z t) [Z/Z^* r]$
$[\forall] (\vec{v})^\iota$
$[\forall] q_{\vec{v}}^k$
$[\lambda] w : q_{\vec{v}} \Vdash \vec{v} \in \text{Term}$
$\xi_{\text{Term}} w : C[q_{\vec{v}}] \Rightarrow \vec{v} \in \text{Term}$
$[\forall] r^\kappa$
$[\lambda] c' : C[pq_{\vec{v}} r]$
$\alpha_2(\alpha_1 c') : C[q_{\vec{v}}]$
$\xi_{\text{Term}} w(\alpha_2(\alpha_1 c')) : \vec{v} \in \text{Term}$
$x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c'))) : Z(\text{Leaf } \vec{v})$
$\lambda c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c'))) : \forall r. C[pq_{\vec{v}} r] \Rightarrow Z(\text{Leaf } \vec{v}) \equiv (pq_{\vec{v}} \Vdash Z(\text{Leaf } \vec{v})) [Z/Z^* r]$
$\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c'))) : \forall \vec{v} q_{\vec{v}}. (q_{\vec{v}} \Vdash \vec{v} \in \text{Term}) \Rightarrow (pq_{\vec{v}} \Vdash Z(\text{Leaf } \vec{v})) [Z/Z^* r]$
$\gamma_1(\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c')))) : \forall \vec{v}. (p \Vdash \vec{v} \in \text{Term} \Rightarrow Z(\text{Leaf } \vec{v})) [Z/Z^* r] \simeq (p \Vdash \forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) [Z/Z^* r]$
$\gamma_3 z(\gamma_1(\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c'))))) : (p \Vdash \text{closZ} \Rightarrow Z t) [Z/Z^* r]$
$\gamma_3(z(\gamma_3 z(\gamma_1(\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c')))))) : (p \Vdash \text{closZ}) [Z/Z^* r] \Rightarrow (p \Vdash Z t) [Z/Z^* r]$
$[\forall] t'_1$
$[\forall] t'_2$
$[\forall] a'$
See below
$\gamma_1(\lambda u. \gamma_1(\lambda v_1. \gamma_1(\lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c')))))) : p \Vdash \text{closZ}$
Write T the term just above so that $T : p \Vdash \text{closZ}$
$\gamma_3(z(\gamma_3 z(\gamma_1(\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c')))))) T : (p \Vdash Z t) [Z/Z^* r] \equiv \forall r^\kappa. C[pr] \Rightarrow Z t$
$\alpha_7 c : C[p1]$
$\gamma_3(z(\gamma_3 z(\gamma_1(\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c')))))) T(\alpha_7 c) : Z t$
$\lambda x y. \gamma_3(z(\gamma_3 z(\gamma_1(\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c')))))) T(\alpha_7 c) : \forall Z. (\forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) \Rightarrow \text{closZ} \Rightarrow Z t \equiv t \in \text{Tree}$
$\lambda z c x y. \gamma_3(z(\gamma_3 z(\gamma_1(\lambda w c'. x(\xi_{\text{Term}} w(\alpha_2(\alpha_1 c')))))) T(\alpha_7 c) : (p \Vdash t \in \text{Tree}) \Rightarrow (C[p] \Rightarrow t \in \text{Tree})$

For the second part of the proof, recall that we have the following context:

$$\begin{aligned} z &: p \Vdash t \in \text{Tree} \\ c &: C[p] \\ x &: \forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v}) \\ y &: \text{closZ} \end{aligned}$$

$$\begin{aligned}
& \boxed{[\forall] t_1^t} \\
& \boxed{[\forall] t_2^t} \\
& \boxed{[\forall] a^t} \\
& \boxed{[\forall] q_0^\kappa} \\
& [\lambda] u : q_0 \models a \in \text{Atom} \\
& \xi_{\text{Atom}} u : C[q_0] \Rightarrow a \in \text{Atom} \\
& \boxed{[\forall] q_1^\kappa} \\
& [\lambda] v_1 : (q_1 \models Z t_1)[Z/Z^*r] \simeq \forall r^\kappa. C[q_1 r] \Rightarrow Z t_1 \\
& \boxed{[\forall] q_2^\kappa} \\
& [\lambda] v_2 : (q_2 \models Z t_2)[Z/Z^*r] \simeq \forall r^\kappa. C[q_2 r] \Rightarrow Z t_2 \\
& \boxed{[\forall] r^\kappa} \\
& [\lambda] c' : C[pq_0 q_1 q_2 r] \\
& \alpha_2(\alpha_1^3 c') : C[q_0] \\
& \xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')) : a \in \text{Atom} \\
& y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))) : Z t_1 \Rightarrow Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2) \\
& \alpha_9(\alpha_{16} c') : C[q_1 r] \\
& v_1(\alpha_9(\alpha_{16} c')) : Z t_1 \\
& y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))) : Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2) \\
& \alpha_{10} c' : C[q_2 r] \\
& v_2(\alpha_{10} c') : Z t_2 \\
& y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10} c')) : Z(\text{Node } a t_1 t_2) \\
& \lambda c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c')))) : (pq_0 q_1 q_2 \models Z(\text{Node } a t_1 t_2))[Z/Z^*r] \\
& \lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c')))) : (\forall q_2^\kappa. (q_2 \models Z t_2) \Rightarrow (pq_0 q_1 q_2 \models Z(\text{Node } a t_1 t_2))[Z/Z^*r]) \\
& \gamma_1(\lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c'))))) : (pq_0 q_1 \models Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2))[Z/Z^*r] \\
& \lambda v_1. \gamma_1(\lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c'))))) \\
& \quad : (\forall q_1^\kappa. (q_1 \models Z t_1) \Rightarrow (pq_0 q_1 \models Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2))[Z/Z^*r]) \\
& \gamma_1(\lambda v_1. \gamma_1(\lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c')))))) : (pq_0 \models Z t_1 \Rightarrow Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2))[Z/Z^*r] \\
& \lambda u. \gamma_1(\lambda v_1. \gamma_1(\lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c')))))) \\
& \quad : \forall q_0^\kappa. (q_0 \models a \in \text{Atom}) \Rightarrow (pq_0 \models Z t_1 \Rightarrow Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2))[Z/Z^*r] \\
& \gamma_1(\lambda u. \gamma_1(\lambda v_1. \gamma_1(\lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c')))))) \\
& \quad : (p \models a \in \text{Atom} \Rightarrow Z t_1 \Rightarrow Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2))[Z/Z^*r]) \\
& \gamma_1(\lambda u. \gamma_1(\lambda v_1. \gamma_1(\lambda v_2 c'. y(\xi_{\text{Atom}} u(\alpha_2(\alpha_1^3 c')))(v_1(\alpha_9(\alpha_{16} c')))(v_2(\alpha_{10}(\alpha_{16} c')))))) : p \models \text{closZ}
\end{aligned}$$

$[\lambda] z : C[p] \Rightarrow t \in \text{Tree}$
$[\forall] (Z^*)^{t \rightarrow \kappa \rightarrow o}$
$[\forall] q_L^\kappa$
$[\lambda] x : q_L \Vdash \forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})$
$[\forall] q_N^\kappa$
$[\lambda] y : q_N \Vdash \text{closZ}$
$\beta_3 x : \forall(\vec{v})^t. q_L q_N \Vdash \vec{v} \in \text{Term} \Rightarrow Z(\text{Leaf } \vec{v})$
$\gamma_3 (\beta_3 x) : \forall(\vec{v})^t. (q_L q_N \Vdash \vec{v} \in \text{Term}) \Rightarrow (q_L q_N \Vdash Z(\text{Leaf } \vec{v}))$
$[\forall] (\vec{v})^t$
$[\lambda] u : \vec{v} \in \text{Term}$
$\lambda_{_.u} : C[q_L q_N] \Rightarrow \vec{v} \in \text{Term}$
$\xi'_{\text{Term}} (\lambda_{_.u}) : q_L q_N \Vdash \vec{v} \in \text{Term}$
$\gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u})) : q_L q_N \Vdash Z(\text{Leaf } \vec{v})$
$\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u})) : \forall \vec{v} \in \text{Term}. q_L q_N \Vdash Z(\text{Leaf } \vec{v})$
$\beta_4 y : \forall t_1^t \forall t_2^t \forall a^t. q_L q_N \Vdash a \in \text{Atom} \Rightarrow Z t_1 \Rightarrow Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2)$
$\gamma_3 (\beta_4 y) : \forall t_1^t \forall t_2^t \forall a^t. (q_L q_N \Vdash a \in \text{Atom}) \Rightarrow (q_L q_N \Vdash Z t_1 \Rightarrow Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2))$
$[\forall] t_1^t$
$[\forall] t_2^t$
$[\forall] a^t$
$[\lambda] u : a \in \text{Atom}$
$[\lambda] v : q_L q_N \Vdash Z t_1$
$[\lambda] w : q_L q_N \Vdash Z t_2$
$\lambda_{_.u} : C[q_L q_N] \Rightarrow a \in \text{Atom}$
$\xi'_{\text{Atom}} (\lambda_{_.u}) : q_L q_N \Vdash a \in \text{Atom}$
$\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u})) : q_L q_N \Vdash Z t_1 \Rightarrow Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2)$
$\gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) : (q_L q_N \Vdash Z t_1) \Rightarrow (q_L q_N \Vdash Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2))$
$\gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v : q_L q_N \Vdash Z t_2 \Rightarrow Z(\text{Node } a t_1 t_2)$
$\gamma_3 (\gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) : (q_L q_N \Vdash Z t_2) \Rightarrow (q_L q_N \Vdash Z(\text{Node } a t_1 t_2))$
$\gamma_3 (\gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w : q_L q_N \Vdash Z(\text{Node } a t_1 t_2)$
$\lambda u v w. \gamma_3 (\gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w : \forall t_1^t \forall t_2^t \forall a \in \text{Atom}. (q_L q_N \Vdash Z t_1) \Rightarrow (q_L q_N \Vdash Z t_2) \Rightarrow q_L q_N \Vdash Z(\text{Node } a t_1 t_2)$
$[\forall] r^k$
$[\lambda] c : C[pq_L q_N r]$
$\alpha_1^3 c : C[p]$
$z(\alpha_1^3 c) : t \in \text{Tree}$
$z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w : q_L q_N \Vdash Z t$
$z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w : C[q_L q_N r] \Rightarrow Z^* t r$
$\alpha_{16} c : C[q_L q_N r]$
$z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c) : Z^* t r$
$\lambda c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c)$
$: \forall r^k. C[pq_L q_N r] \Rightarrow Z^* t r \equiv pq_L q_N \Vdash Z t$
$\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c)$
$: \forall q_N^\kappa. (q_N \Vdash \text{closZ}) \Rightarrow (pq_L q_N \Vdash Z t)$
$\gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c)) : pq_L \Vdash \text{closZ} \Rightarrow Z t$
$\lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c))$
$: \forall q_L^\kappa. (q_L \Vdash \forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) \Rightarrow (pq_L \Vdash \text{closZ} \Rightarrow Z t)$
$\gamma_1 (\lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c)))$
$: p \Vdash (\forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) \Rightarrow \text{closZ} \Rightarrow Z t$
$\gamma_1 (\lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c)))$
$: \forall Z^*. p \Vdash (\forall \vec{v} \in \text{Term}. Z(\text{Leaf } \vec{v})) \Rightarrow \text{closZ} \Rightarrow Z t) \simeq p \Vdash t \in \text{Tree}$
$\lambda z. \gamma_1 (\lambda x. \gamma_1 (\lambda y c. z(\alpha_1^3 c) (\lambda u. \gamma_3 (\beta_3 x) (\xi'_{\text{Term}} (\lambda_{_.u}))) (\lambda u v w. \gamma_3 (\gamma_3 (\beta_4 y) (\xi'_{\text{Atom}} (\lambda_{_.u}))) v) w) (\alpha_{16} c)))$
$: (C[p] \Rightarrow t \in \text{Tree}) \Rightarrow (p \Vdash t \in \text{Tree})$

5.2 subH is absolute

Remember that

$$\text{subH } p := \exists t \in \text{Tree}. \text{ subHtree } p t = 1 \equiv (\forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp) \Rightarrow \perp$$

$$\begin{aligned}
& [\lambda] z : q \models \text{subH } p \\
& [\lambda] c : C[q] \\
& [\lambda] x : \forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \\
& \gamma_3 z : (q \models \forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp) \Rightarrow (q \models \perp) \\
& [[\forall] t^t] \\
& [\forall] q_T^\kappa \\
& [\lambda] u : q_T \models t \in \text{Tree} \\
& [[\forall] q_p^\kappa] \\
& [\lambda] v : q_p \models \text{subHtree } p t = 1 \\
& [\lambda] c' : C[qq_T q_p] \\
& \xi_{\text{Tree}} u : C[q_T] \Rightarrow t \in \text{Tree} \\
& \alpha_2(\alpha_1 c') : C[q_T] \\
& \xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')) : t \in \text{Tree} \\
& x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c'))) : \text{subHtree } p t = 1 \Rightarrow \perp \\
& \alpha_2 c' : C[q_p] \\
& \xi_{\text{eq}} v : C[q_p] \Rightarrow \text{subHtree } p t = 1 \\
& \xi_{\text{eq}} v(\alpha_2 c') : \text{subHtree } p t = 1 \\
& x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2(\alpha_1 c'))) : \perp \\
& \lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')) : C[qq_T q_p] \Rightarrow \perp \\
& \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')) : qq_T q_p \models \perp) \\
& \lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')) : \forall q_p^\kappa. (q_p \models \text{subHtree } p t = 1) \Rightarrow (qq_T q_p \models \perp) \\
& \gamma_1(\lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))) : qq_T \models \text{subHtree } p t = 1 \Rightarrow \perp \\
& \lambda u. \gamma_1(\lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))) : \forall q_T^\kappa. (q_T \models t \in \text{Tree}) \Rightarrow (qq_T \models \text{subHtree } p t = 1 \Rightarrow \perp) \\
& \gamma_1(\lambda u. \gamma_1(\lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c'))))) : q \models t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \\
& \gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))) : q \models \perp \\
& \xi_\perp(\gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))))) : C[q] \Rightarrow \perp \\
& \xi_\perp(\gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))))) c : \perp \\
& \lambda x. \xi_\perp(\gamma_3 z(\gamma_1(\lambda u. \gamma_1(\lambda v. \xi'_\perp(\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))))) c : \text{subH } p \\
& \lambda z c. z @ (\lambda^* u. (\lambda^* v. (\lambda c'. \langle \xi_{\text{Tree}} u(\alpha_2(\alpha_1(\alpha_1 c'))), \xi_{\text{eq}} v(\alpha_2(\alpha_1 c')) \rangle))) (\alpha_7 c) : (q \models \text{subH } p) \Rightarrow (C[q] \Rightarrow \text{subH } p)
\end{aligned}$$

$$\begin{array}{l}
[\lambda] z : q \models \text{subH } p \\
[\lambda] c : C[q] \\
[[\lambda] x : \forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \\
\gamma_3 z : (q \models \forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp) \Rightarrow (q \models \perp) \\
[[\forall] t' \\
[[\forall] q_T^\kappa \\
[\lambda] u : q_T \models t \in \text{Tree} \\
[[\forall] q_p^\kappa \\
[\lambda] v : q_p \models \text{subHtree } p t = 1 \\
[[\lambda] c' : C[qq_T q_p] \\
\xi_{\text{Tree}} u : C[q_T] \Rightarrow t \in \text{Tree} \\
\alpha_2(\alpha_1 c') : C[q_T] \\
\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')) : t \in \text{Tree} \\
x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c'))) : \text{subHtree } p t = 1 \Rightarrow \perp \\
\alpha_2 c' : C[q_p] \\
\xi_{\text{eq}} v : C[q_p] \Rightarrow \text{subHtree } p t = 1 \\
\xi_{\text{eq}} v(\alpha_2 c') : \text{subHtree } p t = 1 \\
x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2(\alpha_1 c'))) : \perp \\
\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')) : C[qq_T q_p] \Rightarrow \perp \\
\xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c'))) : qq_T q_p \models \perp \\
\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c'))) : \forall q_p^\kappa. (q_p \models \text{subHtree } p t = 1) \Rightarrow (qq_T q_p \models \perp) \\
\gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))) : qq_T \models \text{subHtree } p t = 1 \Rightarrow \perp \\
\lambda u. \gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))) : q \models t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \\
\gamma_1 (\lambda u. \gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c'))))) : q \models t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \\
\gamma_3 z (\gamma_1 (\lambda u. \gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))) : q \models \perp \\
\xi_\perp (\gamma_3 z (\gamma_1 (\lambda u. \gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))))) : C[q] \Rightarrow \perp \\
\xi_\perp (\gamma_3 z (\gamma_1 (\lambda u. \gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))))) c : \perp \\
\lambda x. \xi_\perp (\gamma_3 z (\gamma_1 (\lambda u. \gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))))) c : \text{subH } p \\
\lambda z cx. \xi_\perp (\gamma_3 z (\gamma_1 (\lambda u. \gamma_1 (\lambda v. \xi'_\perp (\lambda c'. x(\xi_{\text{Tree}} u(\alpha_2(\alpha_1 c')))(\xi_{\text{eq}} v(\alpha_2 c')))))))) c : (q \models \text{subH } p) \Rightarrow (C[q] \Rightarrow \text{subH } p)
\end{array}$$

$$\begin{aligned}
& [\lambda] z : C[q] \Rightarrow \text{subH } p \\
& [\forall] r^k \\
& [\lambda] x : r \models \forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \simeq \forall t. r \models t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \\
& \beta_4 x : \forall t. qr \models t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp \\
& \gamma_3(\beta_4 x) : \forall t. (qr \models t \in \text{Tree}) \Rightarrow (qr \models \text{subHtree } p t = 1 \Rightarrow \perp) \\
& [\lambda] c : C[qr] \\
& \alpha_1 c : C[q] \\
& z(\alpha_1 c) : \text{subH } p \equiv (\forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp) \Rightarrow \perp \\
& [\forall] t^t \\
& [\lambda] u : t \in \text{Tree} \\
& [\lambda] v : \text{subHtree } p t = 1 \\
& \lambda_. u : C[qr] \Rightarrow t \in \text{Tree} \\
& \xi'_{\text{Tree}}(\lambda_. u) : qr \models t \in \text{Tree} \\
& \gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)) : qr \models \text{subHtree } p t = 1 \Rightarrow \perp \\
& \gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u))) : (qr \models \text{subHtree } p t = 1) \Rightarrow (qr \models \perp) \\
& \lambda_. v : C[qr] \Rightarrow \text{subHtree } p t = 1 \\
& \xi_{\text{eq}}(\lambda_. v) : qr \models \text{subHtree } p t = 1 \\
& \gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v)) : qr \models \perp \\
& \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v))) : C[qr] \Rightarrow \perp \\
& \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v)))c : \perp \\
& \lambda u v. \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v)))c : \forall t. (t \in \text{Tree}) \Rightarrow (\text{subHtree } p t = 1) \Rightarrow \perp \\
& z(\alpha_1 c)(\lambda u v. \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v))))c : \perp \\
& \lambda c. z(\alpha_1 c)(\lambda u v. \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v))))c : C[qr] \Rightarrow \perp \\
& \xi'_{\perp}(\lambda c. z(\alpha_1 c)(\lambda u v. \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v))))c) : qr \models \perp \\
& \lambda x. \xi'_{\perp}(\lambda c. z(\alpha_1 c)(\lambda u v. \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v))))c) \\
& \quad : \forall r. (r \models \forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp) \Rightarrow (qr \models \perp) \\
& \gamma_1(\lambda x. \xi'_{\perp}(\lambda c. z(\alpha_1 c)(\lambda u v. \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v))))c)) \\
& \quad : q \models (\forall t. t \in \text{Tree} \Rightarrow \text{subHtree } p t = 1 \Rightarrow \perp) \Rightarrow \perp \simeq q \models \text{subH } p \\
& \lambda z. \gamma_1(\lambda x. \xi'_{\perp}(\lambda c. z(\alpha_1 c)(\lambda u v. \xi_{\perp}(\gamma_3(\gamma_3(\beta_4 x)(\xi'_{\text{Tree}}(\lambda_. u)))(\xi_{\text{eq}}(\lambda_. v))))c)) : (C[q] \Rightarrow \text{subH } p) \Rightarrow (q \models \text{subH } p)
\end{aligned}$$

5.3 FVal is absolute

The definition of FVal is

$$\begin{aligned}
p \in \text{FVal} := & \forall Z^{t \rightarrow o}. Z \emptyset \Rightarrow \\
& (\forall p''. \forall a \in \text{Atom}. \text{mem } a p' =_t 0 \mapsto Z p' \Rightarrow Z(p' a^1)) \Rightarrow \\
& (\forall p''. \forall a \in \text{Atom}. \text{mem } a p' =_t 1 \mapsto Z p' \Rightarrow Z(p' a^0)) \Rightarrow \\
& Z p
\end{aligned}$$

To ease writing, we set

$$\begin{aligned}
\text{closZ}^+ &:= \forall r' \forall a \in \text{Atom}. \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1) \\
\text{closZ}^- &:= \forall r' \forall a \in \text{Atom}. \text{mem } a r =_t 1 \mapsto Z r \Rightarrow Z(ra^0).
\end{aligned}$$

$\boxed{[\lambda] z : q \Vdash p \in \text{FVal}}$
 $[\lambda] c : C[q]$
 $[[\forall] Z^{\iota \rightarrow o}]$
 $[\lambda] x : Z \emptyset$
 $[\lambda] y^+ : \text{closZ}^+$
 $[\lambda] y^- : \text{closZ}^-$
 $z : (q \Vdash Z \emptyset \Rightarrow \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p) [Z/Z^* r]$
 $\gamma_3 z : (q \Vdash Z \emptyset) [Z/Z^* r] \Rightarrow (q \Vdash \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p) [Z/Z^* r]$
 $\lambda_. x : \forall r^k. C[qr] \Rightarrow Z \emptyset \simeq (q \Vdash Z \emptyset) [Z/Z^* r]$
 $\gamma_3 z (\lambda_. x) : (q \Vdash \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p) [Z/Z^* r]$
 $\gamma_3 (\gamma_3 z (\lambda_. x)) : (q \Vdash \text{closZ}^+) [Z/Z^* r] \Rightarrow (q \Vdash \text{closZ}^- \Rightarrow Z p) [Z/Z^* r]$
 $[[\forall] (p')^\iota]$
 $[[\forall] a^\iota]$
 $[[\forall] q_0^\kappa]$
 $[\lambda] u : q_0 \Vdash a \in \text{Atom}$
 $\xi_{\text{Atom}} u : C[q_0] \Rightarrow a \in \text{Atom}$
 $[[\forall] q_1^\kappa]$
 $[\lambda] v : (q_1 \Vdash Z r) [Z/Z^* r] \simeq \forall s^\kappa. C[q_1 s] \Rightarrow Z r$
 $[[\forall] s^\kappa]$
 $[\lambda] c' : C[qq_0 q_1 s]$
 $\alpha_2(\alpha_1^2 c') : C[q_0]$
 $\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c')) : a \in \text{Atom}$
 $y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) : \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1)$
 $y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) : Z r \Rightarrow Z(ra^1)$
 $\alpha_{10} c' : C[q_1 s]$
 $v(\alpha_{10} c') : Z r$
 $y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c')) : Z(ra^1)$
 $\lambda c'. y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c')) : \forall s^\kappa. C[qq_0 q_1 s] \Rightarrow Z(ra^1) \equiv (qq_0 q_1 \Vdash Z(ra^1)) [Z/Z^* r]$
 $\lambda vc'. y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c')) : \forall q_1^\kappa. (q_1 \Vdash Z r) [Z/Z^* r] \Rightarrow (qq_0 q_1 \Vdash Z(ra^1)) [Z/Z^* r]$
 $\gamma_1 (\lambda vc'. y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c'))) : (qq_0 \Vdash Z r \Rightarrow Z(ra^1)) [Z/Z^* r]$
 $\gamma_1 (\lambda vc'. y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c')))$
 $\quad : \text{mem } a r =_t 0 \mapsto (qq_0 \Vdash Z r \Rightarrow Z(ra^1)) [Z/Z^* r] \simeq (qq_0 \Vdash \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1)) [Z/Z^* r]$
 $\lambda u. \gamma_1 (\lambda vc'. y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c'))) : \forall q_0^\kappa. q_0 \Vdash a \in \text{Atom} \Rightarrow (qq_0 \Vdash \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1)) [Z/Z^* r]$
 $\gamma_1 (\lambda u. \gamma_1 (\lambda vc'. y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c')))) : (q \Vdash a \in \text{Atom} \Rightarrow \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1)) [Z/Z^* r]$
 $\gamma_1 (\lambda u. \gamma_1 (\lambda vc'. y^+ (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c')))) : (q \Vdash \text{closZ}^+) [Z/Z^* r]$
 $\text{Write } T^{+/-} := \gamma_1 (\lambda u. \gamma_1 (\lambda vc'. y^{+/-} (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c'))))$
 $T^+ : (q \Vdash \text{closZ}^+) [Z/Z^* r]$
 idem
 $T^- : (q \Vdash \text{closZ}^-) [Z/Z^* r]$
 $\gamma_3 (\gamma_3 z (\lambda_. x)) T^+ : (q \Vdash \text{closZ}^- \Rightarrow Z t) [Z/Z^* r]$
 $\gamma_3 (\gamma_3 (\gamma_3 z (\lambda_. x)) T^+) : (q \Vdash \text{closZ}^-) [Z/Z^* r] \Rightarrow (q \Vdash Z p) [Z/Z^* r]$
 $\gamma_3 (\gamma_3 (\gamma_3 z (\lambda_. x)) T^+) T^- : (q \Vdash Z p) [Z/Z^* r] \equiv \forall s^\kappa. C[q s] \Rightarrow Z p$
 $\alpha_7 c : C[q 1]$
 $\gamma_3 (\gamma_3 (\gamma_3 z (\lambda_. x)) T^+) T^- (\alpha_7 c) : Z p$
 $\lambda xy^+ y^- . \gamma_3 (\gamma_3 (\gamma_3 z (\lambda_. x)) T^+) T^- (\alpha_7 c) : \forall Z^{\iota \rightarrow o}. Z \emptyset \Rightarrow \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p \equiv p \in \text{FVal}$
 $\lambda z c x y^+ y^- . \gamma_3 (\gamma_3 (\gamma_3 z (\lambda_. x)) T^+) T^- (\alpha_7 c) : (q \Vdash p \in \text{FVal}) \Rightarrow (C[q] \Rightarrow p \in \text{FVal})$

remembering that $T^{+/-} := \gamma_1 (\lambda u. \gamma_1 (\lambda vc'. y^{+/-} (\xi_{\text{Atom}} u (\alpha_2(\alpha_1^2 c'))) (v(\alpha_{10} c'))))$

$\boxed{[\lambda] z : C[q] \Rightarrow p \in \text{FVal}}$
 $\boxed{[\forall] (Z^*)^{t \rightarrow \kappa \rightarrow o}}$
 $\boxed{[\forall] q_0^\kappa}$
 $\boxed{[\lambda] x : q_0 \Vdash Z \emptyset}$
 $\boxed{[\forall] (q^+)^\kappa}$
 $\boxed{[\lambda] y^+ : q^+ \Vdash \text{closZ}^+}$
 $\boxed{[\forall] (q^-)^\kappa}$
 $\boxed{[\lambda] y^- : q^- \Vdash \text{closZ}^-}$
 $\beta_3 x : q_0 q^+ q^- \Vdash Z \emptyset$
 $\beta_4 y^+ : \forall r^t \forall a^t. q_0 q^+ q^- \Vdash a \in \text{Atom} \Rightarrow \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1)$
 $\gamma_3(\beta_4 y^+) : \forall r^t \forall a^t. (q_0 q^+ q^- \Vdash a \in \text{Atom}) \Rightarrow (q_0 q^+ q^- \Vdash \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1))$
 $\boxed{[\forall] r^t}$
 $\boxed{[\forall] a^t}$
 $\boxed{[\lambda] u : a \in \text{Atom}}$
 $\boxed{[\text{Eq.}] \text{ mem } a r =_t 0}$
 $\boxed{[\lambda] v : q_0 q^+ q^- \Vdash Z r}$
 $\lambda_- u : C[q_0 q^+ q^-] \Rightarrow a \in \text{Atom}$
 $\xi'_{\text{Atom}}(\lambda_- u) : q_0 q^+ q^- \Vdash a \in \text{Atom}$
 $\gamma_3(\beta_4 y^+) (\xi'_{\text{Atom}}(\lambda_- u)) : q_0 q^+ q^- \Vdash \text{mem } a r =_t 0 \mapsto Z r \Rightarrow Z(ra^1) \simeq \text{mem } a r =_t 0 \mapsto q_0 q^+ q^- \Vdash Z r \Rightarrow Z(ra^1)$
 $\gamma_3(\beta_4 y^+) (\xi'_{\text{Atom}}(\lambda_- u)) : q_0 q^+ q^- \Vdash Z r \Rightarrow Z(ra^1)$
 $\gamma_3(\gamma_3(\beta_4 y^+) (\xi'_{\text{Atom}}(\lambda_- u))) : (q_0 q^+ q^- \Vdash Z r) \Rightarrow (q_0 q^+ q^- \Vdash Z(ra^1))$
 $\gamma_3(\gamma_3(\beta_4 y^+) (\xi'_{\text{Atom}}(\lambda_- u))) v : q_0 q^+ q^- \Vdash Z(r a^1)$
 $\lambda u v. \gamma_3(\beta_4 y^+) (\xi'_{\text{Atom}}(\lambda_- u)) v : \forall r^t \forall a \in \text{Atom}. \text{ mem } a r =_t 0 \mapsto (q_0 q^+ q^- \Vdash Z r) \Rightarrow (q_0 q^+ q^- \Vdash Z(ra^1))$
 $\text{Write } U^{+/-} := \lambda u v. \gamma_3(\beta_4 y^{+/-}) (\xi'_{\text{Atom}}(\lambda_- u)) v$
 $U^+ : \forall r^t \forall a \in \text{Atom}. \text{ mem } a r =_t 0 \mapsto (q_0 q^+ q^- \Vdash Z r) \Rightarrow (q_0 q^+ q^- \Vdash Z(ra^1))$
idem
 $U^- : \forall r^t \forall a \in \text{Atom}. \text{ mem } a r =_t 0 \mapsto (q_0 q^+ q^- \Vdash Z r) \Rightarrow (q_0 q^+ q^- \Vdash Z(ra^0))$
 $\boxed{[\forall] r^\kappa}$
 $\boxed{[\lambda] c : C[qq_0 q^+ q^- r]}$
 $\alpha_1^4 c : C[q]$
 $z(\alpha_1^4 c) : p \in \text{FVal}$
 $z(\alpha_1^4 c)(\beta_3 x) U^+ U^- : q_0 q^+ q^- \Vdash Z p \equiv \forall r^\kappa. C[q_0 q^+ q^- r] \Rightarrow Z^* p r$
 $\alpha_6(\alpha_{16}(\alpha_5 c)) : C[q_0 q^+ q^- r]$
 $z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c))) : Z^* p r$
 $\lambda c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c))) : \forall r^\kappa. C[qq_0 q^+ q^- r] \Rightarrow Z^* p r \equiv pq_0 q^+ q^- \Vdash Z p$
 $\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c))) : \forall q^-. q^- \Vdash \text{closZ}^- \Rightarrow qq_0 q^+ q^- \Vdash Z p$
 $\gamma_1(\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c)))) : qq_0 q^+ \Vdash \text{closZ}^- \Rightarrow Z p$
 $\lambda y^+. (\gamma_1(\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c))))) : \forall q^+. (q^+ \Vdash \text{closZ}^+) \Rightarrow (qq_0 q^+ \Vdash \text{closZ}^- \Rightarrow Z p)$
 $\gamma_1(\lambda y^+. (\gamma_1(\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c)))))) : qq_0 \Vdash \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p$
 $\lambda x. \gamma_1(\lambda y^+. (\gamma_1(\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c)))))) : \forall q_0. (q_0 \Vdash Z \emptyset) \Rightarrow (qq_0 \Vdash \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p)$
 $\gamma_1(\lambda x. \gamma_1(\lambda y^+. (\gamma_1(\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c))))))) : q \Vdash Z \emptyset \Rightarrow \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p$
 $\gamma_1(\lambda x. \gamma_1(\lambda y^+. (\gamma_1(\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c))))))) : \forall Z^*. q \Vdash Z \emptyset \Rightarrow \text{closZ}^+ \Rightarrow \text{closZ}^- \Rightarrow Z p \simeq q \Vdash p \in \text{FVal}$
 $\lambda z. \gamma_1(\lambda x. \gamma_1(\lambda y^+. (\gamma_1(\lambda y^- c. z(\alpha_1^4 c)(\beta_3 x) U^+ U^- (\alpha_6(\alpha_{16}(\alpha_5 c))))))) : (C[q] \Rightarrow p \in \text{FVal}) \Rightarrow (q \Vdash p \in \text{FVal})$

remembering that $U^{+/-} := \lambda u v. \gamma_3(\beta_4 y^{+/-}) (\xi'_{\text{Atom}}(\lambda_- u)) v$.

5.4 C is absolute

The set C is defined by $C[p] := p \in \text{FVal} \wedge (\text{subH } p \Rightarrow \text{subH } \emptyset)$.

We have

$$\begin{aligned} \xi_{\text{FVal}} &: (q \Vdash p \in \text{FVal}) \Rightarrow (C[q] \Rightarrow p \in \text{FVal}) \\ \xi'_{\text{FVal}} &: (C[q] \Rightarrow p \in \text{FVal}) \Rightarrow (q \Vdash p \in \text{FVal}) \\ \xi_{\text{subH}} &: (q \Vdash \text{subH } p) \Rightarrow (C[q] \Rightarrow \text{subH } p) \\ \xi'_{\text{subH}} &: (C[q] \Rightarrow \text{subH } p) \Rightarrow (q \Vdash \text{subH } p) \end{aligned}$$

but also

$$\begin{aligned}\xi_{\text{subH}} : (q \models \text{subH } \emptyset) &\Rightarrow (C[q] \Rightarrow \text{subH } \emptyset) \\ \xi'_{\text{subH}} : (C[q] \Rightarrow \text{subH } \emptyset) &\Rightarrow (q \models \text{subH } \emptyset)\end{aligned}$$

We then get

$$\begin{aligned}\xi \wedge \xi_{\text{FVal}} (\xi \Rightarrow \xi'_{\text{subH}} \xi_{\text{subH}}) : (q \models C[p]) &\Rightarrow (C[q] \Rightarrow C[p]) \\ \xi' \wedge \xi'_{\text{FVal}} (\xi' \Rightarrow \xi_{\text{subH}} \xi'_{\text{subH}}) : (C[q] \Rightarrow C[p]) &\Rightarrow (q \models C[p])\end{aligned}$$

6 Proof of forcing the extra axiom (A)

We want to prove $1 \models \forall a \in \text{Atom}. (\forall pb. p \in G \Rightarrow b \in \text{Bool} \Rightarrow \text{test } p \ a \ b = 1 \Rightarrow \perp) \Rightarrow \perp$.

6.1 A textual proof

We first give a textual proof before the formal one in order to convey the essential ideas.

Proof. Given a, q_a, q and proof terms

$$\begin{aligned}x : q_a \models a \in \text{Atom} \\ y : q \models \forall p \forall b. p \in G \Rightarrow b \in \text{Bool} \Rightarrow \text{test } p \ a \ b = 1 \Rightarrow \perp,\end{aligned}$$

we want to prove $1 q_a q \models \perp$. Since 1 is neutral for the product, this is the same as proving that $q_a q \models \perp$. With repeated use of γ_3 , we can turn y into $y' := \lambda u v. \gamma_3(\gamma_3(\beta_4 y) u) v$ which is a proof term for

$$\forall q' pb. (q' q \models p \in G) \Rightarrow (q' q \models b \in \text{Bool}) \Rightarrow (q' q \models \text{test } p \ a \ b = 1) \Rightarrow (q' q \models \perp).$$

Since test is total (assumed in the interface), mem is total too and we have $\text{mem } a(q_a q) \in \text{Bool}$. We then proceed by case analysis, noting that $\text{mem } a(q_a q) = 1$ can itself be decomposed into $\text{test}(q_a q) a 1 = 1$ or $\text{test}(q_a q) a 0 = 1$.

- $\text{test}(q_a q) a 1 = 1$: We take $p := q_a q$ and $b := 1$. We want to use y' to prove $q_a q \models \perp$ so that we have to prove its premises:
 - $\lambda x. x : q_a q \models q_a q \in G \equiv q_a q \leq q_a q$,
 - $\gamma_1(\lambda u. \gamma_1(\lambda v. \beta_4(\beta_3 u))) : q_a q \models 1 \in \text{Bool}$,
 - $\xi'_{\text{eq}}(\lambda x. x) : q_a q \models \text{test}(q_a q) a 1 = 1$ because the equality holds in the equational theory.
- $\text{test}(q_a q) a 0 = 1$: It is similar to the previous case.
- $\text{mem } a(q_a q) = 0$: We use ξ'_\perp and we are left with $C[q_a q] \Rightarrow \perp$ to prove. We first prove that we have $C[(q_a a^1)q] \Rightarrow \perp$ by using first ξ_\perp then y' with $q := q_a a^1$, $p := a^1$ and $b := 1$.
 - $\alpha_{10} \circ \alpha_9 : (q_a a^1)q \models a^1 \in G \equiv (q_a a^1)q \leq a^1$ because product is the glb for \leq ,
 - $(q_a a^1)q \models 1 \in \text{Bool}$ proved as before,
 - $(q_a a^1)q \models \text{test } a^1 a 1 = 1$ proved as before.

In a similar fashion, we prove that $C[(q_a a^0)q] \Rightarrow \perp$.

Back to the proof of $C[q_a q] \Rightarrow \perp$. Assume $C[q_a q]$. Applying the proof term of $C[q_a a^1 q] \Rightarrow \perp$, we have to prove $C[q_a a^1 q] \equiv q_a a^1 q \in \text{FVal} \wedge (\text{subH } q_a a^1 q \Rightarrow \text{subH } \emptyset)$. The first part present no difficulty because we have $C[q_a q]$ and $\text{mem } a(q_a q) = 0$: we just need to apply the second constructor of FVal. For the second one, we assume $\text{subH } q_a a^1 q'$ and we want to prove $\text{subH } \emptyset$. Instead we choose to prove \perp . Applying again the same method with $C[q_a a^0 q] \Rightarrow \perp$, we end up proving $\text{subH } \emptyset$ with $\text{subH } q_a a^1 q$ and $\text{subH } q_a a^0 q$ as extra hypotheses. $\text{subH } q_a a^1 q$ and $\text{subH } q_a a^0 q$ allow us to prove $\text{subH } q_a q$. Finally, from $C[q_a q]$, we deduce $\text{subH } q_a q \Rightarrow \text{subH } \emptyset$, which allows us to conclude. ■

6.2 The formal proof

Given q^* and

$$\begin{aligned} f : \forall pb. q \Vdash p \in G \Rightarrow b \in \text{Bool} \Rightarrow M = N \Rightarrow \perp \\ x_b : q \Vdash b \in \text{Bool} \end{aligned}$$

we have

$$\left[\begin{array}{l} [\text{Eq.}] M = N \\ I : q \Vdash q \in G \equiv q \leq q \\ f @ I @ x_b : q \Vdash M = N \Rightarrow \perp \\ I : M = M \simeq M = N \\ I^* : q \Vdash M = N \\ f @ I @ x_b @ I^* : \perp \\ f @ I @ x_b @ I^* : M \doteq N \mapsto q \Vdash \perp \end{array} \right]$$

We will use the lemma four times in the coming proof, always with $x_b = \text{true}^*$ or $x_b = \text{false}^*$ and $M = \text{test } q \ a \ b$.

$\boxed{[\forall] r^\kappa}$
 $[\forall] q_a^\kappa$
 $[\forall] q_f^\kappa$
 $[\lambda] c : C[q(q_a(q_f r))]$
 $[\forall] a^t$
 $[\lambda] x_a : q_a \text{ IF } a \in \text{Atom}$
 $[\lambda] f : q_f \text{ IF } \forall pb. p \in G \Rightarrow b \in \text{Bool} \Rightarrow \text{test } p \ a \ b = 1 \Rightarrow \perp \simeq \forall pb. q \text{ IF } p \in G \Rightarrow b \in \text{Bool} \Rightarrow \text{test } p \ a \ b = 1 \Rightarrow \perp$
 $\alpha_1(\alpha_2 c) : C[q_a]$
 $\alpha_6(\alpha_6 c) : C[((qq_a)q_f)r]$
 $\hat{c} := \alpha_1(\alpha_6(\alpha_6 c)) : C[((qq_a)q_f)]$
 $[\text{Let}] c_1 : (qq_a)q_f \in \text{FVal}$
 $[\text{Let}] c_2 : \text{subH}(qq_a)q_f \Rightarrow \text{subH } \emptyset$
 $\hat{x}_a := \xi_{\text{Atom}} x_a (\alpha_1(\alpha_2 c)) : a \in \text{Atom}$
 $\text{Tot}_{\text{mem}} c_1 \hat{x}_a : \text{mem } a (qq_a)q_f \in \text{Bool}$
First case: mem $a (qq_a)q_f = 1$
 $\text{Tot}_{\text{test}} c_1 \hat{x}_a \text{ true} : \text{test } (qq_a)q_f a 1 \in \text{Bool}$
 $[\text{Eq.}] \text{mem } a (qq_a)q_f = 1$
First sub-case: test $(qq_a)q_f a 1 = 1$
 $[\text{Eq.}] \text{test } (qq_a)q_f a 1 = 1$
 $\beta_4 f : (qq_a)q_f \text{ IF } (qq_a)q_f \in G \Rightarrow 1 \in \text{Bool} \Rightarrow \text{test } (qq_a)q_f a 1 = 1 \Rightarrow \perp$
 $\text{true}^* : (qq_a)q_f \text{ IF } 1 \in \text{Bool}$
 $((\beta_4 f) @ I @ \text{true}^* @ I^*) : \text{test } (qq_a)q_f a 1 \doteq 1 \mapsto (qq_a)q_f \text{ IF } \perp \simeq (qq_a)q_f \text{ IF } \perp$
 $((\beta_4 f) @ I @ \text{true}^* @ I^*) c : \perp$
Second sub-case: test $(qq_a)q_f a 1 = 0$
 $[\text{Eq.}] \text{test } (qq_a)q_f a 1 = 0$
 From $\text{mem } a (qq_a)q_f =_E 1$ and $\text{test } (qq_a)q_f a 1 =_E 0$, we deduce $\text{test } (qq_a)q_f a 0 =_E 1$
 $\beta_4 f : (qq_a)q_f \text{ IF } (qq_a)q_f \in G \Rightarrow 0 \in \text{Bool} \Rightarrow \text{test } (qq_a)q_f a 0 = 1 \Rightarrow \perp$
 $\text{false}^* : (qq_a)q_f \text{ IF } 0 \in \text{Bool}$
 $((\beta_4 f) @ I @ \text{false}^* @ I^*) : \text{test } (qq_a)q_f a 0 \doteq 1 \mapsto (qq_a)q_f \text{ IF } \perp \simeq (qq_a)q_f \text{ IF } \perp$
 $((\beta_4 f) @ I @ \text{false}^* @ I^*) : \perp$
 if $\text{Tot}_{\text{test}} c_1 \hat{x}_a \text{ true}$ then $((\beta_4 f) @ I @ \text{true}^* @ I^*) c$ else $((\beta_4 f) @ I @ \text{false}^* @ I^*) c : \perp$
Second case: mem $a (qq_a)q_f = 0$
 $[\text{See below}]$
 $(\beta_3(\beta_4 f) @ I @ \text{true}^* @ I^*)(\alpha_7 \langle \text{consT } \hat{x}_a c_1, \lambda u. B \rangle) : C[((qq_a)q_f)a^1 1] : \perp$
 $\text{if } \text{Tot}_{\text{mem}} c_1 \hat{x}_a$
 then if $\text{Tot}_{\text{test}} c_1 \hat{x}_a \text{ true}$ then $((\beta_4 f) @ I @ \text{true}^* @ I^*) c$ else $((\beta_4 f) @ I @ \text{false}^* @ I^*) c$
 $\text{else } (\beta_3(\beta_4 f) @ I @ \text{true}^* @ I^*)(\alpha_7 \langle \text{consT } \hat{x}_a c_1, \lambda u. B \rangle) : \perp$
 $\text{let } (c_1, c_2) := \hat{c} \text{ in if } \text{Tot}_{\text{mem}} c_1 \hat{x}_a$
 then if $\text{Tot}_{\text{test}} c_1 \hat{x}_a \text{ true}$ then $((\beta_4 f) @ I @ \text{true}^* @ I^*) c$ else $((\beta_4 f) @ I @ \text{false}^* @ I^*) c$
 $\text{else } (\beta_3(\beta_4 f) @ I @ \text{true}^* @ I^*)(\alpha_7 \langle \text{constT } \hat{x}_a c_1, \lambda u. B \rangle) : \perp$
 $\lambda c x_a f.$
 $\text{let } (c_1, c_2) := \hat{c} \text{ in if } \text{Tot}_{\text{mem}} c_1 \hat{x}_a$
 then if $\text{Tot}_{\text{test}} c_1 \hat{x}_a \text{ true}$ then $((\beta_4 f) @ I @ \text{true}^* @ I^*) c$ else $((\beta_4 f) @ I @ \text{false}^* @ I^*) c$
 $\text{else } (\beta_3(\beta_4 f) @ I @ \text{true}^* @ I^*)(\alpha_7 \langle \text{consT } \hat{x}_a c_1, \lambda u. B \rangle)$
 $: q \text{ IF } \forall a \in \text{Atom}. (\forall pb. p \in G \Rightarrow b \in \text{Bool} \Rightarrow \text{test } p \ a \ b = 1 \Rightarrow \perp) \Rightarrow \perp$

For the second case, recall that we have

$$\begin{aligned}
& \text{mem } a(qq_a)q_f =_{\mathcal{E}} 0 \\
& c : C[q(q_a(q_f r))] \\
& c_1 : (qq_a)q_f \in \text{FVal} \\
& c_2 : \text{subH}(qq_a)q_f \Rightarrow \text{subH } \emptyset \\
& x_a : q_a \text{ IF } a \in \text{Atom} \\
& \hat{x}_a := \xi_{\text{Atom}} x_a (\alpha_1 (\alpha_2 c)) : a \in \text{Atom} \\
& f : \forall p b. q_f \text{ IF } p \in G \Rightarrow b \in \text{Bool} \Rightarrow \text{test } p a b = 1 \Rightarrow \perp
\end{aligned}$$

$$\begin{aligned}
& \hat{x}_a : a \in \text{Atom} \\
& \beta_3(\beta_4 f) : ((qq_a)q_f)a^1 \text{ IF } ((qq_a)q_f)a^1 \in G \Rightarrow 1 \in \text{Bool} \Rightarrow \text{test}((qq_a)q_f)a^1 a 1 = 1 \Rightarrow \perp \\
& \beta_3(\beta_4 f) @ I @ \text{true}^* @ I^* : ((qq_a)q_f)a^1 \text{ IF } \perp \\
& \beta_3(\beta_4 f) : ((qq_a)q_f)a^0 \text{ IF } ((qq_a)q_f)a^0 \in G \Rightarrow 0 \in \text{Bool} \Rightarrow \text{test}((qq_a)q_f)a^0 a 1 = 1 \Rightarrow \perp \\
& \beta_3(\beta_4 f) @ I @ \text{false}^* @ I^* : ((qq_a)q_f)a^0 \text{ IF } \perp \\
& \text{consT } \hat{x}_a c_1 : ((qq_a)q_f)a^1 \in \text{FVal} \\
& [\lambda] u : \text{subH}((qq_a)q_f)a^1 \\
& \text{consF } \hat{x}_a c_1 : ((qq_a)q_f)a^0 \in \text{FVal} \\
& [\lambda] v : \text{subH}((qq_a)q_f)a^0 \\
& \text{merge } \hat{x}_a u v : \text{subH}(qq_a)q_f \\
& c_2 (\text{merge } \hat{x}_a u v) : \text{subH } \emptyset \\
& \lambda v. c_2 (\text{merge } \hat{x}_a u v) : \text{subH}((qq_a)q_f)a^0 \Rightarrow \text{subH } \emptyset \\
& \langle \text{consF } \hat{x}_a c_1, \lambda v. c_2 (\text{merge } \hat{x}_a u v) \rangle : C[((qq_a)q_f)a^0] \\
& \alpha_7 \langle \text{consF } \hat{x}_a c_1, \lambda v. c_2 (\text{merge } \hat{x}_a u v) \rangle : C[((qq_a)q_f)a^0 1] \\
& B := (\beta_3(\beta_4 f) @ I @ \text{false}^* @ I^*) (\alpha_7 \langle \text{consF } \hat{x}_a c_1, \lambda v. c_2 (\text{merge } \hat{x}_a u v) \rangle) : \perp \\
& B : \text{subH } \emptyset \\
& \lambda u. B : \text{subH}((qq_a)q_f)a^1 \Rightarrow \text{subH } \emptyset \\
& \langle \text{consT } \hat{x}_a c_1, \lambda u. B \rangle : C[((qq_a)q_f)a^1] \\
& \alpha_7 \langle \text{consT } \hat{x}_a c_1, \lambda u. B \rangle : C[((qq_a)q_f)a^1 1] \\
& (\beta_3(\beta_4 f) @ I @ \text{true}^* @ I^*) (\alpha_7 \langle \text{consT } \hat{x}_a c_1, \lambda u. B \rangle : C[((qq_a)q_f)a^1 1]) : \perp
\end{aligned}$$

6.3 Simplifying the combinators

We now use a canonical implementation for finite valuations featuring associativity, commutativity and idempotency of union whose neutral element is \emptyset . For instance, one can take ordered list without repetition. This implies that most primitive combinators of the forcing structure are the identity. More precisely, only α_1 and α_2 are not the identity among the first 8 combinators which triggers a similar result for derived ones. All in all, there is only one real combinator, which we call α , that remove a part of the finite list (and thus is idempotent).

$$\begin{aligned}
& \alpha_1 = \alpha_2 = \alpha_9 = \alpha_{10} = \alpha_{14} = \alpha_{15} = \alpha_{16} = \alpha \\
& \alpha_3 = \alpha_4 = \alpha_5 = \alpha_6 = \alpha_7 = \alpha_8 = \alpha_{11} = \alpha_{12} = \alpha_{13} = I \\
& \beta_2 = \beta_3 = \beta_4 = \lambda x c. x(\alpha c) \\
& \gamma_1 = \lambda x c y. x y c \quad \gamma_2 = \gamma_3 = \lambda x c y. x c y \quad \gamma_4 = \lambda x c y. x(y(\alpha c))
\end{aligned}$$

In this context, the previous program becomes

$$\begin{aligned}
& \lambda c a f. \text{let } (c_1, c_2) := \alpha c \text{ in} \\
& \quad \text{if } \text{Tot}_{\text{mem}} c_1 \hat{x}_a \text{ then} \\
& \quad \quad \text{if } \text{Tot}_{\text{test}} c_1 \hat{x}_a \text{ true then } f(\alpha c) I \text{ true}^* I^* \text{ else } f(\alpha c) I \text{ false}^* I^* \\
& \quad \quad \text{else } f(\alpha \langle \text{consT } \hat{x}_a c_1, \lambda u. f(\alpha \langle \text{consF } \hat{x}_a c_1, \lambda v. c_2 (\text{merge } \hat{x}_a u v) \rangle) I \text{ false}^* I^* \rangle) I \text{ true}^* I^*
\end{aligned}$$

Let us now look at the implementation of α :

$$\lambda c. \text{let } (c_1, c_2) := c \text{ in } \langle \text{Up}_{\text{FVal}} c_1, \lambda x. \text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle \rangle$$

The term $\text{Mon}_{\text{subHtree}} : \forall pqt. \text{subHtree } pt = 1 \Rightarrow \text{subHtree } pq t = 1$ proves a true Horn formula that can be realized by the identity. Indeed, any Herbrand tree below p is also a Herbrand tree below pq . Thus, from a realizer point of view, we have

$$\begin{aligned}\alpha &:= \lambda c. \text{let } (c_1, c_2) := c \text{ in } \langle \text{Up}_{\text{FVal}} c_1, \lambda x. \text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, \text{Mon}_{\text{subHtree}} x_2 \rangle \rangle \\ &\equiv \lambda c. \text{let } (c_1, c_2) := c \text{ in } \langle \text{Up}_{\text{FVal}} c_1, \lambda x. \text{let } (x_1, x_2) := x \text{ in } c_2 \langle x_1, x_2 \rangle \rangle \\ &\equiv \lambda c. \text{let } (c_1, c_2) := c \text{ in } \langle \text{Up}_{\text{FVal}} c_1, \lambda x. c_2 x \rangle \\ &\equiv \lambda c. \text{let } (c_1, c_2) := c \text{ in } \langle \text{Up}_{\text{FVal}} c_1, c_2 \rangle\end{aligned}$$

This shows that the second component of $C[p]$ is never touched by α and the last line of the program becomes

$$f \langle \text{Up}_{\text{FVal}} \text{consT } \hat{x}_a c_1, \lambda u. f \langle \text{Up}_{\text{FVal}} \text{consF } \hat{x}_a c_1, \lambda v. c_2 (\text{merge } \hat{x}_a u v) \rangle \text{Ifalse}^* \text{I}^* \rangle \text{Itrue}^* \text{I}^*$$