



HAL
open science

Compensated integrability Applications to the Vlasov–Poisson equation

Denis Serre

► **To cite this version:**

Denis Serre. Compensated integrability Applications to the Vlasov–Poisson equation. 2018. ensl-01723852v1

HAL Id: ensl-01723852

<https://ens-lyon.hal.science/ensl-01723852v1>

Preprint submitted on 5 Mar 2018 (v1), last revised 5 May 2018 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Compensated integrability

Applications to the Vlasov–Poisson equation

Denis Serre
École Normale Supérieure de Lyon^{*}

Abstract

We extend and improve our analysis of divergence-free positive symmetric tensors (DPT). In particular, we show that the Vlasov–Poisson equation for a plasma is eligible for our theory, contrary to the opinion that we expressed in [8].

1 Introduction

Let $d \geq 2$ be an integer. In a previous paper [8], we began the study of divergence-free positive symmetric tensors, in short DPTs. These objects are measurable fields

$$T : \Omega \longrightarrow \mathbf{Sym}_d^+$$

over an open domain $\Omega \subset \mathbb{R}^d$ (or a torus $\Omega = \mathbb{R}^d / \Gamma$), with values in the cone of positive semi-definite matrices, whose row-wise divergence $\text{Div } T$ vanishes:

$$\forall i = 1, \dots, d \quad \sum_{j=1}^d \partial_j T_{ij} \equiv 0.$$

We also considered situations where the condition above is replaced by some control of $\text{Div } T$ in the space of vector fields with values in bounded measures.

In the periodic case, our main result was that when such a field is integrable, then $(\det T)^{\frac{1}{d-1}}$ is integrable too. This is remarkable since a product like $(T_{11} \cdots T_{dd})^{\frac{1}{d-1}}$ does not need to be integrable. Such a result can therefore be called a *Compensated Integrability*. At the algebraic level, the relation

^{*}U.M.P.A., UMR CNRS–ENSL # 5669. 46 allée d’Italie, 69364 Lyon cedex 07. France. denis.serre@ens-lyon.fr

[†]The author thanks Meiji University (Tokyo) and the Chinese University (Hong Kong) for their hospitality when a part of this paper was developed.

between the differential constraint and the function $\det^{\frac{1}{d-1}}$ is that the latter is concave in the characteristic directions of the operator Div . The inequality associated with the property of compensated integrability,

$$\int_{\mathbb{R}^d/\Gamma} (\det T)^{\frac{1}{d-1}} dx \leq \left(\det \int_{\mathbb{R}^d/\Gamma} T(x) dx \right)^{\frac{1}{d-1}},$$

is a *non-diagonal* extension of Gagliardo's Inequality. It tells us that the map $T \mapsto (\det T)^{\frac{1}{d-1}}$ is *Divergence- quasiconcave*, in the terminology of Fonseca & Müller [3].

For a bounded domain, we established a similar inequality, where the right-hand side involves instead a boundary integral. Despite its correctness, our Theorem 2.3 had two flaws. On the one hand, we proved only the functional inequality, but failed to establish the gain of integrability and had to make it an assumption. On the other hand, we imposed the useless restriction that the domain Ω be convex.

The paper below addresses several aspects of DPTs, concerning either general statements or their applications to Physics and Fluid dynamics.

In Section 2, we complete the results of [8] and give more elegant proofs, both in the periodic and in the bounded cases. We drop the assumption of convexity and establish the Compensated Integrability in all situations. We treat the case where Ω is a slab $(0, \tau) \times \mathbb{R}^{d-1}$, which is tailored for the applications to models of continuum mechanics, especially inviscid compressible fluids ; our statement can be applied whenever the total mass and energy of the fluid are finite. As shown in [8], this yields new *a priori* estimates, where the gain of integrability comes to the price of a time integration. We answer a question that was left aside, of whether the positivity of $T(x)$ is a meaningful assumption. The answer is positive when $d \geq 3$, as shown by an example, constructed by S. Klainerman & M. Machedon in the context of the wave equation.

Section 3 is a rather general study of those functions $T \mapsto f(T)$ which display a similar gain of integrability as $\det^{\frac{1}{d-1}}$. We first show that they must vanish on rank-one tensors $v \otimes v$. It is therefore natural to consider appropriate powers of *immanants*. Our main result is that among this collection, $\det^{\frac{1}{d-1}}$ is the only one to gain maximal integrability. This paragraph makes use of the classification of the locally integrable DPTs that are homogeneous of a given degree. We show in particular that this degree may not belong to the interval $(-d, 1-d)$. This is an indication that singularities of DPTs should not be too strong.

Section 4 is devoted to divergence-free symmetric tensors that occur in mathematical Physics ; they often come as energy-momentum tensors. We warn the reader that not all of them are positive. We mentioned above the case of the wave equation. We present here a few other examples, among which a rather interesting one is the non-linear Maxwell system of an electro-magnetic field ; it is remarkable that the symmetry of the tensor is a consequence of the invariance of the model under the action of the Lorentz group of special relativity. We also describe in full generality those first-order systems of conservation laws having the DPT form, and that are compatible with a convex entropy. We show that they can be described in terms of a single potential function.

Last, but not least, Section 5 deals with kinetic models of the Vlasov family in $d = 1 + n$ space-time dimensions. We hastily made a remark in [8], that our theory does not apply to the Vlasov–Poisson equation, despite the presence of a divergence-free symmetric tensor, because the latter is not positive (except in one space dimension). We show here that this claim followed from a wrong perspective: when the self-induced force $F = -\nabla\phi$ is given in terms of the density ρ by the Poisson formula¹ $\Delta\phi = \pm\rho$, we chose a tensor containing a diagonal block $F \otimes F - \frac{1}{2}|F|^2 I_n$, which is neither positive nor negative. However this choice is specific to the coupling above ; it could not be extended to a more general one when the potential is given by $\phi = \chi * \rho$, where χ is a radial kernel. We show here that the two first moments of the Vlasov equation can always be put into the form $\text{Div}_{t,y} T = 0$ where the symmetric tensor T is a non-local function of the macroscopic density ρ . This tensor turns out to be positive whenever the force field is *repulsive*. This leads us to a new space-time estimate, for instance for plasma flows (though not for galaxies). As in gas dynamics, this estimates displays a higher integrability property, by comparison with the mass and energy estimates, to the price of a time integration. Remark that our argument suggests that the same strategy could be tried for other physically relevant divergence-free (or divergence-controlled) symmetric tensors T that are not positive semi-definite : find another symmetric tensor T' with the same row-wise divergence, and see if this T' fulfills the positivity requirement.

2 Compensated integrability

2.1 Periodic case

Let us recall Theorem 2.1 of [8] :

Theorem 2.1 *Let the DPT $x \mapsto A(x)$ be Γ -periodic, with $A \in L^1(\mathbb{R}^d/\Gamma)$. Then $(\det A)^{\frac{1}{d-1}} \in L^1(\mathbb{R}^d/\Gamma)$ and there holds*

$$(1) \quad \int_{\mathbb{R}^d/\Gamma} (\det A(x))^{\frac{1}{d-1}} dx \leq \left(\det \int_{\mathbb{R}^d/\Gamma} A(x) dx \right)^{\frac{1}{d-1}}.$$

The fact that $(\det A)^{\frac{1}{d-1}}$ is integrable, is not implied by the sole assumption $A \in L^1(\mathbb{R}^d/\Gamma)$. There is no reason why other homogeneous functions of A of degree $\frac{d}{d-1}$ should be integrable. The theorem therefore expresses a compensation property, which may be called *Compensated Integrability*.

A similar phenomenon was established by Coifman & al. [1], who proved that if two vector fields $u, v \in L^2(\mathbb{R}^d)$ satisfy $\text{div} u = 0$ and $\text{curl} v = 0$, then the scalar product $u \cdot v$ belongs to the Hardy space $\mathcal{H}^1(\mathbb{R}^d)$, a strict subspace of the obvious $L^1(\mathbb{R}^d)$. If $f = u \cdot v$ is non-negative, this implies that $f \log(1 + f) \in L^1_{\text{loc}}(\mathbb{R}^d)$. It is interesting to compare the result of Coifman & al. with ours when $d = 2$, because their differential constraints read $\text{Div} T = 0$ for $T = \begin{pmatrix} u_1 & u_2 \\ -v_2 & v_1 \end{pmatrix}$, the function $f = u \cdot v$

¹The sign determines whether the force is attractive (gravity) or repulsive (electric force).

is nothing but $\det T$, and we both make the assumption that $f \geq 0$. Their integrability is higher than ours, both in the assumption and in the conclusion. We have an extra assumption (symmetry), but our gain is better (from $L^{1/2}$ to L^1 , instead of from L^1 to $L \log L$).

The following proof of Theorem 2.1 is much more direct than the one in [8]. Let $f > 0$ be an arbitrary smooth, Γ -periodic function. Let $S \in \mathbf{SPD}_d$ satisfy the constraint

$$(2) \quad \det S = \int_{\mathbb{R}^d/\Gamma} f(x) dx.$$

By Yan Yan Li's Theorem [6], the Monge–Ampère equation $\det D^2\theta = f$ admits a smooth, convex solution of the form $\theta = \frac{1}{2}x^T Sx + \rho(x)$ where ρ is periodic too. With the same arguments as in [8], we have

$$(f \det A)^{\frac{1}{d}} \leq \frac{1}{d} \operatorname{Tr}(AD^2\theta) = \frac{1}{d} (\operatorname{Tr}(AS) + \operatorname{div}(A\nabla\rho)).$$

Integrating, we have

$$\int_{\mathbb{R}^d/\Gamma} (f \det A)^{\frac{1}{d}} dx \leq \frac{1}{d} \operatorname{Tr}(S \int_{\mathbb{R}^d/\Gamma} A(x) dx) =: \frac{1}{d} \operatorname{Tr}(SA_+).$$

We optimize the choice of S by taking $S = \lambda A_+^{-1}$, where $\lambda > 0$ satisfies (2). We obtain

$$\int_{\mathbb{R}^d/\Gamma} (f \det A)^{\frac{1}{d}} dx \leq \lambda = \left(\int_{\mathbb{R}^d/\Gamma} f(x) dx \det A_+ \right)^{\frac{1}{d}}.$$

Denoting $\phi := f^{\frac{1}{d}}$, this is recast as

$$\int_{\mathbb{R}^d/\Gamma} \phi(\det A)^{\frac{1}{d}} dx \leq \lambda = \|\phi\|_{L_{\text{per}}^d} (\det A_+)^{\frac{1}{d}}.$$

By density, this inequality remains true for every non-negative function $\phi \in L_{\text{per}}^p$. Because $\det A \geq 0$, we infer that $(\det A)^{\frac{1}{d}} \in L_{\text{per}}^{d'}$, where $d' = \frac{d}{d-1}$ is the conjugate exponent, that is $(\det A)^{\frac{1}{d-1}} \in L_{\text{per}}^1$, and we have

$$\|(\det A)^{\frac{1}{d}}\|_{L_{\text{per}}^{d'}} \leq (\det A_+)^{\frac{1}{d}}.$$

This is exactly our functional inequality (1).

Remark. The proof given here has the advantage to apply under a weaker hypothesis, when we assume that the entries of A are bounded measures on \mathbb{R}^d/Γ , instead of being integrable. The positive-ness means that for every $\xi \in S^{d-1}$, the bounded measure $\sum_{i,j} \xi_i \xi_j a_{ij}$ is non-negative. The only point to clarify is the definition of $(\det A)^{\frac{1}{d}}$. By assumption, the entries are absolutely continuous with respect to the non-negative measure $\operatorname{Tr} A$. One can therefore introduce the densities f_{ij} by $a_{ij} = f_{ij} \operatorname{Tr} A$.

These functions f_{ij} are integrable with respect to $\text{Tr} A$ and the matrix $F = (f_{ij})_{1 \leq i, j \leq d}$ takes values in \mathbf{Sym}_d^+ . Then

$$(\det A)^{\frac{1}{d}} := (\det F)^{\frac{1}{d}} \text{Tr} A$$

defines a bounded measure. Once again, there is a compensated integrability: $(\det A)^{\frac{1}{d}}$ is absolutely continuous with respect to the Lebesgue measure, and its density belongs to $L^{\frac{d}{d-1}}(\mathbb{R}^d/\Gamma)$.

The present remark can be made for the Theorems 2.2 and 2.3 below. It is also meaningful when treating renormalized solutions of Boltzman flow (see Section 3.4 of [8]), where the conservation of momentum includes a *defect measure* Σ , since our analysis provided an estimate of

$$\int_0^\tau \int_{\mathbb{R}^n} (\rho \det \Sigma)^{\frac{1}{n}} dy dt.$$

2.2 DPTs in a bounded domain

The analysis above works the same in a convex bounded domain Ω . For the sake of completeness, we may assume that $\text{Div} A$ is a vector-valued bounded measure. Let $f > 0$ be an arbitrary smooth function over $\bar{\Omega}$. Let $\nabla \theta$ be the Brenier's transport from $(\Omega, f(x)dx)$ to (B_r, dx) , where the constraint

$$\frac{r^d}{d} |S^{d-1}| = \int_{\Omega} f(x) dx$$

expresses that both measures have the same mass.

Following the same lines, we have

$$(f \det A)^{\frac{1}{d}} \leq \frac{1}{d} (\text{div}(A \nabla \theta) + (\text{Div} A) \nabla \theta).$$

Integrating, there comes

$$\int_{\Omega} (f \det A)^{\frac{1}{d}} dx \leq \frac{1}{d} \left(\int_{\partial \Omega} (A \nabla \theta) \cdot \vec{n} ds(x) + \int_{\Omega} (\text{Div} A) \nabla \theta dx \right) \leq \frac{r}{d} \left(\int_{\partial \Omega} |A \vec{n}| ds(x) + \|\text{Div} A\|_{\mathcal{M}} \right).$$

Let us proceed as above: we introduce $\phi = f^{\frac{1}{d}}$ and rewrite

$$\int_{\Omega} \phi (\det A)^{\frac{1}{d}} dx \leq K \|\phi\|_{L^d}$$

where

$$K = \frac{1}{d} \left(\frac{d}{|S^{d-1}|} \right)^{\frac{1}{d}} \left(\int_{\partial \Omega} |A \vec{n}| ds(x) + \|\text{Div} A\|_{\mathcal{M}} \right).$$

We deduce that $(\det A)^{\frac{1}{d}} \in L^{d'}(\Omega)$, with $\|(\det A)^{\frac{1}{d}}\|_{L^{d'}} \leq K$. This proves that $(\det A)^{\frac{1}{d-1}}$ is integrable, a point that was left open in Theorem 2.3 of [8].

2.2.1 Non-convex domains

This Theorem 2.3 can actually be improved in another direction, by dropping the assumption that the domain be convex. Our ultimate result is therefore:

Theorem 2.2 *Let Ω be a bounded open subset in \mathbb{R}^d with smooth boundary. Let $A \in L^1(\Omega; \mathbf{Sym}_d^+)$ be given, such that $\text{Div} A$ is a (vector-valued) bounded measure. We assume that the normal trace $A\vec{n}$ is a bounded measure. Then there holds*

$$(3) \quad \int_{\Omega} (\det A(x))^{\frac{1}{d-1}} dx \leq \frac{1}{d|\mathcal{S}^{d-1}|^{\frac{1}{d-1}}} \left(\|A\vec{n}\|_{\mathcal{M}(\partial\Omega)} + \|\text{Div} A\|_{\mathcal{M}(\Omega)} \right)^{\frac{d}{d-1}}.$$

Proof

The Theorem has already been proven when the domain is convex and the normal trace is integrable. We may therefore apply this restricted version to the pair $(\tilde{\Omega}, \tilde{A})$, where $\tilde{\Omega}$ is a ball that contains $\overline{\Omega}$, and \tilde{A} is the extension of A by 0_d over $\tilde{\Omega} \setminus \Omega$. The assumption tells us that $\text{Div} \tilde{\Omega}$ is a bounded measure, and we have

$$\|\text{Div} \tilde{A}\|_{\mathcal{M}(\tilde{\Omega})} = \|A\vec{n}\|_{\mathcal{M}(\partial\Omega)} + \|\text{Div} A\|_{\mathcal{M}(\Omega)}.$$

Besides, the normal trace of \tilde{A} vanishes identically. The inequality (3), valid for \tilde{A} in $\tilde{\Omega}$, yields the same inequality for A in Ω . ■

Of course, the same remark as in the periodic case holds true: we can actually replace the integrability of A , by the assumption that A is a symmetric tensor of bounded measures, non-negative in the sense that for every vector $\xi \in \mathbb{R}^d$, the scalar measure $\xi^T A \xi$ is non-negative. This assumption allows us to define a non-negative measure $(\det A)^{\frac{1}{d}}$, and the theorem tells us that the latter is actually a function of class $L^{\frac{d}{d-1}}$. In particular, it does not display a singular part.

Remark also that the isoperimetric inequality follows immediately and for every smooth domain, by applying Theorem 2.2 to the tensor $A \equiv I_d$.

2.3 The case of a slab

We consider now a domain of the form $\Omega = (0, \tau) \times \mathbb{R}^n$, where $\tau > 0$ and $d = 1 + n$. We split the coordinates as $x = (t, y)$ and may think of t and y as time and space variables, respectively. Suppose that A is a DPT over Ω , with $A \in L^1(\Omega)$. Assume also that its normal traces on the top/bottom boundaries are bounded measures.

Let us choose a function $\phi \in \mathcal{D}(\mathbb{R}^n)$ with support in the ball B_2 of radius 2, which satisfies $0 \leq \phi \leq 1$ everywhere, and $\phi \equiv 1$ in the unit ball B_1 . If $R > 0$, define $\phi_R(y) := \phi(y/R)$. Let us apply Theorem

2.2 to the tensor $\phi_R A$, in the domain $\Omega_R = (0, \tau) \times B_{2R}$. Its divergence is $A \nabla \phi_R$ and its trace on the lateral boundary $(0, \tau) \times \partial B_{2R}$ vanishes, while $|\phi_R A \xi| \leq |A \xi|$. We have therefore

$$\begin{aligned} \int_{\Omega_R} (\phi_R \det A(x))^{\frac{1}{d-1}} dx &\leq \frac{1}{d |S^{d-1}|^{\frac{1}{d-1}}} \left(\|\phi_R A \vec{n}\|_{\mathcal{M}(\partial \Omega_R)} + \|\text{Div}(\phi_R A)\|_{\mathcal{M}(\Omega_R)} \right)^{\frac{d}{d-1}} \\ &\leq \frac{1}{d |S^{d-1}|^{\frac{1}{d-1}}} \left(\|A \vec{n}\|_{\mathcal{M}(\{0\} \times \mathbb{R}^n)} + \|A \vec{n}\|_{\mathcal{M}(\{\tau\} \times \mathbb{R}^n)} + \|A \nabla \phi_R\|_{\mathcal{M}(\Omega_R)} \right)^{\frac{d}{d-1}}. \end{aligned}$$

We infer

$$\int_0^\tau \int_{B_R} (\det A)^{\frac{1}{d-1}} dy dt \leq \frac{1}{d |S^{d-1}|^{\frac{1}{d-1}}} \left(\|A \vec{n}\|_{\mathcal{M}(\{0\} \times \mathbb{R}^n)} + \|A \vec{n}\|_{\mathcal{M}(\{\tau\} \times \mathbb{R}^n)} + \frac{c}{R} \|A\|_{L^1} \right)^{\frac{d}{d-1}}.$$

Passing to the limit as $R \rightarrow +\infty$, we may state:

Theorem 2.3 *Let A be a DPT over the slab $(0, \tau) \times \mathbb{R}^n$. We assume that it is integrable, and that its normal traces on the top/bottom boundaries are bounded measures. Then $(\det A)^{\frac{1}{n}}$ is integrable and one has*

$$(4) \quad \int_0^\tau \int_{\mathbb{R}^n} (\det A)^{\frac{1}{n}} dy dt \leq \frac{1}{(n+1) |S^n|^{\frac{1}{n}}} \left(\|A \vec{n}\|_{\mathcal{M}(\{0\} \times \mathbb{R}^n)} + \|A \vec{n}\|_{\mathcal{M}(\{\tau\} \times \mathbb{R}^n)} \right)^{1+\frac{1}{n}}.$$

This is the fundamental inequality that we apply to various models in physics, mostly in gas dynamics.

3 Singularities vs gain of integrability

This section intends to show that the determinant is essentially the only homogeneous polynomial over \mathbf{Sym}_d^+ to display a maximal gain of integrability when applied to DPTs.

3.1 Homogeneous singularities of DPTs

We begin by considering those DPTs that are homogeneous, with a negative degree $-m$; we call m the *order of singularity*. Denoting $r = |x|$ and $e = \frac{x}{r}$, we look for tensors of the form

$$T(x) = r^{-m} S(e)$$

where $e \mapsto S(e)$ takes values in \mathbf{Sym}_d^+ . Of course, we restrict our attention to integrable singularities, more generally to singularities that are bounded measures. We assume therefore that $m < d$ and S is a bounded measure over the unit sphere S^{d-1} .

We begin with elementary examples:

Lemma 3.1 *When $m \leq d - 1$, the tensor*

$$T_m = \frac{1}{r^m} (m e \otimes e + (d - 1 - m) I_d)$$

is an integrable DPT.

The proof is elementary, using the identity $\text{Div}(fS) = f \text{Div} S + S \nabla f$ and the fact $\text{Div}(x \otimes x) = (d + 1)x$. The family T_m includes the special cases

$$T_0 = (d - 1)I_d, \quad T_{d-1} = \frac{d - 1}{r^{d-1}} e \otimes e,$$

where the constant factors $d - 1$ are irrelevant. Remark that if $m \in (d - 1, d)$, the divergence-free tensor T_m is integrable but not positive. This is consistent with the first part of the following statement.

Proposition 3.1 *1. In the class of homogeneous DPTs, the order of singularity $m < d$ must be less than or equal to $d - 1$.*

2. The homogeneous DPTs with order of singularity $d - 1$ are of the form

$$(5) \quad T(x) = \frac{\lambda(e)}{r^{d-1}} e \otimes e$$

for some non-negative measure λ . For every such measure, the formula above provides a DPT whose entries are bounded measures.

3. If $\theta : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex and positively homogeneous of degree one (for instance if θ is a semi-norm), then $\widehat{D^2\theta}$ is a homogeneous DPT of maximal order of singularity, that is $d - 1$.

Remark. A partial converse of point 3 above holds true: suppose T is in the form (5) for some measure $\lambda \geq 0$. Assume in addition that

$$(6) \quad \int_{S^{d-1}} e_i \lambda(e) = 0, \quad \forall i = 1, \dots, d.$$

Then there exists a convex function θ , positively homogeneous of degree one, such that $T = \widehat{D^2\theta}$. The map $e \mapsto \nabla\theta(e)$ solves Minkowski's Problem, which consists in finding a convex hypersurface whose Gauß curvature is a prescribed function of the normal direction, here $\kappa = \frac{1}{\lambda(e)}$. The necessary condition (6) turns out to be sufficient, see Pogorelov [7]. The solution of the problem is related to an equation of Monge-Ampère type on the sphere, namely

$$\det D^2\theta|_{e^\perp} = \lambda(e),$$

where the convexity of θ as a function of $x \in \mathbb{R}^d$ ensures the ellipticity. If λ is even, the condition (6) is obviously satisfied and the solution θ is a semi-norm.

Proof

Let T be a homogeneous DPT of order $m < d$. The divergence-free condition writes

$$\text{Div } S = \frac{m}{r^2} Sx,$$

where we view S as a function of x , homogeneous of degree zero. Let us make the scalar product with x ,

$$me^T Se = x \cdot \text{Div } S = \text{div}(Sx) - \text{Tr } S.$$

Let us integrate over the unit ball B ,

$$\int_B (me^T Se + \text{Tr } S) dx = \int_{S^{d-1}} e^T Se ds(e).$$

With $dx = r^{d-1} dr ds(e)$, this yields

$$\int_{S^{d-1}} ((m+1-d)e^T Se + \text{Tr } S|_{e^\perp}) ds(e) = 0.$$

When $m \geq d-1$, the integrand is non-negative (recall that $S(e) \in \mathbf{Sym}_d^+$) and the equality above is equivalent to $(m+1-d)e^T Se + \text{Tr } S|_{e^\perp} \equiv 0$. If $m > d-1$, this means $S \equiv 0_d$, which proves the first item. If $m = d-1$, it says only that S vanishes over e^\perp , that is S is of the form $\lambda(e)e \otimes e$ with $\lambda(e) \geq 0$.

Conversely, let λ be a non-negative measure, so that $T := r^{1-d}\lambda(e)e \otimes e$ is a symmetric, positive semi-definite tensor. If $\lambda \equiv 1$, we verify easily that the corresponding tensor T^0 equals $\widehat{D^2\theta}$ where $\theta \equiv |x|$. Therefore T^0 is a DPT. For a general λ , we have

$$\text{Div } T = \text{Div}(\lambda T^0) = T^0 \nabla \lambda = r^{-d}(x \cdot \nabla \lambda)e \equiv 0,$$

where the last identity is that of Euler for homogeneous functions. Thus T is a DPT.

Finally, if θ is convex and positively homogeneous of degree one, then we already know that $T = \widehat{D^2\theta}$ is a DPT. Since $\widehat{D^2\theta}$ is homogeneous of degree -1 , T is homogeneous of degree $1-d$. ■

3.2 Are there other gains of integrability ?

Definition 3.1 Let $f : \mathbf{Sym}_d^+ \rightarrow \mathbb{R}^+$ be a continuous, homogeneous function of degree $\frac{d}{d-1}$. We say that f exhibits a maximal gain of regularity for DPTs if there exists a finite constant c_f such that for every divergence-free $T \in L^1(B; \mathbf{Sym}_d^+)$ with an integrable normal trace, one has

$$(7) \quad \int_B f(T) dx \leq c_f \|T\vec{n}\|_{L^1(S^{d-1})}^{\frac{d}{d-1}}.$$

The degree of homogeneity is justified by the fact that if *all* the derivatives ∂T are under control, and not only $\text{Div } T$, then $T \in W^{1,1}(B)$, which is contained in $L^{\frac{d}{d-1}}(B)$ by Sobolev embedding. Besides, a dimensional analysis shows that such an inequality would be valid in every ball of \mathbb{R}^d , with the same constant c_f .

We begin by testing (7) against the tensors T_m . Because of $\vec{n} = e$, we have $T_m \vec{n} = (d-1)e$ and thus $\|T_m \vec{n}\|_{L^1(S^{d-1})} = (d-1)|S^{d-1}|$. On the other hand,

$$\int_B f(T_m) dx = \frac{d-1}{d(d-1-m)} \int_{S^{d-1}} f(me \otimes e + (d-1-m)I_d) ds(e).$$

We deduce therefore

Lemma 3.2 *If f displays a maximal gain of integrability, then there exists a finite constant c'_f such that*

$$(8) \quad \int_{S^{d-1}} f(me \otimes e + (d-1-m)I_d) ds(e) \leq c'_f(d-1-m)$$

for every $m < d-1$.

By letting $m \rightarrow d-1$, f must vanish on the cone of rank-one tensors $v \otimes v$.

3.3 The case of immanants

The determinant is not the only homogeneous polynomial to vanish on the cone of rank-one tensors. Most of the immanants² share this property. We recall their definition:

Definition 3.2 *Let G be a subgroup of the symmetric group \mathfrak{S}_d and χ be an irreducible character over G . The immanant $J_\chi^G : \mathbf{M}_d(\mathbb{R}) \rightarrow \mathbb{R}$ is the polynomial*

$$J_\chi^G(M) = \sum_{g \in G} \chi(g) \prod_{i=1}^d m_{ig(i)}.$$

If $G = \mathfrak{S}_d$, an immanant has integral coefficients. Two examples of immanants are the determinant, for which χ is the signature, and the *permanent*, denoted Per , where $\chi \equiv 1$. When $G = (1)$, there is only one immanant, the diagonal product.

When G is a proper subgroup, and if there exists a $g \in G$ such that g^{-1} and g are in different conjugacy classes, an immanant may have non-real coefficients. However, the restriction of an immanant to \mathbf{Sym}_d does have real coefficients; the reason is that $\chi(g^{-1}) = \overline{\chi(g)}$, while for symmetric matrices

$$\prod_{i=1}^d s_{ig^{-1}(i)} = \prod_{i=1}^d s_{ig(i)}.$$

²There are two spellings in the literature. *Immanant* recalls the determinant, while *immanent* recalls the permanent.

If $v \in \mathbb{R}^n$, we have

$$J_\chi^G(v \otimes v) = \left(\sum_{g \in G} \chi(g) \right) v_1^2 \cdots v_d^2.$$

For $\chi \neq 1$, the sum above equals zero by orthogonality with the trivial character ; therefore J_χ^G vanishes identically over the cone of tensors $v \otimes v$.

Let us recall Schur's property of immanants over the cone of positive semi-definite matrices. We use the digit 1 to denote the unit element (the identity) of G .

Proposition 3.2 (I. Schur, 1918.) *For every $S \in \mathbf{Sym}_d^+$, one has*

$$\det S \leq \frac{1}{\chi(1)} J_\chi^G(S).$$

The particular case when $G = (1)$,

$$\det S \leq s_{11} \cdots s_{dd}, \quad \forall S \in \mathbf{Sym}_d^+,$$

is known as the Hadamard inequality. The *permanent dominance conjecture* tells on the contrary that

$$J_\chi^G(S) \leq \text{Per}(S), \quad \forall S \in \mathbf{Sym}_d^+.$$

Schur's property allows us to define

$$f_\chi^G(S) = \left(J_\chi^G(S) \right)^{\frac{1}{d-1}},$$

which is homogeneous of degree $\frac{d}{d-1}$ over \mathbf{Sym}_d^+ . A natural question is whether f_χ^G can display a maximal gain of integrability. An obvious necessary condition is $\chi \neq 1$.

To begin with, we examine the degree of the polynomial $p_\chi^G(X) := J_\chi^G(I_d + X e \otimes e)$.

Lemma 3.3 *The degree of p_χ^G is ≥ 2 , unless $G = \mathfrak{S}_n$ and $\chi = \varepsilon$.*

Remark that this degree is 1 for the pair $(\mathfrak{S}_d, \varepsilon)$.

Proof

The coefficient of X^2 in p_χ^G ,

$$h_\chi^G(e) := \chi(1) \sum_{i < j} e_i^2 e_j^2 + \sum_{\tau = (ij) \in G} \chi(\tau) e_i^2 e_j^2,$$

is itself a polynomial in e , a linear combination of monomials $e_i^2 e_j^2$. The coefficient of a monomial is either $\chi(1)$ or $\chi(1) + \chi(\tau)$ if $\tau = (ij) \in G$ (notice that $\chi(\tau)$ is real because $\tau^{-1} = \tau$).

Let $\rho : G \rightarrow GL(V)$ denote the representation associated with χ . We recall that $|\chi(g)| \leq \chi(1)$ for every $g \in G$, and the equality implies that $\rho(g)$ is a homothety, $\rho(g) = \frac{\chi(g)}{\chi(1)} \text{id}_V$. In particular h_χ^G has non-negative coefficients, from which we deduce $h_\chi^G \geq 0$.

If $\deg p_\chi^G \leq 1$, we have $h_\chi^G \equiv 0$. Since $\chi(1) > 0$, this implies that G contains every transposition, and therefore $G = \mathfrak{S}_d$. It also tells that $\chi(\tau) = -\chi(1)$ for every transposition, which implies $\rho(\tau) = -\text{id}_V$. Because ρ is a morphism and $\varepsilon(g) = (-1)^\ell$ where ℓ is the number of transposition in a factorisation of g , we deduce $\rho(g) = \varepsilon(g)\text{id}_V$. Hence $\chi = \varepsilon$. ■

This yields the main result of this section.

Theorem 3.1 *Among the pairs (G, χ) , only one yields a function f_χ^G that displays a maximal gain of integrability, namely the pair $(\mathfrak{S}_d, \varepsilon)$, for which the function is $\det^{\frac{1}{d-1}}$.*

Proof

Denote $h_k(e)X^k$ the dominant monomial in p_χ^G . When $m \rightarrow (d-1)^-$, we have

$$J_\chi^G(me \otimes e + (d-1-m)I_d) \sim (d-1-m)^{d-k} m^k h_k(e).$$

In the formula

$$h_k(e) = \sum_{|A|=k} e_A^2 \sum_{g \in G | A^c \subset \text{Fix}(g)} \chi(g), \quad e_A = \prod_{i \in A} e_i,$$

each sum of characters is a non-negative integer. For the sake of completeness, it equals the multiplicity of the trivial representation in $\text{Res}_H^G(\rho)$, where H is the subgroup of all $g \in G$ such that $A^c \subset \text{Fix}(g)$ (the stabilizer of A^c).

By definition $h_k \not\equiv 0$ and we obtain $\int_{S^{d-1}} (h_k(e))^{\frac{1}{d-1}} ds(e) > 0$. Therefore

$$\int_{S^{d-1}} f_\chi^G(me \otimes e + (d-1-m)I_d) ds(e) \sim \mu(d-1-m)^{\frac{d-k}{d-1}}$$

as $m \rightarrow (d-1)^-$, where μ is some positive constant. If f_χ^G displays a maximal gain of integrability, this together with Lemma 3.2 implies $\frac{d-k}{d-1} \leq 1$, that is $k \leq 1$.

We conclude by using Lemma 3.3. ■

4 More examples of divergence-free symmetric tensors

4.1 The wave equation

Let u be a solution of the wave equation :

$$\partial_t^2 u = c^2 \Delta_y u$$

The assumption of bounded energy yields the standard regularity $\nabla_{t,y}u \in L^\infty(\mathbb{R}; L^2(\mathbb{R}^n))$. Then the tensor

$$T = \begin{pmatrix} \frac{1}{2}(u_t^2 + c^2|\nabla u|^2) & -c^2u_t\nabla u^T \\ -c^2u_t\nabla u & c^4\nabla u \otimes \nabla u + \frac{c^2}{2}(u_t^2 - c^2|\nabla u|^2)I_n \end{pmatrix}.$$

is integrable in slabs $(0, \tau) \times \mathbb{R}^n$ and satisfies $\text{Div}_{t,y}T = 0$.

The matrix $T(t, y)$ admits two invariant subspaces, namely the plane $\mathbb{R} \times \mathbb{R}\nabla u$ and its orthogonal. The restriction of T to $\mathbb{R} \times \mathbb{R}\nabla u$ is positive definite. In particular, T is a DPT when $n = 1$. On the contrary, the restriction to $\{0\} \times \nabla u^\perp$ is a homothety of ratio $\frac{c^2}{2}(u_t^2 - c^2|\nabla u|^2)$. When $n \geq 2$, this tensor is therefore positive semi-definite if and only if $c|\nabla u| \leq |u_t|$. One has

$$\det T = c^{2n} \left(\frac{u_t^2 - c^2|\nabla u|^2}{2} \right)^{n+1}.$$

Notice that this determinant has a constant (positive) sign when the space dimension is odd, even if T is not positive semi-definite.

If the compensated integrability held true, no matter of the signature of the tensor, we should receive the estimate

$$\int_{\mathbb{R}} \int_{\mathbb{R}^n} |u_t^2 - c^2|\nabla u|^2|^{1+\frac{1}{n}} dy dt \leq CE_0^{1+\frac{1}{n}},$$

where

$$E_0 \equiv \int_{\mathbb{R}^n} \frac{1}{2}(u_t^2 + c^2|\nabla u|^2) dy$$

is the total energy. If $n = 1$, this estimate is correct since T is a DPT, and the proof is actually an application of Fubini's Theorem to Riemann invariants. On the contrary, in space dimension $n = 2$ or 3, Klainerman & Machedon constructed an example where the energy is finite, but $u_t^2 - c^2|\nabla u|^2 \notin L^{1+\frac{1}{n}}_{t,y}$. See Proposition 5 in [5]. Therefore the compensated integrability may fail when we drop the positivity assumption.

Remark. If we consider the Laplace equation instead, $\Delta_x u = 0$, the symmetric tensor $T = \nabla u \otimes \nabla u - \frac{1}{2}|\nabla u|^2 I_d$ is again divergence-free. Still, it is neither positive nor negative and thus we may not apply the theorems above. However, the gain of integrability holds true. Of course, it is proved from a different argument. The normal trace

$$T\vec{n} = \frac{\partial u}{\partial n} \nabla u - \frac{1}{2}|\nabla u|^2 \vec{n}$$

satisfies $|T\vec{n}| = \frac{1}{4}|\nabla u|^2$. The assumption that $T\vec{n}$ is a bounded measure over $\partial\Omega$ tells us that the normal derivative belongs to $L^2(\partial\Omega)$. By regularity theory, we obtain $u \in H^{3/2}(\Omega)$. Then the Sobolev embedding provides $\nabla u \in L^p(\Omega)$ with $p = \frac{2d}{d-1}$. Therefore

$$|\det T|^{\frac{1}{d-1}} = \left(\frac{1}{2}|\nabla u|^2 \right)^{\frac{d}{d-1}} \in L^1(\Omega).$$

4.2 The nonlinear Maxwell system

For the sake of simplicity, we choose a system of physical units in which the speed of light is $c = 1$.

The variational formulation of the Maxwell's system for the electro-magnetic field in vacuum is

$$\delta\mathcal{L} = 0, \quad \text{where } \mathcal{L} := \int \int L(B, E) dy dt.$$

The ambient space is that of closed 2-forms $\omega := E \cdot (dt \wedge dy) + B \cdot (dy \wedge dy)$ on the 1 + 3-dimensional Minkowski space with coordinates (t, y) . The closedness reads

$$\partial_t B + \text{curl} E = 0, \quad \text{div} B = 0$$

while the variational principle yields

$$\partial_t D - \text{curl} H = 0, \quad \text{div} D = 0 \quad \text{where } D := \frac{\partial L}{\partial E}, \quad H := -\frac{\partial L}{\partial B}.$$

Assuming that $(B, E) \mapsto (B, D)$ is a change of variable, we may define the energy density

$$W(B, D) := D \cdot E - L(B, E).$$

Then we have

$$(9) \quad E = \frac{\partial W}{\partial D}, \quad H := \frac{\partial W}{\partial B}.$$

The relations (9) were the starting point of the thermodynamical approach for non-linear models by Coleman & Dill [2]. Their general form of the Maxwell's system,

$$\partial_t B + \text{curl} \frac{\partial W}{\partial D} = 0, \quad \partial_t D - \text{curl} \frac{\partial W}{\partial B} = 0, \quad \text{div} B = \text{div} = 0$$

implies four extra conservation laws

$$(10) \quad \partial_t W + \text{div}(E \times H) = 0,$$

$$(11) \quad \partial_t(D \times B) - \text{Div}(E \otimes D + H \otimes B) + \nabla(D \cdot E + B \cdot H - W) = 0.$$

The first equation of the system above justifies the terminology that $Q := E \times H$ is the *electromagnetic momentum*. The vector field $P := D \times B$ is called the Poynting vector.

When the Maxwell's equations are consistent with special relativity, the Lagrangian \mathcal{L} must be invariant under the action of the Lorentz group over 2-forms. This means that the density L depends only on two scalar variables (σ, π) ,

$$L(B, E) = \ell(\sigma, \pi), \quad \sigma := \frac{1}{2}(|B|^2 - |E|^2), \quad \pi := B \cdot E.$$

We have therefore

$$D = -\ell_\sigma E + \ell_\pi B, \quad H = -\ell_\sigma B - \ell_\pi E$$

and this implies

$$(12) \quad P = Q.$$

This fundamental identity tells that the Abraham form $E \times H$ of the Poynting vector coincides with its Minkowski form $D \times B$.

Now, the system (10,11) can be rewritten in the form

$$\text{Div}_{t,y} T = 0$$

where T is a symmetric tensor, thanks to (12) :

$$T = \begin{pmatrix} W & P^T \\ P & \ell_\sigma(E \otimes E + B \otimes B) + (\ell + B \cdot H)I_3 \end{pmatrix}.$$

Decomposition of T . When (B, E) are linearly independent, the 4×4 matrix T admits two obvious invariant planes, namely $\mathbb{R} \times \mathbb{R}P$ and its orthogonal $\{0\} \times \text{Span}(B, E)$. We therefore have $T \sim S \oplus^\perp R$, where

$$S = \begin{pmatrix} W & |P|^2 \\ 1 & \ell + B \cdot H \end{pmatrix}, \quad R = \begin{pmatrix} \ell - \pi \ell_\pi & \pi \ell_\sigma \\ \pi \ell_\sigma & \ell - 2\sigma \ell_\sigma - \pi \ell_\pi \end{pmatrix}.$$

The spectrum of T , which is real, is therefore the union of that of S and R . We have

$$\det S = -\det R = \ell_\sigma^2(\sigma^2 + \pi^2) - (\ell - \sigma \ell_\sigma - \pi \ell_\pi)^2.$$

Thus $\det T = -(\det S)^2$ and T cannot be positive semi-definite. It cannot be negative either because in practice W is positive.

In conclusion, we cannot conclude a compensated integrability for the tensor T .

4.3 Systems of conservation laws with a convex extension

The space-time dimension is still $d = 1 + n$, while the field u takes values in a convex domain of \mathbb{R}^N . We consider a system of the form

$$\partial_t u + \sum_\alpha \partial_\alpha f^\alpha(u) = 0$$

and assume that it is formally compatible with an additional conservation law

$$\partial_t \eta(u) + \text{div}_x \vec{Q}(u) = 0$$

where $D^2\eta > 0_N$ (strong convexity). Following Godunov [4], we introduce the conjugate variables $q_j = \partial\eta/\partial u_j$ and we know that there exist potentials L^α ($\alpha = 0, \dots, n$) such that $u = \partial L^0/\partial q$ and $f^\alpha(u) = \partial L^\alpha/\partial q$. The system rewrites

$$\partial_t \frac{\partial L^0}{\partial q} + \sum_{\alpha} \partial_{q^\alpha} \frac{\partial L^\alpha}{\partial q} = 0.$$

The entropy and its fluxes are given by

$$\eta(u) = q \cdot u - L^0(q), \quad Q^\alpha(u) = q \cdot \frac{\partial L^\alpha}{\partial q} - L^\alpha(q).$$

Let us assume $N = n + 1 =: d$. We label the indices j from 0 to n . Then $\text{Div}_{t,y} T = 0$ where $T = \partial L/\partial q$ is a Jacobian matrix. It is symmetric if and only if L derives from a single potential $R(q)$:

$$L^\alpha = \frac{\partial R}{\partial q^\alpha}, \quad T = D_q^2 R.$$

It is positive if and only if $q \mapsto R(q)$ is a convex function. When it is so, Theorem 2.3 yields an estimate of

$$\int_0^\tau \int_{\mathbb{R}^n} (\det D_q^2 R)^{\frac{1}{n}} dx dt$$

in terms of the L^1 -norm of the $\frac{\partial L^0}{\partial q}$ at initial and final times. In practical situations, these norms can be estimated in terms of the integral

$$I_0 := \int_{\mathbb{R}^n} \eta(u_0(y)) dy$$

at initial time, using the second principle of thermodynamics, which tells that

$$t \mapsto I(t) = \int_{\mathbb{R}^n} \eta(u(t,y)) dy$$

is a non-increasing function of time.

The above analysis applies also when $N > d$ and the d first lines of the system write $\text{Div}_{t,y} T = 0$ for a positive symmetric tensor. Then R depends upon the additional parameters (q_d, \dots, q_{N-1}) , which are not essential in the calculations.

Example: barotropic gas dynamics. In terms of the density ρ , the velocity v and the linear momentum $m = \rho v$, we have

$$u = \begin{pmatrix} \rho \\ m \end{pmatrix}, \quad f^\alpha(u) = \begin{pmatrix} m_\alpha \\ m_\alpha v + p(\rho) \bar{e}^\alpha \end{pmatrix}, \quad \eta(u) = \frac{|m|^2}{2\rho} + \varepsilon(\rho).$$

This yields $q = (-\frac{1}{2}|v|^2 + \varepsilon'(\rho), v)$ and

$$L^0(q) = \rho\varepsilon' - \varepsilon, \quad L^\alpha = (\rho\varepsilon' - \varepsilon)v_\alpha.$$

Of course, we find that L derives from a potential R , with

$$dR = (\rho\varepsilon' - \varepsilon)d(\varepsilon').$$

Let us denote $\lambda := \varepsilon'(\rho)$. Then $\rho\varepsilon' - \varepsilon = \varepsilon^*(\lambda)$, where ε^* denotes the convex conjugate of ε . We have $\rho = (\varepsilon^*)'(\lambda)$ and we find $R = R(\lambda)$ where $R' = \varepsilon^*$. With $\lambda = q_0 + \frac{1}{2}(q_1^2 + \dots + q_n^2)$, we find that the convexity of R in terms of q (the positivity of T) amounts to the convexity and monotonicity of R as a function of λ , because of

$$T = R''V \otimes V + R' \begin{pmatrix} 0 & 0 \\ 0 & I_n \end{pmatrix}, \quad V := \begin{pmatrix} 1 \\ q_1 \\ \vdots \\ q_n \end{pmatrix}.$$

To compute $\det T$, we observe that the plane $\text{vec}(\vec{e}^0, V)$ and its orthogonal are T -invariant. On the former, T acts as the matrix

$$M = \begin{pmatrix} 0 & -R' \\ R'' & |V|^2 R'' + R' \end{pmatrix},$$

while on the latter T acts as a homothety of ratio R' . Finally

$$\det T = (R')^{n-1} \det M = (R')^n R''.$$

Therefore we obtain an estimate of

$$\int_0^\tau \int_{\mathbb{R}^n} R'(\lambda) R''(\lambda)^{\frac{1}{n}} dx dt.$$

We point out that $R' = \varepsilon^* = \rho\varepsilon' - \varepsilon$ is the pressure, while $R'' = (\varepsilon^*)' = \rho$, since $(\varepsilon^*)'$ is the reciprocal of ε' .

5 Vlasov models

Vlasov models are kinetic equations for clouds where particles don't collide, but are driven by a self-induced force field $F(t, y)$. In a plasma, the particles are ions, subject to the Coulomb force. In galaxies, the particles are stars and interact through gravity. The Newton equations

$$\frac{dX}{dt} = v, \quad \frac{dv}{dt} = F$$

yield a kinetic model at the mesoscopic scale. The density $f(t, y, v)$ obeys to the equation

$$(\partial_t + v \cdot \nabla_y) f + F(t, y) \cdot \nabla_v f = 0.$$

The macroscopic density

$$\rho(t, y) := \int_{\mathbb{R}^n} f(t, y, v) dv$$

induces the force field:

$$F = -\nabla\phi, \quad \phi := \chi * \rho.$$

The radial kernel χ characterizes the physics at stake. The convolution is meant with respect to the space variable y ,

$$\phi(t, y) = \int_{\mathbb{R}^n} \chi(|y - z|) \rho(t, z) dz = \int_{\mathbb{R}^n} \chi(|z|) \rho(t, y - z) dz.$$

In Coulomb or gravity force, the potential ϕ is given by a Poisson equation, $\Delta\phi = \text{cst} \cdot \rho$, and we speak of the Vlasov–Poisson model. Depending upon the sign of the constant, the force is repulsive or attractive.

As usual, integration with respect to the velocity v yields mass conservation:

$$(13) \quad \partial_t \rho + \text{div}_y m = 0, \quad m := \int_{\mathbb{R}^n} f(t, y, v) v dv.$$

Integrating against $v dv$ gives formally

$$(14) \quad \partial_t m + \text{Div}_y \left(\int_{\mathbb{R}^n} f v \otimes v \right) - \rho F = 0,$$

which implies the conservation of the overall momentum, because of

$$\int_{\mathbb{R}^n} \rho F dy = - \int_{\mathbb{R}^n \times \mathbb{R}^n} \rho(y) \rho(z) \nabla \chi(y - z) dy dz,$$

and $\nabla \chi$ is odd.

The energy estimate. We have formally

$$\partial_t \int_{\mathbb{R}^n} \frac{|v|^2}{2} f dv + \text{div}_y \int_{\mathbb{R}^n} \frac{|v|^2}{2} f v dv = - \int_{\mathbb{R}^n} F \cdot \nabla_v f \frac{|v|^2}{2} dv = F \cdot m = -\nabla_y \phi \cdot m.$$

Therefore

$$\partial_t \int_{\mathbb{R}^n} \frac{|v|^2}{2} f dv + \text{div}_y \left(\int_{\mathbb{R}^n} \frac{|v|^2}{2} f v dv + \phi m \right) = \phi \text{div}_y m = -(\chi * \rho) \partial_t \rho.$$

Integrating in space, there comes

$$\frac{d}{dt} \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|v|^2}{2} f dv dy = - \int_{\mathbb{R}^n} (\chi * \rho) \partial_t \rho dy.$$

Because χ is even, the bilinear form

$$(\rho, \eta) \mapsto \int_{\mathbb{R}^n} (\chi * \rho) \eta \, dy$$

is symmetric. We infer

$$\frac{d}{dt} \int_{\mathbb{R}^n} (\chi * \rho) \rho \, dy = 2 \int_{\mathbb{R}^n} (\chi * \rho) \partial_t \rho \, dy,$$

and deduce formally

$$\frac{d}{dt} \left(\int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|v|^2}{2} f \, dv \, dy + \frac{1}{2} \int_{\mathbb{R}^n} (\chi * \rho) \rho \, dy \right) = 0.$$

The total energy

$$E(t) := \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|v|^2}{2} f \, dv \, dy + \frac{1}{2} \int_{\mathbb{R}^n} (\chi * \rho) \rho \, dy$$

is certainly a non-negative quantity if either χ , or its Fourier transform $\hat{\chi}$ (which is a real function because χ is even), is non-negative. Actually, because of the conservation of mass, it suffices that χ be bounded by below ($\inf \chi(r) > -\infty$), in order to build a non-negative conserved quantity of the form $E + \text{cst} \cdot M$. When this is the case, and the initial energy E_0 is finite, the conservation $E(t) \equiv E_0$ implies an *a priori* estimate. In the sequel, we consider solutions that satisfy the conservation of mass and momentum (equations 13 and 14), together with $E(t) \leq E_0$.

An alternate divergence-free tensor. Our next goal is two-fold. First we identify the expression $-\rho F$ as the space divergence of a symmetric tensor S , so that the following symmetric tensor be divergence-free,

$$T = \begin{pmatrix} \rho & m^T \\ m & \int_{\mathbb{R}^n} f v \otimes v \, dv + S \end{pmatrix}.$$

The second task is to determine those functions χ which guarantee that S is non-negative. When this property occurs, T is a DPT and we have the same estimate as for the Boltzmann equation, together with an estimate of

$$\int_0^\infty \int_{\mathbb{R}^n} (\rho \det S)^{\frac{1}{n}} \, dy \, dt$$

because of $\det T \geq \rho \det S$. Notice that, because $\rho \mapsto F$ is linear, we expect that S be quadratic in ρ . Therefore the new estimate involves a space-time integral of a quantity that is homogenous in ρ , of degree $2 + \frac{1}{n}$. This is a gain of integrability, compared to the energy estimate, which is only quadratic in ρ ; as usual, the price to pay is an extra integration, with respect to the time variable.

Dropping the time variable, we have

$$\begin{aligned} -(\rho F)(y) &= (\rho \nabla \phi)(y) = \rho(y) \int_{\mathbb{R}^n} \nabla \chi(z) \rho(y-z) \, dz = \rho(y) \int_{\mathbb{R}^n} \chi'(|z|) \rho(y-z) \frac{z}{|z|} \, dz \\ &= \frac{1}{2} \rho(y) \int_{\mathbb{R}^n} \chi'(|z|) (\rho(y-z) - \rho(y+z)) \frac{z}{|z|} \, dz \end{aligned}$$

At fixed z , we have

$$\rho(y)(\rho(y-z) - \rho(y+z)) = -\operatorname{div}_y \int_{-\frac{1}{2}}^{\frac{1}{2}} \rho(y + (s - \frac{1}{2})z) \rho(y + (s + \frac{1}{2})z) z ds.$$

We infer $-\rho F = \operatorname{Div}_y S$ with

$$S = -\frac{1}{2} \int_{\mathbb{R}^n} \frac{\chi'(|z|)}{|z|} dz \int_{-\frac{1}{2}}^{\frac{1}{2}} \rho(y + (s - \frac{1}{2})z) \rho(y + (s + \frac{1}{2})z) z \otimes z ds.$$

In particular, S is non-negative whenever χ is monotonous non-increasing. This corresponds to a repulsive force. This reminds us the case of the Euler equations, where the monotonicity of the pressure (the fact that the sound speed is a real number) induces a repulsive force too ; a gas tends to rarefy. Notice that the sign $\chi' \leq 0$ is also a necessary condition for the positivity ; think of a density ρ that concentrates at two points.

According to Section 2.3, a space-time estimate can be established whenever the DPT be integrable. For this we need that S be integrable. This comes in general as a consequence of the mass and energy estimates. Actually, using the notation $\phi(r) = r|\chi'(r)|$, we have

$$\|S(y)\| \leq \frac{1}{2} \int_{\mathbb{R}^n} \phi(|z|) dz \int_{-\frac{1}{2}}^{\frac{1}{2}} \rho(y + (s - \frac{1}{2})z) \rho(y + (s + \frac{1}{2})z) ds.$$

This yields

$$\int_{\mathbb{R}^n} \|S(y)\| dy \leq \frac{1}{2} \int_{\mathbb{R}^n \times \mathbb{R}^n} \int_{-\frac{1}{2}}^{\frac{1}{2}} \phi(|z|) \rho(y + (s - \frac{1}{2})z) \rho(y + (s + \frac{1}{2})z) ds dy dz.$$

Let us make the change of variable

$$(y, z, s) \longmapsto (v, w, s), \quad v = y + (s - \frac{1}{2})z, \quad w = y + (s + \frac{1}{2})z.$$

Remarking that $dy dz ds = dv dw ds$, we obtain

$$\int_{\mathbb{R}^n} \|S(y)\| dy \leq \frac{1}{2} \int_{\mathbb{R}^n \times \mathbb{R}^n} \phi(|w-v|) \rho(v) \rho(w) dv dw.$$

Therefore $S \in L_t^\infty(L_y^1)$ whenever the initial data has finite mass and energy and the kernel satisfies

$$|\chi'(r)| r \leq \text{cst} \cdot (1 + \chi(r)).$$

Conclusion. The moral of this example is that we should investigate in more depth other natural examples of divergence-free symmetric tensors, when T is neither positive nor negative semi-definite. We should ask ourselves whether there exists a hidden DPT A such that the equation $\operatorname{Div} T = 0$ is equivalent to $\operatorname{Div} A = 0$. The latter tensor could be defined in terms of the state of the model in a non-local way, while T is often defined in local terms.

References

- [1] R. Coifman, P.-L. Lions, Y. Meyer & S. Semmes. Compacité par compensation et espaces de Hardy. *C. R. Acad. Sci. Paris*, **309** (1989), pp 945–949.
- [2] B. D. Coleman, E. H. Dill. Thermodynamic restrictions on the constitutive equations of electromagnetic theory. *Zeitschrift für angewandte Mathematik und Physik*, **22** (1971), pp 691–702.
- [3] I. Fonseca, S. Müller. \mathcal{A} -Quasiconvexity, lower semicontinuity, and Young measures. *SIAM J. Math. Anal.*, **30** (1999), pp 1355–1390.
- [4] S. K. Godunov. Lois de conservation et intégrales d'énergie des équations hyperboliques, in *Nonlinear hyperbolic problems, Saint-Étienne 1986*, Springer-Verlag LNM **1270** (1987), pp 135–149.
- [5] S. Klainerman, M. Machedon. Space-time estimates for null forms and the local existence theorem. *Comm. Pure & Appl. Math.*, **46** (1993), pp 1221–1268.
- [6] Yan Yan Li. Some existence results of fully nonlinear elliptic equations of Monge–Ampère type. *Comm. Pure & Appl. Math.*, **43** (1990), pp 233–271.
- [7] A. V. Pogorelov. *The Minkowski multidimensional problem*. Scripta Series in Mathematics. V. H. Winston & Sons, Washington, D.C.; Halsted Press (John Wiley & Sons), New York–Toronto–London (1978).
- [8] D. Serre. Divergence-free positive symmetric tensors and fluid dynamics. *Annales de l'Institut Henri Poincaré (analyse non linéaire)*. To appear.
- [9] C. Villani. *Topics in optimal transportation*. Graduate Studies in Mathematics **58**, Amer. Math. Society (2003).