



Estimating the number and the strength of collisions in molecular dynamics

Denis Serre

► To cite this version:

Denis Serre. Estimating the number and the strength of collisions in molecular dynamics. 2019. ensl-02065701v1

HAL Id: ensl-02065701

<https://ens-lyon.hal.science/ensl-02065701v1>

Preprint submitted on 12 Mar 2019 (v1), last revised 18 Feb 2020 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Estimating the number and the strength of collisions in molecular dynamics

Denis Serre
École Normale Supérieure de Lyon*

Abstract

We consider the motion of a finite though large number of particles in the whole space \mathbb{R}^n . Particles move freely until they experience pairwise collisions. We use our recent theory of divergence-controlled positive symmetric tensors in order to establish two estimates regarding the set of collisions. The only information needed from the initial data is the total mass and the total energy.

1 Models of molecular dynamics

We consider a set of N identical particles of mass m , moving in the whole space \mathbb{R}^n . The coordinate in the physical space \mathbb{R}^{1+n} is denoted $x = (t, y)$ where t is the time and $y \in \mathbb{R}^n$ the position.

In practice $n = 3$ and N is of the size of the Avogadro number, but the analysis below is valid in every dimension and for any cardinality. We shall think of the particles as spheres of radius $a > 0$, so that a collision between two particles occurs when their centers y_α approach to a distance $2a$:

$$|y_\beta - y_\alpha| = 2a.$$

The incoming and outgoing velocities ($v_{\alpha,\beta}$ and $v'_{\alpha,\beta}$ respectively) satisfy

$$(1) \quad (v_\beta - v_\alpha) \cdot (y_\beta - y_\alpha) < 0, \quad (v'_\beta - v'_\alpha) \cdot (y_\beta - y_\alpha) \geq 0.$$

Our assumptions are as follows:

- Each particle P_α has a finite internal energy $\varepsilon_\alpha \geq 0$ and a velocity $v_\alpha \in \mathbb{R}^n$. These parameters remain constant between consecutive collisions involving P_α .

*U.M.P.A., UMR CNRS-ENSL # 5669. 46 allée d'Italie, 69364 Lyon cedex 07. France. denis.serre@ens-lyon.fr

- When P_α and P_β collide, at a given time t , their parameters (velocity, internal energy) experience a jump, which obey to the conservation of momentum and energy:

$$\begin{aligned} v'_\alpha + v'_\beta &= v_\alpha + v_\beta, \\ \frac{m}{2} (|v'_\alpha|^2 + |v'_\beta|^2) + \epsilon'_\alpha + \epsilon'_\beta &= \frac{m}{2} (|v_\alpha|^2 + |v_\beta|^2) + \epsilon_\alpha + \epsilon_\beta. \end{aligned}$$

- The collisions are friction-less, meaning that the jump of velocity is orthogonal to the common tangent space to the particles:

$$(2) \quad v'_\alpha - v_\alpha = v_\beta - v'_\beta \parallel y_\beta - y_\alpha.$$

The trajectory of a given particle is a polygonal chain. The conservation of energy implies a bound for the emerging velocities:

$$|v'_\alpha|^2 + |v'_\beta|^2 \leq |v_\alpha|^2 + \frac{2}{m} \epsilon_\alpha + |v_\beta|^2 + \frac{2}{m} \epsilon_\beta.$$

The assumptions made above cover two important cases. On the one hand that of hard spheres, for which there is no exchange of internal energy (we might as well assume that there is no internal energy at all) and therefore

$$|v'_\alpha|^2 + |v'_\beta|^2 = |v_\alpha|^2 + |v_\beta|^2.$$

On the other hand, we have the model of *sticky* particles, for which

$$v'_\alpha = v'_\beta = \frac{1}{2}(v_\alpha + v_\beta).$$

Two important quantities emerge from this considerations, namely the total mass $M = Nm$ and the total energy

$$E = \sum_{\alpha} \left(\frac{m}{2} |v_\alpha(t)|^2 + \epsilon_\alpha(t) \right),$$

which do not depend on the instant at which they are computed. The third conserved quantity (total momentum)

$$Q = m \sum_{\alpha} v_\alpha(t)$$

will not be used below, for several reasons. On the first leg its nature is vectorial, which makes hard its use for an estimate. Besides, it just vanishes if we choose an appropriate inertial frame, a flaw which does not occur to the mass or the energy. Finally, it can be estimated by Cauchy–Schwarz inequality, $|Q| \leq \sqrt{2ME}$, and therefore the knowledge of M and E will always give a sufficient information.

2 Results

Our aim is to estimate the number of collisions during the whole history, given only M and E . We shall partly achieve this goal, in the sense that we establish estimates of the form

$$\sum_{\text{coll.}} F(v_\alpha, v_\beta, v'_\alpha, v'_\beta)$$

in terms of m, M and E , where F is some explicit non-negative function, and we sum over all collisions. We confess that these estimates get poorer when either $v_\beta - v_\alpha$ is small (slow collision), or when the vectors $v_\alpha, v_\beta, v'_\alpha, v'_\beta$ are approximately coplanar (*grazing* collisions). We believe that these limitations are inherent to the context of molecular dynamics, and that they do not reveal a weakness of the mathematical tool which we employ below.

Our main results are as follows. We limit ourselves to the case where every collision involves exactly two particles, and the evolution is defined for every time. This is a generic situation, as shown by Alexander in his PhD thesis [1] ; see Theorem 4.2.1 of [4]. See also Uchiyama's analysis [12] of the Broadwell model, a model with discrete velocity set. Our estimates involve the wedge product of two or three vectors in arbitrary dimensions ; we explain this notion, that comes from exterior calculus, in Paragraph 4.2. The product of ℓ vectors in \mathbb{R}^n always vanishes if $\ell > n$.

Theorem 2.1 *Consider a finite system of particles moving in the physical space \mathbb{R}^n according to the laws described above. Let $M = Nm$ be the total mass and $E < \infty$ be the total energy of the system. Let us make the generic assumption that the collision set is locally finite, and that the motion involves only binary collisions.*

At every collision, denote v, v_1 the incoming velocities and v', v'_1 the outgoing ones (they vary from one collision to another one, although the notation remains the same).

Then there exists a universal constant such that the following inequalities hold true

$$(3) \quad m^2 \sum_{\text{coll.}} \left(\frac{E|v' - v|^2 + M|v \wedge v'|^2}{\sqrt{(E + M|v|^2)(E + M|v'|^2)}} + \frac{E|v'_1 - v_1|^2 + M|v_1 \wedge v'_1|^2}{\sqrt{(E + M|v_1|^2)(E + M|v'_1|^2)}} \right) \leq c_n M E,$$

where the sum runs over the set of collisions. When $n \geq 2$, we also have

$$(4) \quad m^{\frac{3}{2}} \sum_{\text{coll.}} \frac{AB}{C} \leq c_n M^{\frac{1}{2}} E^{\frac{3}{4}},$$

where for each collision,

$$\begin{aligned} A &= (E|(v' - v) \wedge (v_1 - v)|^2 + M|v \wedge v' \wedge v_1|^2)^{\frac{1}{2}} \\ B &= (4E + M(|v|^2 + |v'|^2 + |v_1|^2 + |v'_1|^2))^{1/4} \\ C &= ((E + M|v|^2)(E + M|v'|^2)(E + M|v_1|^2)(E + M|v'_1|^2))^{\frac{1}{4}}. \end{aligned}$$

By using the inequality $|v \wedge v'| \leq |v| \cdot |v' - v|$, we see that (3) implies a simpler estimate:

Corollary 2.1 *With the same assumptions as in Theorem 2.1, we have*

$$(5) \quad m^2 \sum_{\text{coll.}} \frac{|v \wedge v'|^2}{|v| \cdot |v'|} \leq c_n M E$$

for some universal constant c_n .

It seems that (4) cannot be simplified in a similar way.

Comments.

- These estimates are independent of the size a of the particles. They are also dimension-independent, apart for the constants c_n .
- The expressions A, B and C are symmetric functions of the velocities v, v', v_1 and v'_1 (obvious for B and C).
- The estimates above do not tell us whether the number of collisions is finite.
- Nevertheless (3) tells us something about the number of the "strongest" ones. In terms of a typical velocity $\mathbf{v} := \sqrt{E/M}$, the number of collisions for which $|v \wedge v'|^2 \geq \epsilon \mathbf{v}^2 |v| \cdot |v'|$ is bounded by $c_n N^2 \epsilon^{-1}$.
- Likewise, (4) tells us that the number of collisions for which $|(v' - v) \wedge (v_1 - v)| \geq \epsilon \mathbf{v}^2$ and $|v|, |v'|, |v_1|, |v'_1| \leq A \mathbf{v}$ for some $0 < \epsilon < A < \infty$, is bounded by $C_{\epsilon, A} N^{\frac{3}{2}}$. On the contrary, it says very little about grazing collisions.

These polynomial bounds should be compared with those known for the whole set of collisions. For the hard spheres model with elastic collisions, Alexander [1] and Vaserstein [13] showed that this set is finite, a problem raised by Ya. Sinai. The proof of finiteness was simplified by Illner [6, 7]. So far, these works argued by contradiction and did not give an upper bound of the collision number. Later on Burago & all. [2] established the upper bound

$$(32N^{\frac{3}{2}})^{N^2}.$$

This seems way too large to be accurate, since the fate of the particles is to disperse towards infinity in independent directions.

On the opposite side, some authors constructed configurations for which the number of collisions is superlinear in the number of particles. When the energy is not conserved and may increase arbitrarily, one can even observe infinitely many collisions [6, 5]. For hard spheres with elastic collisions, Burdzy & Duarte [3] announce that some configuration leads to at least

$$\frac{N^3}{27}$$

collisions, a number which sounds reasonable. Let us point out that, according to our estimates, most of these collisions must be weak and grazing.

Plan of the paper. We begin by constructing in Section 3 the mass-momentum tensor associated with the motion. It is a map $x \mapsto T(x)$ taking values in the cone of positive semi-definite matrices Sym_{1+n}^+ , albeit a singular one: the entries are bounded measures and their support is one-dimensional. The conservation of mass and momentum is expressed by the row-wise identity $\text{Div} T = 0$. One striking feature in this construction is the introduction of massless virtual particles (*collitons*) whose role is to carry the exchange of momentum between colliding particles. Section 4 presents the tools that will be used in the analysis. It is mainly a recall of our theory of Compensated Integrability for symmetric positive tensors whose row-wise divergence is a (vector-valued) bounded measure. We fix also a few notations related to exterior calculus. Sections 5 and 6 present the proofs of the binary and ternary estimates, respectively. We combine Compensated Integrability with a trick which we developed for the first time in [11].

Acknowledgement. I am indebted to Laure Saint-Raymond and Reinhard Illner for valuable discussions and their help in gathering the relevant literature.

3 The mass-momentum tensor

From now on, we denote $d = 1 + n$ the time-space dimension.

3.1 Single particle

We begin by considering a single particle P whose constant velocity is $v \in \mathbb{R}^n$. The trajectory $t \mapsto (t, y(t))$ of the center of mass in the physical space \mathbb{R}^{1+n} is a line L , whose direction is

$$\xi = \frac{V}{|V|}, \quad \text{where } V := \begin{pmatrix} 1 \\ v \end{pmatrix}.$$

Let us define a symmetric matrix, whose entries are locally finite measures over \mathbb{R}^{1+n} , by

$$S = mV \otimes \xi \delta_L = m|V| \xi \otimes \xi \delta_L.$$

In other words

$$\langle S_{ab}, \phi \rangle = mV_a \xi_b \int_{\mathbb{R}} \phi(\bar{x} + s\xi) ds, \quad \forall \phi \in C_K(\mathbb{R}^{1+n}),$$

where \bar{x} is any point on the line L .

Lemma 3.1 *One has*

$$\text{Div} S = 0,$$

an equation that stands row-wise. More generally, if $Q \in \mathbb{R}^d$ with $Q \neq 0$, and $\eta = \frac{Q}{|Q|}$, then for every line $L = \bar{x} + \mathbb{R}\eta$, the symmetric tensor

$$S^Q := Q \otimes \eta \delta_L$$

is divergence-free.

Notice that for a particle, S is nothing but S^Q with $Q = mV$. Remark also that S^Q is everywhere positive semi-definite.

Proof

If ϕ is a test function, then

$$\langle \text{Div } S^Q, \phi \rangle = -\langle S^Q, \nabla \phi \rangle = -Q \int_{\mathbb{R}} \eta \cdot \nabla \phi(\bar{x} + s\eta) ds = -Q \int_{\mathbb{R}} \frac{d}{ds} \phi(\bar{x} + s\eta) ds = 0.$$

■

3.2 Multi-line configuration

When L is replaced by a semi-infinite line $L^+ = \bar{x} + \mathbb{R}_+ \eta$, the tensor

$$S^{Q+} := Q \otimes \eta \delta_{L^+}$$

is no longer divergence-free. The calculation above yields

$$(6) \quad \langle \text{Div } S^{Q+}, \phi \rangle = \phi(\bar{x})Q,$$

which is recast as

$$\text{Div } S^{Q+} = Q \delta_{\bar{x}}.$$

Now, if finitely many vectors Q_1, Q_2, \dots are given, together with a point $\bar{x} \in \mathbb{R}^d$, we may form the converging semi-lines $L_j^+ = \bar{x} + \mathbb{R}_+ Q_j$ and define a symmetric tensor

$$S_{\text{mult}} := \sum_j S^{Q_j+}.$$

Then (6) tells us that S_{mult} is divergence-free whenever

$$(7) \quad \sum_j Q_j = 0.$$

Application to 1-D molecular dynamics. When $n = 1$, we may simplify the model by setting $a = 0$. At a binary collision, the point particles meet at some point $\bar{x} \in \mathbb{R}^{1+1}$, with incoming velocities v, w and outgoing ones v', w' . Let us choose

$$(8) \quad V_1 = -\begin{pmatrix} 1 \\ v \end{pmatrix}, \quad V_2 = -\begin{pmatrix} 1 \\ w \end{pmatrix}, \quad V_3 = \begin{pmatrix} 1 \\ v' \end{pmatrix}, \quad V_4 = \begin{pmatrix} 1 \\ w' \end{pmatrix}$$

and $Q_j = mV_j$. Then the positive semi-definite tensor

$$S^{Q_1+} + S^{Q_2+} + S^{Q_3+} + S^{Q_4+}$$

associated with this pair of particles is divergence-free ; the compatibility condition $Q_1 + Q_2 + Q_3 + Q_4 = 0$ is ensured by the conservation of mass and momentum through the collision. Its support is the union of the trajectories.

3.3 Binary collisions ($n \geq 2$)

When $n \geq 2$ instead, the radius a must be positive, in order that collisions take place.

Let two particles P_i and P_j collide at some time t^* . The trajectory of P_i displays a kink at a point $\bar{x}_i = (t^*, \bar{y}_i)$, and that of P_j does at $\bar{x}_j = (t^*, \bar{y}_j)$ at the same instant t^* . We have $|\bar{y}_j - \bar{y}_i| = 2a$. Let us define V_1, \dots, V_4 as in (8). Locally, the trajectories are made of segments of the semi-lines

$$L_1^+ = \bar{x}_i + \mathbb{R}_+ V_1, \quad L_2^+ = \bar{x}_j + \mathbb{R}_+ V_2, \quad L_3^+ = \bar{x}_i + \mathbb{R}_+ V_3, \quad L_4^+ = \bar{x}_j + \mathbb{R}_+ V_4.$$

Let us denote again $Q_j = mV_j$. Because these do not meet at a single point, the tensor $S = S^{\mathcal{Q}_1^+} + S^{\mathcal{Q}_2^+} + S^{\mathcal{Q}_3^+} + S^{\mathcal{Q}_4^+}$ is not divergence-free. We have instead

$$\text{Div } S = (Q_1 + Q_3)\delta_{x_i} + (Q_2 + Q_4)\delta_{x_j} = (Q_1 + Q_3)(\delta_{x_i} - \delta_{x_j}).$$

In order to recover a divergence-free tensor, we introduce a vector Q

$$Q = \begin{pmatrix} 0 \\ q \end{pmatrix}, \quad q = m(v' - v) = m(w - w').$$

Because of (2), the segment $C = [\bar{x}_i, \bar{x}_j]$ has direction Q . In the neighbourhood of the collision, we can define the tensor

$$T = S^{\mathcal{Q}_1^+} + S^{\mathcal{Q}_2^+} + S^{\mathcal{Q}_3^+} + S^{\mathcal{Q}_4^+} + Q \otimes \eta \delta_C.$$

Each of the five terms in the sum above is divergence-free away from either \bar{x}_i or \bar{x}_j . At \bar{x}_i , $\text{Div } T$ is a sum of three Dirac masses, whose weight is

$$(9) \quad m(V_1 + V_3) - Q = \begin{pmatrix} 0 \\ -mv + mv' - q \end{pmatrix} = 0,$$

where the minus sign in front of Q comes from the fact that Q is oriented from x_j to x_i . A similar identity holds true at \bar{x}_j , though with a plus sign. We conclude that

$$\text{Div } T = 0.$$

We may interpret the contribution $Q \otimes \eta \delta_C$ as that of a virtual particle. This particle is mass-less, because the first component of Q vanishes. It carries a momentum which is exchanged instantaneously between P_i and P_j . We suggest the name *colliton* for this object. If we took in account relativistic effects, there would be instead a pair of virtual particles (a particle and its anti-particle), travelling at the speed of light.

3.4 The complete construction

Assuming again that only binary collisions happen, we consider the union of trajectories of the centers of the N particles. Each trajectory is a polygonal chain whose kinks occur where and when the particle

suffers a collision. Each segment J of a trajectory between two consecutive collisions contributes, as explained above, with the tensor

$$mV \otimes \xi \delta_J, \quad V = \begin{pmatrix} 1 \\ v \end{pmatrix}, \quad \xi = \frac{V}{|V|},$$

where v is the particle velocity along J . At a collision we also have the contribution of the corresponding colliton, as described in the previous paragraph. The sum T of all these contributions is a divergence-free positive semi-definite tensor, which we call the *mass-momentum* tensor of the configuration.

We point out that the support of T is a graph, a one-dimensional object in \mathbb{R}^{1+n} . Thus T vanishes almost everywhere in the Lebesgue sense. The support can be equipped with the positive measure $\text{Tr } T$, with respect to which T is rank-1 almost everywhere. We point out however that in order to apply Compensated Integrability (this tool is described in the next Section), we need to work with tensors of full rank $1+n$ (rather than rank-one), positive definite over a set of positive Lebesgue measure. To reach these goals, one option is to regularize T by means of a convolution ; at a kink, this will allow us to combine three contributions, associated with V_1, V_3 and Q with the notations of (8) (two branches of a trajectory, plus the colliton). Such a combination has rank 2 only since $m(V_1 + V_3) = Q$. If we increase slightly the support of the convolution kernel, we may benefit of the combination of five contributions, those associated with V_1, \dots, V_4 and Q , whose rank is generically 3. This is still not sufficient if $n = 3$ (the size of the mass-momentum tensors is $1+n$). We thus need an extra trick in order to conclude. The price to pay is that the resulting tensor will no longer be divergence-free ; this is the reason why we shall state Theorem 4.1 in the more general context of divergence-controlled tensors.

4 Miscellaneous tools

The determinant over Sym_d^+

The cone Sym_d^+ of $d \times d$ positive semi-definite symmetric matrices is convex. On this cone, the map $K \mapsto \det K$ takes non-negative values, and it is monotonous: if $0_d \prec K \prec K'$ where the order is that between the associated quadratic forms, then

$$\det K \leq \det K'.$$

Actually, the d -th root $K \mapsto (\det K)^{\frac{1}{d}}$ is a concave function over Sym_d^+ (Theorem 6.10 in [8]), homogeneous of degree one:

$$(K_1, \dots, K_\ell \in \text{Sym}_d^+) \implies \left((\det(K + \dots + K_\ell))^{\frac{1}{d}} \geq (\det K)^{\frac{1}{d}} + \dots + (\det K_\ell)^{\frac{1}{d}} \right).$$

Since non-negative numbers p_1, \dots, p_ℓ satisfy $(p_1 + \dots + p_\ell)^\alpha \geq p_1^\alpha + \dots + p_\ell^\alpha$ for every exponent $\alpha \geq 1$, we infer (take $\alpha = \frac{d}{d-1}$)

$$(10) \quad (K_1, \dots, K_\ell \in \mathbf{Sym}_d^+) \implies \left((\det(K + \dots + K_\ell))^{\frac{1}{d-1}} \geq (\det K)^{\frac{1}{d-1}} + \dots + (\det K_\ell)^{\frac{1}{d-1}} \right).$$

Notice that since this inequality is homogeneous of degree $\alpha > 1$, we cannot deduce the concavity of $K \mapsto (\det K)^{\frac{1}{d-1}}$. The latter function is actually superlinear, thus non-concave.

Let S be a symmetric tensor whose entries are distributions. Suppose that for every $\xi \in \mathbb{R}^d$, $\xi^T S \xi$ is non-negative, and therefore is a locally bounded measure. Then the entries of S are locally bounded measures. In particular $\mu := \text{Tr } S$ is a non-negative locally bounded measure, and every S_{ij} is absolutely continuous with respect to μ . This allows us to define uniquely the expression $(\det S)^{\frac{1}{d}}$ as a locally bounded measure, absolutely continuous with respect to μ ; just write $S_{ij} = s_{ij}\mu$ and define

$$(\det S)^{\frac{1}{d}} = (\det s)^{\frac{1}{d}} \mu.$$

This definition inherits the properties of the determinant over \mathbf{Sym}_d^+ , in particular the monotonicity and the concavity.

4.1 Compensated integrability

We shall make use of our recent theory of Compensated Integrability for divergence-controlled positive symmetric tensors, for which we refer to [9, 10]. The appropriate version is given in the theorem below. We denote $\|\mu\|$ for the total mass of a (vector-valued) bounded measure μ ,

$$\|\mu\| = \langle |\mu|, \mathbf{1} \rangle.$$

This notation applies below in two distinct contexts, whether μ is a measure over an $(n+1)$ -dimensional slab $H = (t_-, t_+) \times \mathbb{R}^n$, or a measure over \mathbb{R}^n .

Theorem 4.1 *Let $H = (t_-, t_+) \times \mathbb{R}^n$ be a slab in $\mathbb{R} \times \mathbb{R}^n$, and S be symmetric positive semi-definite tensor defined over H , whose entries are bounded measures. We assume that the row-wise divergence of S is also a (vector-valued) bounded measure. Finally we assume that the normal traces $S\vec{e}_i$ at the initial and final times $t = t_\pm$ are themselves bounded measures.*

Then $(\det S)^{\frac{1}{n+1}}$ belongs to $L^{1+\frac{1}{n}}(H)$ and we have

$$(11) \quad \int_H (\det S)^{\frac{1}{n}} dy dt \leq c_n (\|S\vec{e}_i(t_-)\| + \|S\vec{e}_i(t_+)\| + \|\text{Div } S\|)^{1+\frac{1}{n}},$$

where c_n is a finite constant independent of S and H .

Remarks.

- The assumption that $\text{Div } S$ is a bounded measure allows us to define a normal trace in a rather weak space, here the dual of $\text{Lip}(H)$. This is reminiscent to the definition of the normal trace of vector fields $\vec{q} \in H_{\text{div}}(\Omega)$, used in the theory of incompressible fluids, which takes values in $H^{-1/2}(\partial\Omega)$.
- The additional assumption that this trace is a bounded measure is equivalent to saying that the extension \tilde{S} by 0 away from H enjoys too the property that $\text{Div } \tilde{S}$ is a bounded measure. Then

$$\text{Div } \tilde{S} = \widetilde{\text{Div } S} - S\vec{e}_t(t_-) \otimes \delta_{t=t_-} + S\vec{e}_t(t_+) \otimes \delta_{t=t_+}$$

- The qualitative part of the theorem is that the bounded measure $(\det S)^{\frac{1}{d}}$ is absolutely continuous with respect to the Lebesgue measure, and that its density is a function of class $L^{\frac{d}{d-1}}$, where $\frac{d}{d-1} = 1 + \frac{1}{n}$. The quantitative part (11) estimates this density.
- This theorem is useless when S is rank-1 almost everywhere ; the estimate (11) is then trivial.

4.2 Exterior calculus

We use a natural generalization of the determinant of a system of vectors. Let $Z^{(1)}, \dots, Z^{(k)}$ be a list of vectors in \mathbb{R}^d , with $k \leq d$, and let $Z \in \mathbf{M}_{d \times k}(\mathbb{R})$ be the matrix whose columns are the $Z^{(j)}$'s. Let us decompose them on the canonical basis,

$$Z^{(j)} = \sum_{i=1}^d z_{ij} \vec{e}_i.$$

Then the exterior product $Z^{(1)} \wedge \dots \wedge Z^{(k)}$ decomposes over the basis $\vec{e}_{i_1} \wedge \dots \wedge \vec{e}_{i_k}$ with $i_1 < \dots < i_k$ of the exterior power $\Lambda^k(\mathbb{R}^d)$. Its coordinates are the $k \times k$ minors of Z . For instance if $d = 3$, $Z \wedge Z'$ identifies naturally with the cross product.

We denote

$$|Z^{(1)} \wedge \dots \wedge Z^{(k)}|$$

the natural euclidian norm of the exterior product, which is invariant upon the action of the orthogonal group:

$$|RZ^{(1)} \wedge \dots \wedge RZ^{(k)}| = |Z^{(1)} \wedge \dots \wedge Z^{(k)}|, \quad \forall R \in O_d(\mathbb{R}).$$

A practical tool is given by the formula

$$|Z^{(1)} \wedge \dots \wedge Z^{(k)}|^2 = \det \left(\langle Z^{(j)}, Z^{(\ell)} \rangle \right)_{1 \leq j, \ell \leq k}.$$

Suppose $k + \ell \leq d$. If $\text{Span}(Z^{(1)}, \dots, Z^{(k)})$ and $\text{Span}(W^{(1)}, \dots, W^{(\ell)})$ are orthogonal to each other, then

$$|Z^{(1)} \wedge \dots \wedge Z^{(k)} \wedge W^{(1)} \wedge \dots \wedge W^{(\ell)}| = |Z^{(1)} \wedge \dots \wedge Z^{(k)}| \cdot |W^{(1)} \wedge \dots \wedge W^{(\ell)}|.$$

In particular, we shall use the formula

$$|Z^{(1)} \wedge \dots \wedge Z^{(k)}| = \left| \det(Z^{(1)}, \dots, Z^{(k)}, v_{k+1}, \dots, v_d) \right|,$$

where (v_{k+1}, \dots, v_d) is any unitary basis of the subspace $\text{Span}(Z^{(1)}, \dots, Z^{(k)})^\perp$.

5 The binary estimate

In this section, we supplement the mass-momentum tensor T with a contribution associated with the changes $v \mapsto v'$ in the direction of particles motions. Let K be a kink of a trajectory, happening at a point x^* . The incoming/outgoing velocities being v, v' respectively, with $v' \neq v$, we complete the free family

$$V = \begin{pmatrix} 1 \\ v \end{pmatrix}, \quad V' = \begin{pmatrix} 1 \\ v' \end{pmatrix}$$

into a basis (V, V', z_2, \dots, z_n) of \mathbb{R}^{1+n} , where (z_2, \dots, z_n) is some orthonormal basis of $\text{Span}(V, V')^\perp$. To define the kink contribution, we introduce the positive semi-definite tensor

$$S_K = \sum_{j=2}^n z_j \otimes z_j \delta_{\sigma_j},$$

where $\sigma_j := (x^* - az_j, x^* + az_j)$ is an interval. Because $\text{Div } S_K$ is a sum of Dirac masses at the end points $x^* \pm az_j$, we have

$$\|\text{Div } S_K\| = 2(n-1).$$

Given a slab H , we form the tensor

$$T' = T + \sum_{\text{kinks in } H} b_K S_K,$$

where the positive numbers b_K will be chosen later. We have

$$\|\text{Div } T'\| \leq 2(n-1) \sum_{\text{kinks in } H} b_K,$$

where the inequality reflects the fact that a few end points could lie away from H .

Now, we make the convolution product $T'_{2a} = \phi_{2a} * T'$ with the non-negative kernel

$$\phi_{2a} = \frac{1}{|B_{2a}|} \mathbf{1}_{B_{2a}},$$

where B_{2a} is the ball of radius $2a$ centered at the origin, and $\mathbf{1}$ denotes the characteristic function. This new tensor is still symmetric, positive semi-definite, and satisfies

$$\|\text{Div } T'_{2a}\| \leq 2(n-1) \sum_{\text{kinks in } H} b_K.$$

Let K be a kink in H , occuring at point x^* . If $x \in B_a(x^*)$, then $B_{2a}(x)$ contains $B_a(x^*)$. In particular $B_{2a}(x)$ meets every branch of the support of T' around x^* along a segment of length $\geq a$. There follows that

$$T'_{2a}(x) \geq \frac{a}{|B_{2a}|} \left(mV \otimes \xi + mV' \otimes \xi' + b_K \sum_2^n z_j \otimes z_j \right), \quad \forall x \in B_a(x^*).$$

By the monotonicity of the determinant, we infer

$$\det T_{2a}(x) \geq \frac{m^2 b_K^{n-1}}{a^{dn} |B_2|^d |V| \cdot |V'|} (\det(V, V', z_2, \dots, z_n))^2 = \frac{m^2 b_K^{n-1}}{a^{dn} |B_2|^d} \frac{|V \wedge V'|^2}{|V| \cdot |V'|}.$$

We now apply Theorem 4.1 to T'_{2a} in the slightly larger slab $H + B_{2a}$. Let us assume first that the distance between two kinks in H is $\geq 2a$, so that the balls $B_a(x^*)$ are disjoint. Then the integral in the left hand side is estimated below by

$$(12) \quad \int_{H+B_{2a}} (\det T'_{2a}(x))^{\frac{1}{n}} dx \geq \frac{2^{-d} m^{2/n}}{|B_2|^{1/n}} \sum_{\text{kinks}} b_K^{1-\frac{1}{n}} \left(\frac{|V \wedge V'|^2}{|V| \cdot |V'|} \right)^{\frac{1}{n}}.$$

When two (or more) balls $B_a(x^*)$ may overlap, we use Inequality (10) to get

$$(\det T_{2a}(x))^{\frac{1}{n}} \geq \frac{m^{2/n}}{a^d |B_2|^{d/n}} \sum' b_K^{1-\frac{1}{n}} \left(\frac{|V \wedge V'|^2}{|V| \cdot |V'|} \right)^{\frac{1}{n}},$$

where the sum runs over all the kinks K for which $x \in B_a(x^*)$. By integrating over $x \in H$ and then rearranging the sum, we deduce again the same lower bound (12).

From (11) and (12), we deduce

$$m^{2/n} \sum_{\text{kinks}} b_K^{1-\frac{1}{n}} \left(\frac{|V \wedge V'|^2}{|V| \cdot |V'|} \right)^{\frac{1}{n}} \leq c_n \left(\|T'_{2a} \vec{e}_t(t_- - 2a)\| + \|T'_{2a} \vec{e}_t(t_+ + 2a)\| + 2(n-1) \sum_{\text{kinks}} b_K \right)^{1+\frac{1}{n}}$$

for some universal constant c_n ; we recall that the kinks in the summation are those in H . There remains to estimate the masses in the right hand side. We notice that

$$\|T'_{2a} \vec{e}_t(t_+ + 2a)\| \leq \sup_{t > t_+} \|T' \vec{e}_t(t)\| = \sup_{t > t_+} \|T \vec{e}_t(t)\| = m \sup_{t > t_+} \sum |V|,$$

where the sum runs over all particles. Because of $|V| = \sqrt{1 + |v|^2} \leq 1 + \frac{1}{2}|v|^2$, we infer

$$\|T'_{2a} \vec{e}_t(t_+ + 2a)\| \leq M + E$$

and we deduce an explicit inequality

$$(13) \quad m^{2/n} \sum_{\text{kinks}} b_K^{1-\frac{1}{n}} \left(\frac{|V \wedge V'|^2}{|V| \cdot |V'|} \right)^{\frac{1}{n}} \leq c_n \left(M + E + (n-1) \sum_{\text{kinks}} b_K \right)^{1+\frac{1}{n}}$$

with a slightly different constant c_n .

We exploit (13) by letting the coefficients b_K vary. Let us choose positive numbers β_K and another positive parameter λ . By setting

$$b_K = \lambda \beta_K^{\frac{n}{n-1}},$$

we have

$$m^{2/n} \sum_{\text{kinks}} \beta_K \left(\frac{|V \wedge V'|^2}{|V| \cdot |V'|} \right)^{\frac{1}{n}} \leq c_n \lambda^{\frac{1}{n}-1} \left(M + E + \lambda(n-1) \sum_{\text{kinks}} \beta_K^{\frac{n}{n-1}} \right)^{1+\frac{1}{n}}.$$

Choosing

$$\lambda = \frac{M+E}{\sum \beta_K^{\frac{n}{n-1}}},$$

we obtain

$$m^{2/n} \sum_{\text{kinks}} \beta_K \left(\frac{|V \wedge V'|^2}{|V| \cdot |V'|} \right)^{\frac{1}{n}} \leq c_n (M+E)^{\frac{2}{n}} \left(\sum_{\text{kinks}} \beta_K^{\frac{n}{n-1}} \right)^{1-\frac{1}{n}} = c_n (M+E)^{\frac{2}{n}} \left\| \vec{\beta} \right\|_{\ell^{n/(n-1)}}.$$

Since $\frac{n}{n-1}$ is the conjugate exponent of n , a convenient choice of the vector $\vec{\beta}$ gives us an estimate of the ℓ^n -norm of the vector whose coordinates are the expressions

$$\left(\frac{|V \wedge V'|^2}{|V| \cdot |V'|} \right)^{\frac{1}{n}}.$$

Specifically, we have proved

$$(14) \quad m^2 \sum_{\text{kinks}} \frac{|V \wedge V'|^2}{|V| \cdot |V'|} \leq c_n (M+E)^2,$$

with again a slightly different constant c_n . This estimate can be recast in terms of the velocities only, by using

$$|V| = \sqrt{1 + |v|^2}, \quad |V \wedge V'|^2 = |v - v'|^2 + |v \times v'|^2.$$

We point out that the constant in (14) depends only on the space dimension and not on the data. In particular, it does not depend upon the width of the slab H . Since the right-hand side depends only on the initial data, we deduce that (14) is valid when we sum over the whole set of kinks in \mathbb{R}^{1+n} .

The last step involves a rescaling. Keeping the same space variable y and changing the time variable into $\bar{t} = \mu^{-1}t$ has the effect of replacing the velocities v by μv . The mass is preserved, while the energy becomes $\mu^2 E$. Applying (14) to the new dependent/independent variables, we obtain a parametrized estimate

$$m^2 \sum_{\text{kinks}} \frac{\mu^2 |v' - v|^2 + \mu^4 |v \wedge v'|^2}{\sqrt{(1 + \mu^2 |v|^2)(1 + \mu^2 |v'|^2)}} \leq c_n (M + \mu^2 E)^2, \quad \forall \mu > 0.$$

Choosing now $\mu^2 = M/E$, we arrive at our final estimate, which is now invariant upon the rescaling of the time and space variables:

$$m^2 \sum_{\text{kinks}} \frac{E|v' - v|^2 + M|v \wedge v'|^2}{\sqrt{(E + M|v|^2)(E + M|v'|^2)}} \leq c_n M E.$$

6 The ternary estimate

We change slightly the strategy described in the previous section, by considering the set of collisions instead of that of kinks. At a collision C between two particles P_α and P_β , we denote \bar{x} the middle point between the kinks $x_{\alpha,\beta}$. The incoming/outgoing velocities are denoted v, v_1, v', v'_1 . When the vectors $v' - v$ and $y_\beta - y_\alpha$ are not colinear¹, the vectors V, V' and $x_\beta - x_\alpha$ are not coplanar; they span a 3-dimensional subspace which contains also V_1 and V'_1 . Then we choose an orthonormal basis (w_3, \dots, w_n) of the subspace $\text{Span}(V, V', x_\beta - x_\alpha)^\perp$, and we define a tensor

$$S_C = \sum_{j=3}^n w_j \otimes w_j \delta_{\theta_j}, \quad \theta_j := (\bar{x} - aw_j, \bar{x} + aw_j).$$

As above, $\text{Div} S_C$ is a sum of Dirac masses at the end points $\bar{x} \pm aw_j$, whose total mass is $2(n-2)$.

Let $H = (t_-, t_+) \times \mathbb{R}^n$ be a slab. We say that a collision C is in H if it happens at a time $t^* \in (t_-, t_+)$. We form the symmetric, positive semi-definite tensor

$$T'' = T + \sum_{\text{coll. in } H} b_C S_C,$$

where the positive numbers b_C are to be chosen. Then we make the convolution T''_{3a} of T'' with the kernel ϕ_{3a} . The total mass of $\text{Div} T''$ is $2(n-2) \sum b_C$.

Consider a collision C in H . For any point x in $B_a(\bar{x})$, the ball $B_{3a}(x)$ contains the balls $B_a(x_{\alpha,\beta})$, thus meets the five segments involved in the collision pattern along intervals of lengths at least a . Therefore we have

$$T''_{3a}(x) \geq \frac{a}{|B_{3a}|} \left(m(V \otimes \xi + V' \otimes \xi' + V_1 \otimes \xi_1 + V'_1 \otimes \xi'_1 + \bar{V} \otimes \bar{\xi}) + b_C \sum_3^n w_j \otimes w_j \right), \quad \forall x \in B_a(\bar{x})$$

where

$$\bar{V} = \begin{pmatrix} 0 \\ v' - v \end{pmatrix}, \quad \bar{\xi} = \frac{\bar{V}}{|\bar{V}|}$$

accounts for the collision. With the monotonicity of the determinant, we infer the coarse lower bound

$$\det T''_{3a}(x) \geq \frac{m^3 b_C^{n-2}}{a^{dn} |B_3|^d} \frac{|V \wedge V' \wedge V_1|^2}{|V| \cdot |V'| \cdot |V_1|}.$$

¹This is the generic situation where the trajectories are not coplanar

More generally, if x is a -close to several collisions, then we have

$$(\det T''_{3a}(x))^{\frac{1}{n}} \geq \frac{m^{3/n}}{a^d |B_3|^{d/n}} \sum' b_C^{1-2/n} \left(\frac{|V \wedge V' \wedge V_1|^2}{|V| \cdot |V'| \cdot |V_1|} \right)^{\frac{1}{n}},$$

where the sum runs over those collisions C in H for which $x \in B_a(\bar{x})$.

Integrating with respect to x in $H + B_a$, and rearranging the sum, we obtain

$$\int_{H+B_a} (\det T''_{3a}(x))^{\frac{1}{n}} dx \geq \frac{m^{3/n} |B_1|}{|B_3|^{d/n}} \sum_{\text{coll. in } H} b_C^{1-2/n} \left(\frac{|V \wedge V' \wedge V_1|^2}{|V| \cdot |V'| \cdot |V_1|} \right)^{\frac{1}{n}},$$

Applying (11) to T''_{3a} in $H + B_a$, we deduce again

$$m^{\frac{3}{n}} \sum_{\text{coll. in } H} b_C^{1-2/n} \left(\frac{|V \wedge V' \wedge V_1|^2}{|V| \cdot |V'| \cdot |V_1|} \right)^{\frac{1}{n}} \leq c_n \left(M + E + (n-2) \sum_{\text{coll. in } H} b_C \right)^{1+\frac{1}{n}}.$$

Introducing as above auxiliary positive parameters λ and β_C , with

$$b_C = \lambda \beta_C^p, \quad p := \frac{n}{n-2},$$

and choosing

$$\lambda = \frac{M+E}{\beta_C^p},$$

we deduce

$$m^{\frac{3}{n}} \sum_{\text{coll. in } H} \beta_C \left(\frac{|V \wedge V' \wedge V_1|^2}{|V| \cdot |V'| \cdot |V_1|} \right)^{\frac{1}{n}} \leq c_n (M+E) \|\vec{\beta}\|_{\ell^p}.$$

An appropriate choice of $\vec{\beta}$ yields an estimate in the $\ell^{p'}$ -norm, where $p' = \frac{n}{2}$. Namely, we have

$$(15) \quad m^{\frac{3}{2}} \sum_{\text{coll. in } H} \frac{|V \wedge V' \wedge V_1|}{\sqrt{|V| \cdot |V'| \cdot |V_1|}} \leq c_n (M+E)^{\frac{3}{2}}.$$

Remark that we used only the three first factors among $V \otimes \xi, V' \otimes \xi', V_1 \otimes \xi_1$ and $V'_1 \otimes \xi'_1$ in order to bound by below the determinant of $T''_{3a}(x)$. We might as well have taken any three of the four. Remarking that the wedge product remains the same, up to the sign, because of $V + V_1 = V' + V'_1$, we deduce that (15) remains valid when we replace $|V| \cdot |V'| \cdot |V_1|$ by any other product of three norms in the denominator. In other terms, we have

$$m^{\frac{3}{2}} \sum_{\text{coll. in } H} \frac{|V \wedge V' \wedge V_1|}{\sqrt{|V| \cdot |V'| \cdot |V_1| \cdot |V'_1|}} \left(|V|^{1/2} + |V'|^{1/2} + |V_1|^{1/2} + |V'_1|^{1/2} \right) \leq c_n (M+E)^{\frac{3}{2}},$$

where we now sum a quantity that is symmetric in the four velocities.

Our next remark is, as before, that the left-hand side does not depend upon the choice of the slab. Thus our inequality is valid when we sum over the set of all collisions in \mathbb{R}^{1+n} .

We end our analysis as above by applying the rescaling argument. This transform an inhomogeneous estimate into an homogeneous one:

$$m^{\frac{3}{2}} \sum_{\text{coll.}} \frac{(E|(v' - v) \wedge (v_1 - v)|^2 + M|v \wedge v' \wedge v_1|^2)^{\frac{1}{2}} (4E + M(|v|^2 + |v'|^2 + |v_1|^2 + |v'_1|^2))^{1/4}}{((E + M|v|^2)(E + M|v'|^2)(E + M|v_1|^2)(E + M|v'_1|^2))^{\frac{1}{4}}} \leq c_n M^{\frac{1}{2}} E^{\frac{3}{4}},$$

where we have used also the identity

$$|V \wedge V' \wedge V_1|^2 = |(v' - v) \wedge (v_1 - v)|^2 + |v \wedge v' \wedge v_1|^2.$$

References

- [1] R. K. Alexander. *The infinite hard sphere system*. Ph.D. thesis, University of California at Berkeley, (1975).
- [2] D. Burago, S. Ferleger, A. Kononenko. Uniform estimates on the number of collisions in semi-dispersing billiards. *Annals of Mathematics*, Second series, **147** (1998), pp 695–708.
- [3] K. Burdzy, M. Duarte. Lower bound for the number of elastic collisions. ArXiv preprint, arXiv:1803.00979, 2018.
- [4] C. Cercignani, R. Illner, M. Pulvirenti. *The mathematical theory of dilute gases*. Springer-Verlag, New York (1994).
- [5] Xinfu Chen, R. Illner. Finite range repulsive systems of finitely many particles. *Arch. Rational Mech. Anal.*, **173** (2004), pp 1–24.
- [6] R. Illner. On the number of collisions in a hard sphere particle system in all space. *Transport Theory and Stat. Phys.*, **18** (1989), pp 71–86.
- [7] R. Illner. Finiteness of the number of collisions in a hard sphere particle system in all space II: arbitrary diameters and masses. *Transport Theory and Stat. Phys.*, **19** (1990), pp 573–579.
- [8] D. Serre. *Matrices. Theory and Applications*, 2nd edition. Graduate texts in Mathematics **216**, Springer-Verlag (2010).
- [9] D. Serre. Divergence-free positive symmetric tensors and fluid dynamics. *Annales de l’Institut Henri Poincaré (analyse non linéaire)*, **35** (2018), pp 1209–1234.
- [10] D. Serre. Compensated integrability. Applications to the Vlasov–Poisson equation and other models of mathematical physics. *J. Maths. Pures & Appl.*, to appear (2019).

- [11] D. Serre. Divergence et déterminant des tenseurs symétriques positifs. *Séminaire Laurent Schwartz*, École Polytechnique (2018–19), Exposé numéro V.
- [12] K. Uchiyama. On the Boltzmann-Grad limit for the Broadwell model of the Boltzmann equation. *J. Stat. Phys.*, **52** (1988), pp 331–355.
- [13] L. N. Vaserstein. On systems of particles with finite-range and/or repulsive interaction. *Commun. Math. Phys.*, **69** (1979), pp 31–56.